

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Cazitel Plus XL Tablets For Dogs

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains:

Active substances:

Praziquantel	175 mg
Pyrantel Embonate	504 mg (equivalent to 175 mg pyrantel)
Febantel	525 mg

Excipients:

Qualitative composition of excipients and other constituents
Lactose monohydrate,
Microcrystalline cellulose,
Magnesium stearate,
Colloidal anhydrous silica,
Croscarmellose sodium,
Sodium laurilsulfate
Pork flavour

A yellow coloured oblong tablet with a breakline on both sides
The tablets can be divided into two equal parts.

3. CLINICAL INFORMATION

3.1 Target species

Dogs.

3.2 Indications for use for each target species

In adult dogs: Treatment of mixed infections by nematodes and cestodes of the following species

Nematodes:

Ascarids: *Toxocara canis*, *Toxascaris leonina* (adult and late immature forms).

Hookworms: *Uncinaria stenocephala*, *Ancylostoma caninum* (adults).

Whipworms: *Trichuris vulpis* (adults).

Cestodes:

Tapeworms: *Echinococcus* species, (*E. granulosus*, *E. multilocularis*), *Taenia* species, (*T. hydatigena*, *T. pisiformis*, *T. taeniformis*), *Dipylidium caninum* (adult and immature forms).

3.3 Contraindications

Do not use simultaneously with piperazine compounds.

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

3.4 Special warnings

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*.

Tapeworm infestation is certain to reoccur unless control of intermediate hosts such as fleas, mice, etc. is undertaken.

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.
- Underdosing, which may be due to underestimation of body weight, misadministration of the product.

3.5 Special precautions for use

Special precautions for safe use in the target species:

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

In the interests of good hygiene, persons administering the tablets directly to the dog, or by adding them to the dog's food, should wash their hands afterwards

Special precautions for the protection of the environment:

Not applicable.

Other precautions:

Echinococcosis represents a hazard for humans. As echinococcosis is a notifiable disease to the World Organisation for Animal Health (OIE), specific guidelines on the treatment and follow-up, and on the safeguard of persons, need to be obtained from the relevant competent authority.

3.6 Adverse events

Dogs:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Digestive tract disorders (diarrhoea, emesis) Lethargy, anorexia, hyperactivity
--------------------------------------------------------------------------------	------------------------------------------------------------------------------------

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See also the last section of the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Pregnancy:

Teratogenic effects attributed to high doses of febantel have been reported in sheep and rats. No studies have been performed in dogs during early pregnancy. Use of the product during

pregnancy should be in accordance with a benefit risk assessment by the responsible veterinarian. The use is not recommended during the first 4 weeks of pregnancy in dogs. Do not exceed the stated dose when treating pregnant bitches.

3.8 Interaction with other medicinal products and other forms of interaction

Do not use simultaneously with piperazine compounds as the anthelmintic effects of pyrantel and piperazine may be antagonised.

Concurrent use with other cholinergic compounds can lead to toxicity.

3.9 Administration routes and dosage

Oral use.

The recommended dose rates are: 15 mg/kg bodyweight febantel, 5 mg/kg pyrantel (equivalent to 14.4 mg/kg pyrantel embonate) and 5 mg/kg praziquantel. This is equivalent to 1 Cazitel Plus XL tablet per 35 kg. bodyweight.

Dogs of > 35 kg. bodyweight should be given 1 Cazitel Plus XL tablet plus the appropriate quantity of Cazitel Plus tablets equivalent to 1 tablet per 10 kg. bodyweight.

Dogs weighing approximately 17.5 kg. bodyweight should be given ½ Cazitel Plus XL tablet

The tablets can be given directly to the dog or disguised in food. No starvation is needed before or after treatment.

The tablets can be divided into two equal parts.

If there is a risk for re-infestation, the advice of a veterinarian should be sought regarding the need for and the frequency of repeat administration.

To ensure a correct dosage, body weight should be determined as accurately as possible.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

The combination of praziquantel, pyrantel embonate and febantel is well tolerated in dogs. In safety studies, a single dose of 5 times the recommended dose or greater gave rise to occasional vomiting.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable

3.12 Withdrawal periods

Not applicable.

4. PHARMACOLOGICAL INFORMATION

4.1 ATCvet code:

QP52AA51

4.2 Pharmacodynamics

This veterinary medicinal product contains anthelmintics active against gastrointestinal roundworms and tapeworms. The veterinary medicinal product contains three active substances, as follows:

1. Febantel, a probenzimidazole
2. Pyrantel embonate (pamoate), a tetrahydropyrimidine derivative
3. Praziquantel, a partially hydrogenated pyrazinoisoquinoline derivative

In this fixed combination, pyrantel and febantel act against all relevant nematodes (ascarids, hookworms, and whipworms) in dogs. In particular, the activity spectrum covers *Toxocara canis*, *Toxascaris leonina*, *Uncinaria stenocephala*, *Ancylostoma caninum* and *Trichuris vulpis*.

This combination shows synergistic activity in the case of hookworms and febantel is effective against *T. vulpis*.

The spectrum of activity of praziquantel covers all important cestode species in dogs, in particular *Taenia* spp., *Dipylidium caninum*, *Echinococcus granulosus* and *Echinococcus multilocularis*.

Praziquantel acts against all adult and immature forms of these parasites.

Praziquantel is very rapidly absorbed through the parasite's surface and distributed throughout the parasite. Both in vitro and in vivo studies have shown that praziquantel causes severe damage to the parasite integument, resulting in the contraction and paralysis of the parasites. There is an almost instantaneous tetanic contraction of the parasite musculature and a rapid vacuolisation of the syncytial tegument. This rapid contraction has been explained by changes in divalent cation fluxes, especially calcium.

Pyrantel acts as a cholinergic agonist. Its mode of action is to stimulate nicotinic cholinergic receptors of the parasite, induce spastic paralysis of the nematodes and thereby allow removal from the gastrointestinal system by peristalsis.

Within the mammalian system, febantel undergoes ring closure, forming fenbendazole and oxfendazole. It is these chemical entities which exert the anthelmintic effect by inhibition of tubulin polymerisation. Formation of microtubules is thereby prevented, resulting in disruption of structures vital to the normal functioning of the helminth. Glucose uptake in particular is affected, leading to a depletion in cell ATP. The parasite dies upon exhaustion of its energy reserves, which occurs 2 – 3 days later

4.3 Pharmacokinetics

Perorally administered praziquantel is absorbed almost completely from the intestinal tract. After absorption, the drug is distributed to all organs. Praziquantel is metabolised into inactive forms in the liver and secreted in bile. It is excreted within 24 hours to more than 95% of the administered dosage. Only traces of non-metabolised praziquantel are excreted.

Following administration of the veterinary medicinal product to dogs, peak plasma concentrations of praziquantel were achieved by approximately 2.5 hours.

The pamoate salt of pyrantel has low aqueous solubility, an attribute that reduces absorption from the gut and allows the drug to reach and be effective against parasites in the large intestine.

Following absorption, pyrantel pamoate is quickly and almost completely metabolised into inactive metabolites that are excreted rapidly in the urine.

Febantel is absorbed relatively rapidly and metabolized to a number of metabolites including fenbendazole and oxfendazole, which have anthelmintic activity.

Following administration of the veterinary medicinal product to dogs, peak plasma concentrations of fenbendazole and oxfendazole were achieved by approximately 7-9 hours.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

Not applicable.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 5 years

Shelf-life of half tablets: 14 days .

5.3 Special precautions for storage

This veterinary medicinal product does not require any special temperature storage conditions. Each time an unused half tablet is stored, it should be returned to the open blister space and the blister inserted back into the outer carton.

5.4 Nature and composition of immediate packaging

The product is presented in:

Blister packs made up of PVC/PE/PCTFE with 20 μ hard tempered aluminium foil with 2, 4, 5, 6, 8, 10, 12, 14, 16, 18 or 20 tablets per blister.

The Blisters are packed into cartons containing either 2, 4, 5, 6, 8, 10, 12, 14, 16, 18, 20, 24, 28, 30, 32, 36, 40, 42, 44, 48, 50, 52, 56, 60, 64, 68, 70, 72, 76, 80, 84, 88, 92, 96, 98, 100, 104, 106, 108, 112, 116, 120, 140, 150, 180, 200, 204, 206, 208, 250, 280, 300, 500 or 1000 tablets. Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

Chanelle Pharmaceuticals Manufacturing Limited.

7. MARKETING AUTHORISATION NUMBER(S)

VPA10987/077/002

8. DATE OF FIRST AUTHORISATION

02/03/2012

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

18/07/2025

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product not subject to prescription.

Detailed information on this veterinary medicinal product is available in the Union Product Database.