

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Daxocox 15 mg tablets for dogs
Daxocox 30 mg tablets for dogs
Daxocox 45 mg tablets for dogs
Daxocox 70 mg tablets for dogs
Daxocox 100 mg tablets for dogs
Daxocox 140 mg tablets for dogs
Daxocox 200 mg tablets for dogs

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains:

Active substances:

Enflicoxib 15 mg
Enflicoxib 30 mg
Enflicoxib 45 mg
Enflicoxib 70 mg
Enflicoxib 100 mg
Enflicoxib 140 mg
Enflicoxib 200 mg

Excipients:

Qualitative composition of excipients and other constituents	Quantitative composition if that information is essential for proper administration of the veterinary medicinal product
Mannitol	
Silicified microcrystalline cellulose	
Sodium laurilsulfate	
Crospovidone	
Copovidone	
Sodium stearyl fumarate	
Talc	
Iron oxide black (E172)	0.26%
Iron oxide yellow (E172)	0.45%
Iron oxide red (E172)	0.50%
Microcrystalline cellulose	
Dried flavour	

Brown, round and convex or capsule shaped tablets.

3. CLINICAL INFORMATION

3.1 Target species

Dogs

3.2 Indications for use for each target species

For the treatment of pain and inflammation associated with osteoarthritis (or degenerative joint disease).

For the treatment of pain and inflammation associated with orthopaedic or soft tissue surgery.

3.3 Contraindications

Do not use in animals suffering from gastrointestinal disorders, protein or blood losing enteropathy or haemorrhagic disorders.

Do not use in cases of impaired renal or hepatic function.

Do not use in cases of cardiac insufficiency.

Do not use in pregnant or lactating dogs.

Do not use in animals intended for breeding purposes.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in cases of known hypersensitivity to sulphonamides.

Do not use in any dehydrated, hypovolemic or hypotensive animal, as there is a potential risk of increased renal toxicity.

3.4 Special warnings

Do not administer other Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) or glucocorticoids concurrently or within 2 weeks of the last administration of this veterinary medicinal product.

3.5 Special precautions for use

Special precautions for safe use in the target species:

Since the safety of the medicinal product has not been fully demonstrated in very young animals, careful monitoring is advised during the treatment of young dogs aged less than 6 months.

The active metabolite of enflcoxib exhibits an extended plasma half-life due to its low rate of elimination. Use this veterinary medicinal product under strict veterinary monitoring where there is a risk of gastrointestinal ulceration, or if the animal previously displayed intolerance to NSAIDs.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product can cause hypersensitivity (allergic) reactions. People with known hypersensitivity to NSAIDs should avoid contact with the veterinary medicinal product.

Some NSAIDs may be harmful for the unborn child, especially during the third trimester of pregnancy. Pregnant women should administer this veterinary medicinal product with care.

Ingestion of this veterinary medicinal product may be harmful, especially for children, and prolonged pharmacological effects leading to e.g. gastrointestinal disorders may be observed. To avoid accidental ingestion, administer the tablet to the dog immediately after removal from the blister packaging and do not split or crush tablets.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Special precautions for the protection of the environment:

Not applicable.

3.6 Adverse events

Dogs:

Common (1 to 10 animals / 100 animals treated):	Vomiting ⁽¹⁾ , Diarrhoea ⁽¹⁾ , Soft stool ⁽¹⁾
Uncommon (1 to 10 animals / 1 000 animals treated):	Apathy, Appetite loss Haemorrhagic diarrhoea, Gastric ulcer
Undetermined frequency (cannot be estimated from the available data)	Elevated blood urea nitrogen (BUN), Elevated cholesterol (total)

⁽¹⁾ Most cases recover without treatment.

In case of adverse reactions, the use of the veterinary medicinal product should be stopped and general supportive therapy, as for clinical overdose with NSAIDs, should be applied until complete resolution of the signs. Particular attention should be paid to maintain haemodynamic status.

Gastrointestinal protectants and parenteral fluids, as appropriate, may be required for animals that experience gastrointestinal or renal adverse reactions.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

The safety of the veterinary medicinal product has not been established during pregnancy, lactation or reproduction in the target species.

Pregnancy and lactation:

Do not use during pregnancy or lactation.

Laboratory studies in rats and rabbits have shown evidence of foetotoxic effects at maternally toxic doses.

Fertility:

Do not use in breeding animals.

3.8 Interaction with other medicinal products and other forms of interaction

No drug-interaction studies have been performed. In common with other NSAIDs, this veterinary medicinal product should not be administered simultaneously with other NSAIDs or glucocorticoids.

Animals should be carefully monitored if this veterinary medicinal product is administered simultaneously with an anticoagulant.

Enflicoxib is highly bound to plasma proteins and may compete with other highly bound substances, such that concomitant administration may result in toxic effects.

Pre-treatment with other anti-inflammatory substances may result in additional or increased adverse reactions. To avoid such adverse reactions when this veterinary medicinal product is to be administered in replacement to another NSAID, ensure an appropriate treatment-free period before

administering the first dose. The treatment-free period should, however, consider the pharmacology of the medicinal products previously used.

Concurrent administration of potentially nephrotoxic veterinary medicinal products should be avoided.

3.9 Administration routes and dosage

Oral use.

Dosing interval is ONCE PER WEEK.

Osteoarthritis:

First dose: 8 mg enflcoxib per kg body weight.

Maintenance dose: repeat the treatment every 7 days at the dose of 4 mg enflcoxib per kg body weight.

Body weight (Kg) /Tablet size (mg)	Number of tablets to be administered													
	FIRST DOSE 8 mg/kg							MAINTENANCE DOSE 4 mg/kg						
	15 mg	30 mg	45 mg	70 mg	100 mg	140 mg	200 mg	15 mg	30 mg	45 mg	70 mg	100 mg	140 mg	200 mg
2.5 - 4.9	2							1						
5 - 7.5		2							1					
7.6 – 11.2			2							1				
11.3 - 17.5				2							1			
17.6 - 25					2							1		
25.1 - 35						2							1	
35.1 - 50							2							1
50.1 - 75						4							2	

For perioperative use:

A single treatment at a dose of 8 mg per kg body weight must be administered one day (at least 24 hours) before surgery is scheduled. If, at 7 days after the initial treatment (6 days post-surgery), the treating veterinarian determines that further post-operative analgesia is necessary, subsequent treatments may be administered at a dose of 4 mg per kg body weight and at a 7-day treatment interval.

The veterinary medicinal product should be given immediately before or with the dog's meal.

To ensure a correct dosage, body weight should be determined as accurately as possible.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

In overdose safety studies at a continuous weekly administration at 12 mg/kg body weight for a period of 7 months and at 20 mg/kg body weight for a period of 3 months, with an initial loading dose, there was evidence of elevated blood urea and serum cholesterol levels. No other associated treatment related effects were detected.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Not applicable.

4. PHARMACOLOGICAL INFORMATION

4.1 ATCvet code: QM01AH95

4.2 Pharmacodynamics

Enflicoxib is a non-steroidal anti-inflammatory drug belonging to the coxib class and acting by selective inhibition of the enzyme cyclooxygenase 2. The cyclooxygenase enzyme (COX) is present in two isoforms. COX-1 is usually a constitutive enzyme expressed in tissues, which synthesize products responsible for normal physiologic functions (e.g. in the gastro-intestinal tract and kidneys), and COX-2 is mainly inducible and synthesized by macrophages and other inflammatory cells after stimulation by cytokines and other mediators of inflammation. COX-2 is involved in the production of mediators, including PGE₂, that induce pain, exudation, inflammation and fever.

4.3 Pharmacokinetics

Enflicoxib is well absorbed after oral administration; bioavailability is high, and it is increased by 40-50% with food. The recommended dose is based on administration with food. After oral administration to fed dogs at the recommended loading dose of 8 mg/kg bw, enflicoxib is readily absorbed and reaches its maximal concentration of 1.8 (\pm 0.4) mcg/ml (C_{max}) after 2 hours (T_{max}). The elimination half-life ($t_{1/2}$) is 20 h.

Enflicoxib is extensively transformed by the hepatic microsomal system into an active pyrazol metabolite, which reaches its maximal concentration of 1.3 (\pm 0.2) mcg/ml (C_{max}) after 6 days (T_{max}). The elimination half-life ($t_{1/2}$) is 17 days.

Enflicoxib and its active metabolite are extensively bound to dog plasma proteins (98-99%) and are mainly excreted in faeces by the biliary route and, to a lesser extent, in urine.

After repeated administrations, systemic exposure to enflicoxib and its pyrazol metabolite rapidly reaches a plateau, with no evidence of time-dependent pharmacokinetics or over-accumulation for either compound.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

Not applicable.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 5 years.

5.3 Special precautions for storage

This veterinary medicinal product does not require any special temperature storage conditions.

Keep the blisters in the outer carton in order to protect from light.

In order to avoid any accidental ingestion, store tablets out of reach of animals.

5.4 Nature and composition of immediate packaging

Blisters are made of a PVC/Aluminium/oriented polyamide blister foil and an aluminium lidding foil.

Package sizes:

Carton boxes containing 4, 5, 10, 12, 20, 24, 50 or 100 tablets for Daxocox 15, 30, 45, 70 and 100 mg.

Carton boxes containing 4, 5, 12 or 20 tablets for Daxocox 140 and 200 mg.

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

Ecuphar NV

7. MARKETING AUTHORISATION NUMBER(S)

EU/2/21/270/001-048

8. DATE OF FIRST AUTHORISATION

Date of first authorisation: 20/04/2021.

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

{MM/YYYY}

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

ANNEX II

OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

None.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

CARTON BOX

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Daxocox 15 mg tablets
Daxocox 30 mg tablets
Daxocox 45 mg tablets
Daxocox 70 mg tablets
Daxocox 100 mg tablets
Daxocox 140 mg tablets
Daxocox 200 mg tablets

2. STATEMENT OF ACTIVE SUBSTANCES

Each tablet contains:

Enflicoxib 15 mg
Enflicoxib 30 mg
Enflicoxib 45 mg
Enflicoxib 70 mg
Enflicoxib 100 mg
Enflicoxib 140 mg
Enflicoxib 200 mg

3. PACKAGE SIZE

4 tablets
5 tablets
10 tablets
12 tablets
20 tablets
24 tablets
50 tablets
100 tablets

4. TARGET SPECIES

Dogs

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Oral use.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {mm/yyyy}

9. SPECIAL STORAGE PRECAUTIONS

Keep the blisters in the outer carton in order to protect from light.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER**14. MARKETING AUTHORISATION NUMBERS**

EU/2/21/270/001 (15 mg, 4 tablets)
EU/2/21/270/044 (15 mg, 5 tablets)
EU/2/21/270/002 (15 mg, 10 tablets)
EU/2/21/270/003 (15 mg, 12 tablets)
EU/2/21/270/004 (15 mg, 20 tablets)
EU/2/21/270/005 (15 mg, 24 tablets)
EU/2/21/270/006 (15 mg, 50 tablets)
EU/2/21/270/007 (15 mg, 100 tablets)
EU/2/21/270/008 (30 mg, 4 tablets)
EU/2/21/270/045 (30 mg, 5 tablets)
EU/2/21/270/009 (30 mg, 10 tablets)
EU/2/21/270/010 (30 mg, 12 tablets)
EU/2/21/270/011 (30 mg, 20 tablets)
EU/2/21/270/012 (30 mg, 24 tablets)
EU/2/21/270/013 (30 mg, 50 tablets)
EU/2/21/270/014 (30 mg, 100 tablets)
EU/2/21/270/015 (45 mg, 4 tablets)
EU/2/21/270/046 (45 mg, 5 tablets)
EU/2/21/270/016 (45 mg, 10 tablets)

EU/2/21/270/017 (45 mg, 12 tablets)
EU/2/21/270/018 (45 mg, 20 tablets)
EU/2/21/270/019 (45 mg, 24 tablets)
EU/2/21/270/020 (45 mg, 50 tablets)
EU/2/21/270/021 (45 mg, 100 tablets)
EU/2/21/270/022 (70 mg, 4 tablets)
EU/2/21/270/047 (70 mg, 5 tablets)
EU/2/21/270/023 (70 mg, 10 tablets)
EU/2/21/270/024 (70 mg, 12 tablets)
EU/2/21/270/025 (70 mg, 20 tablets)
EU/2/21/270/026 (70 mg, 24 tablets)
EU/2/21/270/027 (70 mg, 50 tablets)
EU/2/21/270/028 (70 mg, 100 tablets)
EU/2/21/270/029 (100 mg, 4 tablets)
EU/2/21/270/048 (100 mg, 5 tablets)
EU/2/21/270/030 (100 mg, 10 tablets)
EU/2/21/270/031 (100 mg, 12 tablets)
EU/2/21/270/032 (100 mg, 20 tablets)
EU/2/21/270/033 (100 mg, 24 tablets)
EU/2/21/270/034 (100 mg, 50 tablets)
EU/2/21/270/035 (100 mg, 100 tablets)
EU/2/21/270/036 (140 mg, 4 tablets)
EU/2/21/270/037 (140 mg, 5 tablets)
EU/2/21/270/038 (140 mg, 12 tablets)
EU/2/21/270/039 (140 mg, 20 tablets)
EU/2/21/270/040 (200 mg, 4 tablets)
EU/2/21/270/041 (200 mg, 5 tablets)
EU/2/21/270/042 (200 mg, 12 tablets)
EU/2/21/270/043 (200 mg, 20 tablets)

15. BATCH NUMBER

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS
BLISTER

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Daxocox



2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

15 mg
30 mg
45 mg
70 mg
100 mg
140 mg
200 mg

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Daxocox 15 mg tablets for dogs
Daxocox 30 mg tablets for dogs
Daxocox 45 mg tablets for dogs
Daxocox 70 mg tablets for dogs
Daxocox 100 mg tablets for dogs
Daxocox 140 mg tablets for dogs
Daxocox 200 mg tablets for dogs

2. Composition

Each tablet contains:

Active substances:

Enflicoxib	15 mg
Enflicoxib	30 mg
Enflicoxib	45 mg
Enflicoxib	70 mg
Enflicoxib	100 mg
Enflicoxib	140 mg
Enflicoxib	200 mg

Excipients:

Iron oxide black (E172)	0.26%
Iron oxide yellow (E172)	0.45%
Iron oxide red (E172)	0.50%

Brown, round and convex or capsule shaped tablets.

3. Target species

Dogs



4. Indications for use

For the treatment of pain and inflammation associated with osteoarthritis (or degenerative joint disease).

For the treatment of pain and inflammation associated with orthopaedic or soft tissue surgery.

5. Contraindications

Do not use in animals suffering from gastrointestinal disorders, protein or blood losing enteropathy or haemorrhagic disorders.

Do not use in cases of impaired renal or hepatic function.

Do not use in cases of cardiac insufficiency.

Do not use in pregnant or lactating dogs.

Do not use in animals intended for breeding purposes.
Do not use in cases of hypersensitivity to the active substance or to any of the excipients.
Do not use in cases of known hypersensitivity to sulphonamides.
Do not use in any dehydrated, hypovolemic or hypotensive animal, as there is a potential risk of increased renal toxicity.

6. Special warnings

Special warnings:

Do not administer other Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) or glucocorticoids concurrently or within 2 weeks of the last administration of this veterinary medicinal product.

Special precautions for safe use in the target species:

Since the safety of the medicinal product has not been fully demonstrated in very young animals, careful monitoring is advised during the treatment of young dogs aged less than 6 months.

The active metabolite of enflcoxib exhibits an extended plasma half-life due to its low rate of elimination. Use this veterinary medicinal product under strict veterinary monitoring where there is a risk of gastrointestinal ulceration, or if the animal previously displayed intolerance to NSAIDs.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product can cause hypersensitivity (allergic) reactions. People with known hypersensitivity to NSAIDs should avoid contact with the veterinary medicinal product.

Some NSAIDs may be harmful for the unborn child, especially during the third trimester of pregnancy. Pregnant women should administer this veterinary medicinal product with care.

Ingestion of this veterinary medicinal product may be harmful, especially for children, and prolonged pharmacological effects leading to e.g. gastrointestinal disorders may be observed. To avoid accidental ingestion, administer the tablet to the dog immediately after removal from the blister packaging and do not split or crush tablets.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Pregnancy, lactation and fertility:

The safety of the veterinary medicinal product has not been established during pregnancy, lactation or reproduction in the target species.

Pregnancy and lactation:

Do not use during pregnancy or lactation.

Laboratory studies in rats and rabbits have shown evidence of foetotoxic effects at maternally toxic doses.

Fertility:

Do not use in breeding animals.

Interaction with other medicinal products and other forms of interaction:

No drug-interaction studies have been performed. In common with other NSAIDs, this veterinary medicinal product should not be administered simultaneously with other NSAIDs or glucocorticoids.

Animals should be carefully monitored if this veterinary medicinal product is administered simultaneously with an anticoagulant.

Enflicoxib is highly bound to plasma proteins and may compete with other highly bound substances, such that concomitant administration may result in toxic effects.

Pre-treatment with other anti-inflammatory substances may result in additional or increased adverse reactions. To avoid such adverse reactions when this veterinary medicinal product is to be administered in replacement to another NSAID, ensure an appropriate treatment-free period before administering the first dose. The treatment-free period should, however, consider the pharmacology of the medicinal products previously used.

Concurrent administration of potentially nephrotoxic veterinary medicinal products should be avoided.

Overdose:

In overdose safety studies at a continuous weekly administration at 12 mg/kg body weight for a period of 7 months and at 20 mg/kg body weight for a period of 3 months, with an initial loading dose, there was evidence of elevated blood urea and serum cholesterol levels. No other associated treatment related effects were detected.

7. Adverse events

Dogs:

Common (1 to 10 animals / 100 animals treated):	Vomiting ⁽¹⁾ , Diarrhoea ⁽¹⁾ , Soft stool ⁽¹⁾
Uncommon (1 to 10 animals / 1 000 animals treated):	Apathy, Appetite loss Haemorrhagic diarrhoea, Gastric ulcer
Undetermined frequency (cannot be estimated from the available data)	Elevated blood urea nitrogen (BUN), Elevated cholesterol (total)

⁽¹⁾ Most cases recover without treatment.

In case of adverse reactions, the use of the veterinary medicinal product should be stopped and general supportive therapy, as for clinical overdose with NSAIDs, should be applied until complete resolution of the signs. Particular attention should be paid to maintain haemodynamic status.

Gastrointestinal protectants and parenteral fluids, as appropriate, may be required for animals that experience gastrointestinal or renal adverse reactions.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or its local representative using the contact details at the end of this leaflet, or via your national reporting system: {national system details}.

8. Dosage for each species, routes and method of administration

Oral use.

Dosing interval is ONCE PER WEEK.

Osteoarthritis:

First dose: 8 mg enflcoxib per kg body weight.

Maintenance dose: repeat the treatment every 7 days at the dose of 4 mg enflcoxib per kg body weight.

Body weight (Kg) /Tablet size (mg)	Number of tablets to be administered													
	FIRST DOSE							MAINTENANCE DOSE						
	8 mg/kg							4 mg/kg						
	15 mg	30 mg	45 mg	70 mg	100 mg	140 mg	200 mg	15 mg	30 mg	45 mg	70 mg	100 mg	140 mg	200 mg
2.5 - 4.9	2							1						
5 - 7.5		2							1					
7.6 – 11.2			2							1				
11.3 - 17.5				2							1			
17.6 - 25					2							1		
25.1 - 35						2							1	
35.1 - 50							2							1
50.1 - 75						4							2	

For perioperative use:

A single treatment at a dose of 8 mg per kg body weight must be administered one day (at least 24 hours) before surgery is scheduled. If, at 7 days after the initial treatment (6 days post-surgery), the treating veterinarian determines that further post-operative analgesia is necessary, subsequent treatments may be administered at a dose of 4 mg per kg body weight and at a 7-day treatment interval.

To ensure a correct dosage, body weight should be determined as accurately as possible.

9. Advice on correct administration

The veterinary medicinal product should be given immediately before or with the dog’s meal.

10. Withdrawal periods

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special temperature storage conditions. Keep the blisters in the outer carton in order to protect from light.

In order to avoid any accidental ingestion, store tablets out of reach of animals.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and blister after Exp. The expiry date refers to the last day of that month.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack sizes

EU/2/21/270/001-048

Carton boxes containing 4, 5, 10, 12, 20, 24, 50 or 100 tablets for Daxocox 15, 30, 45, 70 and 100 mg.
Carton boxes containing 4, 5, 12 and 20 tablets for Daxocox 140 and 200 mg.

Not all pack sizes may be marketed.

15. Date on which the package leaflet was last revised

{MM/YYYY}

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

16. Contact details

Marketing authorisation holder:

Ecuphar NV
Legeweg 157-i
B-8020 Oostkamp, Belgium

Manufacturer responsible for batch release:

Lelypharma B.V.
Zuiveringweg 42
8243 PZ
Lelystad
The Netherlands

Local representatives and contact details to report suspected adverse events:

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

België/Belgique/Belgien

Ecuphar NV
Legeweg 157-i
B-8020 Oostkamp
Tél/Tel: +32 50314269
E-mail: info@ecuphar.be

Република България

САМ БС ЕООД
Бул. "Д-р Петър Дертлиев" 25, Търговски
център Лабиринт, ет. 5, офис САМ БС ЕООД
BG София 1335
Тел: +359 2 810 0173
E-mail: sambs@sambs.bg

Česká republika

VIRBAC Czech Republic s.r.o.
Žitavského 496
156 00 Praha 5
Česká republika
Tel.: +420 608 836 529

Danmark

VIRBAC Danmark A/S
Profilvej 1
DK-6000 Kolding
Tlf: +45 75521244
E-mail: virbac@virbac.dk

Deutschland

Ecuphar GmbH
Brandteichstraße 20
DE-17489 Greifswald
Tel: +49 3834835840
E-mail: info@ecuphar.de

Eesti

OÜ ZOOVETVARU
Uusaru 5
EE-76505 Saue/Harjumaa
Eesti
Tel: +372 56480207
E-mail: pv@zoovet.eu

Ελλάδα

ΧΕΛΛΑΦΑΡΜ VET ΑΕ
1° χλμ. Λ Παιανίας-Μαρκοπούλου,
ΤΘ 100, 19002, Παιανία, Ελλάδα
Τηλ.: +30 2106895188, +30 2114041436
E-mail: info@hellafarmvet.gr

Lietuva

OÜ ZOOVETVARU
Uusaru 5
EE-76505 Saue/Harjumaa
Estija
Tel: +372 56480207
E-mail: pv@zoovet.eu

Luxembourg/Luxemburg

Ecuphar NV
Legeweg 157-i
B-8020 Oostkamp
Tél/Tel: +32 50314269
E-mail: info@ecuphar.be

Magyarország

VIRBAC HUNGARY KFT
Dózsa György út 84. B épület
HU-1068 Budapest
Tel.: +36703387177
E-mail: akos.csoman@virbac.hu

Malta

AGRIMED LIMITED
Mdina Road,
Żebbuġ ZBG 9016
Tel: +356 21465797
E-mail: info@agrimedltd.com

Nederland

Ecuphar BV
Verlengde Poolseweg 16
NL-4818 CL Breda
Tel: +31 880033800
E-mail: info@ecuphar.nl

Norge

VIRBAC Danmark A/S
Profilvej 1
DK-6000 Kolding
Tlf: +45 75521244
E-mail: virbac@virbac.dk

Österreich

VIRBAC Österreich GmbH
Hildebrandgasse 27
AT-1180 Wien
Tel: +43-(0)121834260

España

Ecuphar Veterinaria S.L.U.
C/Cerdanya, 10-12 Planta 6º
ES- 08173 Sant Cugat del Vallés,
Barcelona
Tel: +34 935955000
E-mail: info@ecuphar.es

France

VIRBAC France
13^e rue LID
FR-06517, Carros
Tél: 0 800 73 09 10

Hrvatska

CENTRALNA VETERINARSKA AGENCIJA
d.o.o. (CVA)
Prve Ravnice 2e, 10000 Zagreb
Republika Hrvatska
Tel: + 385 91 46 55 112
E-mail: cva@cva.hr

Ireland

Duggan Veterinary Supplies,
Unit 9 Thurles Retail Park,
Thurles,
Co. Tipperary,
E41 E7K7.
Tel: +353 (0)504 43169
E-mail: pv@dugganvet.ie

Ísland

Ecuphar NV
Legeweg 157-i
B-8020 Oostkamp
Sími: +32 50314269
Netfang: info@ecuphar.be

Italia

Ecuphar Italia S.r.l.
Viale Francesco Restelli, 3/7
IT-20124 Milano
Tel: +39 0282950604
E-mail: info@ecuphar.it

Κύπρος

Panchris Feeds (Veterinary) Ltd
Γόρδιου Δεσμού 15, Βιομηχανική περιοχή
Αραδίππου, Λάρνακα, 7100, (Τ.Θ. 45189, 7112,
Αραδίππου), Κύπρος.
Τηλ.: +357 24813333
E-mail: pharma.safety@panchris.com

Polska

VIRBAC Sp. z o.o.
ul. Puławska 314
PL 02-819 Warszawa
Tel.: +48 22 855 40 46

Portugal

Belphar Lda
Sintra Business Park 7, Edifício 1, Escritório 2K
Zona Industrial de Abrunheira
PT-2710-089 Sintra
Tel: +351 308808321
E-mail: info@ecuphar.pt

România

Altius SA
Str. Zăgazului nr. 21-25, Corp A, et 8 si 8A
Ap. A.8.2, sect 1, cod 014261, București,
Romania
Tel: +40 21 310 88 80

Slovenija

MEDICAL INTERTRADE d.o.o.
Brodišče 12, 1236 Trzin
Slovenija
Tel: +386 1 2529 113
E-mail: farmakovigilanca@medical-intertrade.si

Slovenská republika

VIRBAC Czech Republic s.r.o.
Žitavského 496
156 00 Praha 5
Česká republika
Tel.: +420 608 836 529

Suomi/Finland

BIOFARM OY
Yrittäjätie 20
FI-03600 Karkkila
Puh/Tel: +358-9-225 2560
E-mail: haittavaikutukset@biofarm.fi

Sverige

VIRBAC Danmark A/S Filial Sverige
Box 1027
SE-171 21 Solna
Tel: +45 75521244
E-mail: virbac@virbac.dk

Latvija

OÜ ZOOVETVARU

Uusaru 5

EE-76505 Saue/Harjumaa

Igaunija

Tel: +372 56480207

E-mail: pv@zoovet.eu**United Kingdom (Northern Ireland)**

Ecuphar NV

Legeweg 157-i

B-8020 Oostkamp

Tel: +32 50314269

E-mail: info@ecuphar.be