

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

ERYSENG suspension for injection for pigs

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each dose of 2 ml contains:

Active substance:

Inactivated *Erysipelothrix rhusiopathiae*, strain R32E11,
* IE_{50%} – Inhibition ELISA 50%.

> 3.34 log₂ IE_{50%}*

Adjuvants:

Aluminium hydroxide
DEAE-Dextran
Ginseng

5.29 mg (aluminium)

Excipients:

Qualitative composition of excipients and other constituents
Disodium phosphate dodecahydrate
Potassium chloride
Potassium dihydrogen phosphate
Simethicone
Sodium chloride
Sodium hydroxide
Water for injections

Whitish suspension.

3. CLINICAL INFORMATION

3.1 Target species

Pigs.

3.2 Indications for use for each target species

For the active immunisation of male and female pigs to reduce clinical signs (skin lesions and fever) of swine erysipelas caused by *Erysipelothrix rhusiopathiae*, serotype 1 and serotype 2.

Onset of immunity: three weeks after completion of the basic vaccination scheme.

Duration of immunity: six months.

3.3 Contraindications

Do not use in cases of hypersensitivity to the active substance, to the adjuvants or to any of the excipients.

3.4 Special warnings

Vaccinate healthy animals only.

3.5 Special precautions for use

Special precautions for safe use in the target species:

None.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Special precautions for the protection of the environment:

Not applicable.

3.6 Adverse events

Pigs:

Very common (> 1 animal / 10 animals treated):	Injection site inflammation ¹
Common (1 to 10 animals / 100 animals treated):	Elevated temperature ²
Very rare (< 1 animal / 10 000 animals treated, including isolated reports):	Anaphylactic-type reaction ³

¹Mild to moderate inflammation at the injection site that typically resolves within 4 days but in some cases may persist for up to 12 days post-vaccination.

²A transient increase in body temperature within the first 6 hours after vaccination, which spontaneously resolves within 24 hours.

³An appropriate symptomatic treatment is recommended.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Pregnancy and lactation:

Can be used during pregnancy and lactation.

3.8 Interaction with other medicinal products and other forms of interaction

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

3.9 Administration routes and dosage

Intramuscular use.

Allow the vaccine to reach room temperature (15 °C – 25 °C) before administration.
Shake well before use.

Administer one dose of 2 ml by intramuscular injection in the neck muscles according to the following schedule:

Basic vaccination:

Pigs from 6 months of age which have not been previously vaccinated with the product should be given two injections with an interval of 3 – 4 weeks. The second injection should be administered 3 – 4 weeks before mating.

Revaccination:

A single injection should be given 2 – 3 weeks prior to each subsequent mating (approximately every 6 months).

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

No adverse reactions other than those mentioned in section 3.6 were observed after the administration of a 2-fold vaccine dose.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Zero days.

4. IMMUNOLOGICAL INFORMATION

4.1 ATCvet code: QI09AB03.

To stimulate the development of active immunity in pigs against *E. rhusiopathiae*.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

Do not mix with any other veterinary medicinal product.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

Shelf life after first opening the immediate packaging: use immediately.

5.3 Special precautions for storage

Store and transport refrigerated (2 °C – 8 °C).

Do not freeze.

Protect from light.

5.4 Nature and composition of immediate packaging

Type I colourless glass vials of 20, 50 and 100 ml. The vials are closed with a rubber stopper and aluminium cap.

Polyethylene (PET) bottles of 20, 50, 100 and 250 ml.

Pack sizes:

Cardboard box with 1 glass vial of 10 doses (20 ml).

Cardboard box with 1 glass vial of 25 doses (50 ml).

Cardboard box with 1 glass vial of 50 doses (100 ml).

Cardboard box with 1 PET bottle of 10 doses (20 ml).

Cardboard box with 1 PET bottle of 25 doses (50 ml).

Cardboard box with 1 PET bottle of 50 doses (100 ml).

Cardboard box with 1 PET bottle of 125 doses (250 ml).

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

LABORATORIOS HIPRA, S.A.

7. MARKETING AUTHORISATION NUMBER(S)

EU/2/14/166/001-007

8. DATE OF FIRST AUTHORISATION

Date of first authorisation: 04/07/2014

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

{DD/MM/YYYY}

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

ANNEX II

OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

None

ANNEX III

LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Cardboard box (20 ml, 50 ml, 100 ml, and 250 ml)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

ERYSENG suspension for injection

2. STATEMENT OF ACTIVE SUBSTANCES

One dose (2 ml) contains:

Inactivated *Erysipelothrix rhusiopathiae*, strain R32E11,

> 3.34 log₂ IE₅₀ %*

* IE₅₀ % – Inhibition ELISA 50 %.

3. PACKAGE SIZE

10 doses (20 ml)

25 doses (50 ml)

50 doses (100 ml)

125 doses (250 ml)

4. TARGET SPECIES

Pigs

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Intramuscular use.

7. WITHDRAWAL PERIODS

Withdrawal period: Zero days.

8. EXPIRY DATE

Exp. {mm/yyyy}

Once broached use immediately.

9. SPECIAL STORAGE PRECAUTIONS

Store and transport refrigerated.

Do not freeze.

Protect from light.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

LABORATORIOS HIPRA, S.A.

14. MARKETING AUTHORISATION NUMBERS

EU/2/14/166/001 10 doses
EU/2/14/166/002 25 doses
EU/2/14/166/003 50 doses
EU/2/14/166/004 10 doses
EU/2/14/166/005 25 doses
EU/2/14/166/006 50 doses
EU/2/14/166/007 125 doses

15. BATCH NUMBER

Lot {number}

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

Bottles (100 ml and 250 ml) and vials (100 ml)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

ERYSENG suspension for injection

2. STATEMENT OF ACTIVE SUBSTANCES

One dose (2 ml) contains:

Inactivated *Erysipelothrix rhusiopathiae*, strain R32E11,

> 3.34 log₂ IE₅₀ %*

* IE₅₀% – Inhibition ELISA 50 %.

3. TARGET SPECIES

Pigs

4. ROUTES OF ADMINISTRATION

Read the package leaflet before use.

Intramuscular use.

5. WITHDRAWAL PERIODS

Withdrawal period: Zero days.

6. EXPIRY DATE

Exp. {mm/yyyy}

Once broached use immediately.

7. SPECIAL STORAGE PRECAUTIONS

Store and transport refrigerated.

Do not freeze.

Protect from light.

8. NAME OF THE MARKETING AUTHORISATION HOLDER

LABORATORIOS HIPRA, S.A.

9. BATCH NUMBER

Lot {number}

10. PACKAGE SIZE

50 doses (100 ml)

125 doses (250 ml)

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

Bottles (20 ml, 50 ml) and vials (20 ml, 50 ml)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

ERYSENG

2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

Inactivated *Erysipelothrix rhusiopathiae*, strain R32E11,
* IE_{50%} – Inhibition ELISA 50%.

> 3.34 log₂ IE_{50%}*

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

Once broached use immediately.

5. PACKAGE SIZE

10 doses (20 ml)
25 doses (50 ml)

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

ERYSENG suspension for injection for pigs

2. Composition

Each dose of 2 ml contains:

Active substances:

Inactivated *Erysipelothrix rhusiopathiae*, strain R32E11,

> 3.34 log₂ IE_{50%} *

* IE_{50%} - Inhibition ELISA 50%.

Adjuvants:

Aluminium hydroxide

5.29 mg (aluminium)

Whitish suspension for injection.

3. Target species

Pigs.

4. Indications for use

For the active immunisation of male and female pigs to reduce clinical signs (skin lesions and fever) of swine erysipelas caused by *Erysipelothrix rhusiopathiae*, serotype 1 and serotype 2.

Onset of immunity: three weeks after completion of the basic vaccination scheme.

Duration of immunity: six months.

5. Contraindications

Do not use in cases of hypersensitivity to the active substance, to the adjuvants or to any of the excipients.

6. Special warnings

Special warnings:

Vaccinate healthy animals only.

Special precautions for safe use in the target species:

None.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Pregnancy and lactation:

Can be used during pregnancy and lactation.

Interaction with other medicinal products and other forms of interaction:

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Overdose:

No adverse reactions other than already mentioned under section “Adverse events” can be expected after the administration of a 2-fold vaccine dose.

Major incompatibilities:

Do not mix with any other veterinary medicinal product.

7. Adverse events

Pigs:

Very common (> 1 animal / 10 animals treated):
Injection site inflammation ¹
Common (1 to 10 animals / 100 animals treated):
Elevated temperature ²
Very rare (< 1 animal / 10 000 animals treated, including isolated reports):
Anaphylactic-type reaction (severe allergic reaction) ³

¹Mild to moderate inflammation at the injection site that typically resolves within 4 days but in some cases may persist for up to 12 days post-vaccination.

²A transient increase in body temperature within the first 6 hours after vaccination, which spontaneously resolves within 24 hours.

³An appropriate symptomatic treatment is recommended.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: [{national system details}](#).

8. Dosage for each species, routes and method of administration

Intramuscular use.

Administer one dose of 2 ml by intramuscular injection in the neck muscles according to the following schedule:

Basic vaccination:

Pigs from 6 months of age which have not been previously vaccinated with the product should be given two injections with an interval of 3 – 4 weeks. The second injection should be administered 3 – 4 weeks before mating.

Revaccination:

A single injection should be given 2 – 3 weeks prior to each subsequent mating (approximately every 6 months).

9. Advice on correct administration

Allow the vaccine to reach room temperature (15 °C – 25 °C) before administration.
Shake well before use.

10. Withdrawal periods

Zero days.

11. Special storage precautions

Keep out of the sight and reach of children.
Store and transport refrigerated (2 °C – 8 °C).
Do not freeze.
Protect from light.

Do not use this veterinary medicinal product after the expiry date which is stated on the label and the cardboard box after Exp. The expiry date refers to the last day of that month.
Shelf life after first opening the container: use immediately.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack sizes

Marketing authorisation numbers: EU/2/14/166/001 - 007

Pack sizes:

Cardboard box with 1 glass vial of 10 doses (20 ml).
Cardboard box with 1 glass vial of 25 doses (50 ml).
Cardboard box with 1 glass vial of 50 doses (100 ml).

Cardboard box with 1 PET bottle of 10 doses (20 ml).
Cardboard box with 1 PET bottle of 25 doses (50 ml).
Cardboard box with 1 PET bottle of 50 doses (100 ml).
Cardboard box with 1 PET bottle of 125 doses (250 ml).

Not all pack sizes may be marketed.

15. Date on which the package leaflet was last revised

{DD/MM/YYYY}

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

16. Contact details

Marketing authorisation holder and manufacturer responsible for batch release and contact details to report suspected adverse reactions:

LABORATORIOS HIPRA, S.A.
Avda. la Selva, 135
17170 AMER (Girona) SPAIN
TEL: +34 972 43 06 60

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

Local representatives and contact details to report suspected adverse reactions:

België/Belgique/Belgien

HIPRA BENELUX NV
Nieuwewandeling 62
9000 Gent
BELGIË
Tel: +32 09 2964464

Lietuva

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ISPANIJA
Tel: +34 972 43 06 60

Република България

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ИСПАНИЯ
Тел: +34 972 43 06 60

Luxembourg/Luxemburg

HIPRA BENELUX NV
Nieuwewandeling 62
9000 Gent
BELGIQUE
Tel: +32 09 2964464

Česká republika

HIPRA SLOVENSKO, s.r.o.
Zochova 5,
811 03 Bratislava,
SLOVENSKO
Tel: +421 02 32 335 223

Magyarország

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPANYOLORSZÁG
Tel: +34 972 43 06 60

Danmark

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPANIEN
Tel: +34 972 43 06 60

Malta

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPANJA
Tel: +34 972 43 06 60

Deutschland

HIPRA DEUTSCHLAND GmbH
Am Wehrhahn 28-30
40211 Düsseldorf
DEUTSCHLAND
Tel: +49 211 698236 – 0

Eesti

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
HISPAANIA
Tel: +34 972 43 06 60

Ελλάδα

HIPRA ΕΛΛΑΣ Α.Ε.
Λεωφ. Αθηνών 80 & Μηριόνου 2-4,
104 41 Κολωνός - ΑΘΗΝΑ - ΕΛΛΑΣ
Τηλ: +30 210 4978660

España

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ESPAÑA
Tel: +34 972 43 06 60

France

HIPRA FRANCE
7 rue Roland Garros, Batiment H
44700 - Orvault -
FRANCE
Tél: +33 02 51 80 77 91

Hrvatska

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ŠPANJOLSKA
Tel: +34 972 43 06 60

Ireland

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPAIN
Tel: +34 972 43 06 60

Nederland

HIPRA BENELUX NV
Nieuwewandeling 62
9000 Gent
BELGIË
Tel: +32 09 2964464

Norge

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPANIA
Tlf: +34 972 43 06 60

Österreich

HIPRA DEUTSCHLAND GmbH
Am Wehrhahn 28-30
40211 Düsseldorf
DEUTSCHLAND
Tel: +49 211 698236 – 0

Polska

HIPRA POLSKA Sp.z.o.o.
Ul. Wincentego Rzymowskiego 31
02-697 Warszawa - POLSKA
Tel: +48 22 642 33 06

Portugal

HIPRA ANIMAL HEALTH PORTUGAL,
PRODUTOS FARMACÊUTICOS, LDA
Portela de Mafra, Abrunheira
2665 – 191 Malveira - PORTUGAL
Tel:+351 219 663 450

România

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPANIA
Tel: +34 972 43 06 60

Slovenija

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ŠPANIIJA
Tel: +34 972 43 06 60

Ísland

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPÁNN
Sími: +34 972 43 06 60

Italia

Hipra Italia S.r.l.
Enrico Mattei, 2
25030 Coccaglio (BS)
ITALIA
Tel: +39 030 7241821

Κύπρος

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ΙΣΠΑΝΙΑ
Τηλ: +34 972 43 06 60

Latvija

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPĀNIJA
Tel. +34 972 43 06 60

Slovenská republika

HIPRA SLOVENSKO, s.r.o.
Zochova 5,
811 03 Bratislava,
SLOVENSKO
Tel: +421 02 32 335 223

Suomi/Finland

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ESPANJA
Puh/Tel: +34 972 43 06 60

Sverige

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPANIEN
Tel. +34 972 43 06 60

United Kingdom (Northern Ireland)

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPAIN
Tel: +34 972 43 06 60