

[Version 9.1,11/2024]

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

AXENTYL 200 mg/ml solution for injection for cattle, sheep, goats and pigs (AT, FR, IT)
TYLMASIN 200 mg/ml solution for injection for cattle, sheep, goats and pigs (BG, DE, EL, HU, PT, RO)
Tylovet 200 mg/ml solution for injection for cattle, sheep, goats and pigs (DK)
TYLOSIN BIOVET JSC 200 mg/ml solution for injection for cattle, sheep, goats and pigs (IE)
TYLOZYNA BIOVET JSC 200 mg/ml solution for injection for cattle, sheep, goats and pigs (PL)
TILOSINA BIOVET JSC 200 mg/ml solution for injection for cattle, sheep, goats and pigs (ES)

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml contains:

Active substance:

Tylosin 200 000 IU

Excipients:

Qualitative composition of excipients and other constituents	Quantitative composition if that information is essential for proper administration of the veterinary medicinal product
Benzyl alcohol (E1519)	40 mg
Propylene glycol	
Water for injections	

A pale yellow to amber-coloured liquid.

3. CLINICAL INFORMATION

3.1 Target species

Cattle, sheep, goats, pigs.

3.2 Indications for use for each target species

Infections caused by microorganisms susceptible to tylosin.

Cattle:

- Treatment of respiratory infections, metritis caused by Gram-positive micro-organisms, mastitis caused by *Streptococcus* spp., *Staphylococcus* spp. and interdigital necrobacillosis i.e. panaritum or foot rot

Cattle (calves):

- Treatment of respiratory infections and necrobacillosis.

Pigs:

- Treatment of enzootic pneumonia, haemorrhagic enteritis, erysipelas and metritis.
- Treatment of arthritis caused by *Mycoplasma* and *Staphylococcus* spp.

Sheep and goats:

Treatment of respiratory infections, metritis caused by Gram-positive microorganisms, mastitis caused by Gram-positive microorganisms or *Mycoplasma* spp.

3.3 Contraindications

Do not use in horses. Intramuscular injection can be fatal in chickens and turkeys.
Do not use in cases of hypersensitivity to tylosin, to other macrolides or to any of the excipients.

3.4 Special warnings

None.

3.5 Special precautions for use

Special precautions for safe use in the target species:

Due to likely variability (time, geographical) in susceptibility of bacteria to tylosin, bacteriological sampling and susceptibility testing are recommended.

Use of the veterinary medicinal product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to tylosin and may decrease the effectiveness of treatment with other macrolide antibiotics due to the potential for cross resistance. Official, national and regional antimicrobial policies should be taken into account when the veterinary medicinal product is used.

The efficacy data do not support the use of tylosin for the treatment of bovine mastitis caused by *Mycoplasma* spp. Haemorrhagic enteritis caused by *Brachyspira hyodysenteriae* must be treated with caution due to a high rate of in vitro resistance in European strains.

Where repeat injections are to be administered, use different sites for each injection.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Care should be taken to avoid accidental self-injection.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

Wash hands after use.

Tylosin may induce irritation. Macrolides, such as tylosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tylosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

Do not handle the veterinary medicinal product if you are allergic to ingredients in the veterinary medicinal product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice immediately and show the package leaflet or the label to the physician. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Special precautions for the protection of the environment:

Not applicable.

3.6 Adverse events

Cattle:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Injection site swelling, Injection site inflammation Swollen vulva, anaphylactic shock Death
Undetermined frequency (cannot be estimated from available data)	Injection site skin change (blemishes) ¹

¹can persist for up to 21 days following administration.

Pigs:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Injection site swelling, Injection site inflammation Anaphylactic shock Rectal oedema ¹ , rectal prolapse (partial) Erythema, pruritus Death
Undetermined frequency (cannot be estimated from available data)	Injection site skin change (blemishes) ²

¹oedema of the rectal mucosae

²can persist for up to 21 days following administration.

Sheep, goats:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Injection site swelling, Injection site inflammation Anaphylactic shock Death
Undetermined frequency (cannot be estimated from available data)	Injection site skin change (blemishes) ¹

¹can persist for up to 21 days following administration.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder <or its local representative> or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Pregnancy:

Studies in laboratory animals have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. No studies have been conducted in the target species. Use only according to the benefit/risk assessment by the responsible veterinarian.

3.8 Interaction with other medicinal products and other forms of interaction

None known.

3.9 Administration routes and dosage

Intramuscular or slow intravenous use (only in cattle)

Cattle:

5-10 mg tylosin/kg bodyweight per day during 3 days (2.5 to 5 ml solution for injection per 100 kg bodyweight) Maximum injection volume per injection site should not exceed 15 ml.

Sheep and goats:

10 mg tylosin/kg bodyweight per day during 3 days (5 ml solution for injection per 100 kg bodyweight)

For sheep over 50 kg of bodyweight, the injection should be divided over two injection sites (maximum 2.5 ml injection volume per injection site).

Pigs:

5-10 mg tylosin/kg bodyweight per day during 3 days (2.5 to 5 ml solution for injection per 100 kg bodyweight)

In pigs do not administer more than 5 ml per injection site.

The closures should not be broached more than 15 times. In order to prevent excessive broaching of the stopper, a suitable multiple dosing device should be used.

To ensure a correct dosage, body weight should be determined as accurately as possible.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

In pigs and cattle (calves) an intramuscular injection of 30 mg/kg bodyweight per day during 5 consecutive days produced no adverse effects.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Cattle

Meat and offal: 28 days.

Milk: 108 hours.

Sheep and goats:

Meat and offal: 42 days.

Milk: 108 hours

Pigs:

Meat and offal: 16 days.

4. PHARMACOLOGICAL INFORMATION

4.1 ATCvet code :

QJ01FA90

4.2 Pharmacodynamics

Tylosin is a macrolide antibiotic with a pKa of 7.1. Tylosin is structurally similar to erythromycin. It is produced by *Streptomyces fradiae*. Tylosin has a low solubility in water. Tylosin exerts its antibiotic activity by a similar mechanism to other macrolides, i.e. by binding the 50 S fraction of the ribosomes resulting, in an inhibition of the synthesis of proteins. Tylosin has mainly a bacteriostatic activity. Tylosin has an antibiotic effect against Gram-positive cocci (*Staphylococci*, *Streptococci*), Gram-positive bacilli (like *Erysipelothrix*), certain Gram-negative bacilli and *Mycoplasma*.

Resistance to macrolides is usually plasmid-mediated but modification of ribosomes may occur through chromosomal mutation. Resistance can occur by i) decreased entry into bacteria (most common with the Gram-negative bacteria), ii) synthesis of bacterial enzymes that hydrolyse the drug and, iii) modification of the target (the ribosome).

This latter resistance type may also lead to cross-resistance with other antibiotics that preferentially bind to bacterial ribosome. Gram-negative anaerobic bacteria are often resistant.

4.3 Pharmacokinetics

Absorption:

Following intramuscular injection the tylosin concentration reaches its maximum at 3-4 hours following administration.

Distribution:

The maximum concentration in milk of cattle and sows is 3-6 times higher than the blood concentration about 6 hours following injection. In bovine and porcine lungs maximum tylosin concentrations of 7-8 times higher than the maximum concentrations in serum were found at 6-24 hours following intramuscular injection. In cattle (whether in heat or not) the Mean Residence Time (MRT) in uterus secretions of tylosin injected by intravenous route at a dose rate of 10 mg/kg was about 6-7 times higher than the one measured in serum. This illustrates that in uterine secretions a single tylosin injection at a dose rate of 10 mg/kg during 24 hours can result in concentrations exceeding the MIC₉₀ of tylosin for *Arcanobacterium pyogenes*, one of the pathogens frequently isolated when metritis is diagnosed in cattle.

Elimination:

Tylosin is eliminated in unchanged form in bile and urine.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

Shelf life after first opening of the immediate packaging: 28 days.

5.3 Special precautions for storage

Protect from light.

Store in the original container.

Do not store above 25°C.

Do not freeze.

5.4 Nature and composition of immediate packaging

The veterinary medicinal product is presented in 50 ml, 100 ml or 250 ml Type II colourless glass vials, sealed with a bromobutyl stopper and aluminium cap supplied in a carton. One vial per carton.

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater <or household waste>.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

Biovet JSC

7. MARKETING AUTHORISATION NUMBER(S)

8. DATE OF FIRST AUTHORISATION

<Date of first authorisation:> <{DD/MM/YYYY}> <{DD month YYYY}.>

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

<{MM/YYYY}>

<{DD/MM/YYYY}>

<{DD month YYYY}>

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

{carton of 50/100 /250 ml glass vial}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

AXENTYL 200 mg/ml solution for injection

2. STATEMENT OF ACTIVE SUBSTANCES

Each ml contains:

Active substance:

Tylosin 200 000 IU

3. PACKAGE SIZE

50 ml (100 ml and 250ml)

4. TARGET SPECIES

Cattle, sheep, goats, pigs.

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Intramuscular or intravenous (cattle only) use

7. WITHDRAWAL PERIODS

Withdrawal periods:

Cattle:

Meat and offal: 28 days.

Milk: 108 hours.

Sheep and goats:

Meat and offal: 42 days.

Milk: 108 hours

Pigs:

Meat and offal: 16 days.

8. EXPIRY DATE

Exp {mm/yyyy}

Once broached use within 28 days.

Once opened use by...

9. SPECIAL STORAGE PRECAUTIONS

Protect from light. Store in the original container. Do not store above 25 °C . Do not freeze.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Biovet JSC

14. MARKETING AUTHORISATION NUMBERS

15. BATCH NUMBER

Lot {number}

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

{50/100 /250 ml glass vial}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

AXENTYL 200 mg/ml solution for injection

2. STATEMENT OF ACTIVE SUBSTANCES

Each ml contains:

Active substance:

Tylosin 200 000 IU

3. TARGET SPECIES

Cattle, sheep, goats, pigs.

4. ROUTES OF ADMINISTRATION

Intramuscular or intravenous (cattle only) use

Read the package leaflet before use

5. WITHDRAWAL PERIODS

Cattle:

Meat and offal: 28 days.

Milk: 108 hours.

Sheep and goats:

Meat and offal: 42 days.

Milk: 108 hours

Pigs:

Meat and offal: 16 days.

6. EXPIRY DATE

Exp {mm/yyyy}

Once broached use within 28 days.

Once opened use by...

7. SPECIAL STORAGE PRECAUTIONS

Protect from light. Store in the original container. Do not store above 25 °C. Do not freeze.

8. NAME OF THE MARKETING AUTHORISATION HOLDER

Biovet JSC

9. BATCH NUMBER

Lot {number}

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

AXENTYL 200 mg/ml solution for injection for cattle, sheep, goats and pigs

2. Composition

Each ml contains:

Active substance:

Tylosin 200 000 IU

Excipients:

Benzyl alcohol (E1519) 40 mg

A pale yellow to amber-coloured liquid.

3. Target species

Cattle, sheep, goats, pigs.

4. Indications for use

Infections caused by microorganisms susceptible to tylosin.

Cattle:

- Treatment of respiratory infections, metritis caused by Gram-positive micro-organisms, mastitis caused by *Streptococcus* spp., *Staphylococcus* spp. and interdigital necrobacillosis i.e. panaritum or foot rot

Cattle (calves):

- Treatment of respiratory infections and necrobacillosis.

Pigs:

- Treatment of enzootic pneumonia, haemorrhagic enteritis, erysipelas and metritis.
- Treatment of arthritis caused by *Mycoplasma* and *Staphylococcus* spp.

Sheep and goats:

- Treatment of respiratory infections, metritis caused by Gram-positive microorganisms, mastitis caused by Gram-positive microorganisms or *Mycoplasma* spp.

5. Contraindications

Do not use in horses. Intramuscular injection can be fatal in chickens and turkeys.

Do not use in cases of hypersensitivity to tylosin, to other macrolides or to any of the excipients.

6. Special warnings

Special warnings:

None.

Special precautions for safe use in the target species:

Due to likely variability (time, geographical) in susceptibility of bacteria to tylosin, bacteriological sampling and susceptibility testing are recommended.

Use of the veterinary medicinal product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to tylosin and may decrease the effectiveness of treatment with other macrolide antibiotics due to the potential for cross resistance. Official, national and regional antimicrobial policies should be taken into account when the veterinary medicinal product is used.

The efficacy data do not support the use of tylosin for the treatment of bovine mastitis caused by *Mycoplasma* spp. Haemorrhagic enteritis caused by *Brachyspira hyodysenteriae* must be treated with caution due to a high rate of in vitro resistance in European strains.

Where repeat injections are to be administered, use different sites for each injection.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Care should be taken to avoid accidental self-injection.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

Wash hands after use.

Tylosin may induce irritation. Macrolides, such as tylosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tylosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

Do not handle the veterinary medicinal product if you are allergic to ingredients in the veterinary medicinal product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice immediately and show the package leaflet or the label to the physician. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Pregnancy:

Studies in laboratory animals have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. No studies have been conducted in the target species. Use only according to the benefit/risk assessment by the responsible veterinarian.

Interaction with other medicinal products and other forms of interaction:

None known.

Overdose:

In pigs and cattle (calves) an intramuscular injection of 30 mg/kg bodyweight per day during 5 consecutive days produced no adverse effects.

Major incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

7. Adverse events

Cattle:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Injection site swelling, Injection site inflammation Swollen vulva, anaphylactic shock Death
Undetermined frequency (cannot be estimated from available data)	Injection site skin change (blemishes) ¹

¹can persist for up to 21 days following administration.

Pigs:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Injection site swelling, Injection site inflammation Anaphylactic shock Rectal oedema ¹ , rectal prolapse (partial) Erythema, pruritus Death
Undetermined frequency (cannot be estimated from available data)	Injection site skin change (blemishes) ²

¹oedema of the rectal mucosae

²can persist for up to 21 days following administration.

Sheep, goats:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Injection site swelling, Injection site inflammation Anaphylactic shock Death
Undetermined frequency (cannot be estimated from available data)	Injection site skin change (blemishes) ¹

¹can persist for up to 21 days following administration.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder <or its local representative> using the contact details at the end of this leaflet, or via your national reporting system: { national system details }

8. Dosage for each species, routes and method of administration

Intramuscular or slow intravenous use (only in cattle)

Cattle:

5-10 mg tylosin/kg bodyweight per day during 3 days (2.5 to 5 ml solution for injection per 100 kg bodyweight. Maximum injection volume per injection site should not exceed 15 ml.

Sheep and goats:

10 mg tylosin/kg bodyweight per day during 3 days (5 ml solution for injection per 100 kg bodyweight.

For sheep over 50 kg of bodyweight, the injection should be divided over two injection sites (maximum 2.5 ml injection volume per injection site).

Pigs:

5-10 mg tylosin/kg bodyweight per day during 3 days (2.5 to 5 ml solution for injection per 100 kg bodyweight.

In pigs do not administer more than 5 ml per injection site.

To ensure a correct dosage, body weight should be determined as accurately as possible.

9. Advice on correct administration

The closures should not be broached more than 15 times. In order to prevent excessive broaching of the stopper, a suitable multiple dosing device should be used.

10. Withdrawal periods

Cattle:

Meat and offal: 28 days.

Milk: 108 hours.

Sheep and goats:

Meat and offal: 42 days.

Milk: 108 hours

Pigs:

Meat and offal: 16 days.

11. Special storage precautions

Keep out of the sight and reach of children.

Shelf life after first opening the immediate packaging: 28 days.

Protect from light. Store in the original container. Do not store above 25°C. Do not freeze.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp. The expiry date refers to the last day of that month.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection system. These measures should help to protect the environment.

Ask your veterinary surgeon how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack sizes

The veterinary medicinal product is presented in 50,100 or 250 ml Type II colourless glass vials, sealed with a bromobutyl stopper and aluminium cap supplied in a carton. One vial per carton.

Not all pack sizes may be marketed.

15. Date on which the package leaflet was last revised

<{MM/YYYY}>

<{DD/MM/YYYY}>

<{DD month YYYY}>

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

16. Contact details

Marketing authorisation holder and manufacturer responsible for batch release <and contact details to report suspected adverse events>:

Biovet JSC
39 Petar Rakov Str
4550 Peshtera
Bulgaria

<Local representatives <and contact details to report suspected adverse events>:>

17. Other information