

**B. PACKAGE LEAFLET**



Transient increase of temperature is very rare after vaccination. In very rare cases hypersensitivity reactions may occur. In such case, appropriate treatment should be initiated without delay.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Cats.

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Dose - 1 ml regardless of age, weight and breed of the individual, but not earlier than at 8 weeks of age.

Method of administration - subcutaneous, preferably in the region behind the shoulder blade.

### Vaccination schedule:

Basic vaccination:

Two administrations at an interval of 3 weeks.

The first vaccination with one dose of Biofel PCH in kittens from the age of 8-10 weeks and the second vaccination with one dose of Biofel PCHR from the age of 3 months.

Revaccination:

Further regular revaccination with Biofel PCHR is carried out in 12-month intervals.

## **9. ADVICE ON CORRECT ADMINISTRATION**

The vial contents should be shaken well before use.

## **10. WITHDRAWAL PERIOD(S)**

Not applicable

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Store in a refrigerator (2°C - 8°C).

Protect from light.

Protect from frost.

Shelf life after first opening the immediate packaging: 8 hours.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP.

## **12. SPECIAL WARNING(S)**

### Special warnings for each target species

Vaccinate healthy animals only. Any antiparasitic treatment should precede the vaccination by at least 10 days.

Effect of maternal antibodies on the vaccine efficacy has not been studied.

### Pregnancy:

Vaccination of pregnant cats is recommended in the first half of pregnancy.

### Interaction with other medicinal products and other forms of interaction

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal products. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

### Incompatibilities

Do not mix with other veterinary medicinal products.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals

To the user:

This product contains mineral oil. Accidental injection/self-injection with the product may result in severe pain and swelling, particularly if injected into a joint or finger, and in rare cases can lead to loss of the affected finger if not given prompt medical attention.

If you are accidentally injected with this product, seek prompt medical advice even if only a very small amount is injected and take the package leaflet with you.

If pain persists for more than 12 hours after medical examination, seek medical advice again.

To the physician:

This product contains mineral oil. Even if small amounts have been injected, accidental injection with this product can cause intense swelling, which may, for example, result in ischaemic necrosis and even the loss of a digit. Expert, prompt surgical attention is required and may necessitate early incision and irrigation of the injected area, especially where there is involvement of finger pulp or tendon.

## **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

## **14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

## **15. OTHER INFORMATION**

For animal treatment only.

Veterinary medicinal product subject to prescription.

Package size:

2 x 1 dose, 10 x 1 dose, 20 x 1 dose, 100 x 1 dose,

1 x 5 doses, 5 x 5 dose, 10 x 5 doses

Not all pack sizes may be marketed.

