

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Xylexx 20 mg/ml solution for injection for cattle, horses, dogs and cats (AT, BE, BG, CY, CZ, DE, EE, EL, ES, FR, HU, HR, IE, SI, IT, LT, LU, LV, NL-PL, PT, RO, SK, UK(NI))

Xylexx Vet 20 mg/ml solution for injection for cattle, horses, dogs and cats (DK, FI, IS, NO and SE)

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

1 ml contains:

Active substance:

Xylazine 20.0 mg
(equivalent to 23.31 mg xylazine hydrochloride)

Excipient(s):

Benzethonium chloride 0.11 mg

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for injection.

Clear, colourless to almost colourless solution, practically free from visible particles.

4. CLINICAL PARTICULARS

4.1 Target species

Cattle, horses, dogs and cats.

4.2 Indications for use, specifying the target species

In cattle, horses, dogs and cats:

- sedation;
- premedication in combination with an anaesthetic.

4.3 Contraindications

Do not use in known cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in animals with gastrointestinal obstruction, as it is a muscle relaxant and the properties of the medicinal product appear to enhance the effects of an obstruction, and because of the risk of vomiting.

Do not use in cases of pulmonary disease (breathing deficiency) or cardiac disorders (especially in case of ventricular arrhythmia).

Do not use in cases of impaired liver or renal function. Do not use in cases of predetermined history of seizures. Do not use in cases of hypotension and shock.

Do not use in animals with diabetes mellitus.

Do not administer simultaneously with sympathomimetic amines (e.g. epinephrine).

Do not use in calves less than 1 week of age, foals less than 2 weeks of age or puppies and kittens under 6 weeks of age. Do not use during the last stage of pregnancy (danger of premature birth), except at parturition (see section 4.7).

4.4 Special warnings for each target species

Cattle:

- Ruminants are highly susceptible to the effects of xylazine. Normally cattle remain standing at the lower doses, but some animals may lie down. At the highest recommended doses most animals will lie down and some animals may lapse into lateral recumbency.
- Reticulo-ruminal motor functions are depressed after injection of xylazine. This may result in bloat. It is advisable to withhold feed and water in adult cattle for several hours before administration of xylazine. Fasting in calves might be indicated but should only be done at the discretion of a benefit/risk assessment made by the responsible veterinarian.
- In cattle the ability to eructate, cough and swallow is retained but reduced during the period of sedation, therefore cattle must be closely watched during the recovery period: the animals should be maintained in sternal recumbency.
- In cattle life threatening effects may occur after intramuscular doses above 0.5 mg/kg body weight (respiratory and circulatory failure). Therefore, very precise dosing is required.
- The combination with other pre-anaesthetic agents or anaesthetic agents should be the subject of a benefit/risk assessment. This assessment should consider the composition of the products, their dose and the nature of the surgery. Recommended dosages are likely to vary according to the choice of the anaesthetic combination.

Horses:

- Xylazine inhibits the normal intestinal motility. Therefore, it should only be used in horses with colic that are not responsive to analgesics. The use of xylazine should be avoided in horses with caecal malfunction.
- After treatment of horses with xylazine, the animals are reluctant to walk, so whenever possible the drug should be administered in the place where the treatment/investigation is going to take place.
- Caution should be taken in the administration of the product to horses susceptible to laminitis.
- Horses with airway disease or malfunction may develop life-threatening dyspnoea.
- The dose should be kept as low as possible.
- The combination with other pre-anaesthetic agents or anaesthetic agents should be the subject of a benefit/risk assessment. This assessment should consider the composition of the products, their dose and the nature of the surgery. Recommended dosages are likely to vary according to the choice of the anaesthetic combination.

Dogs and cats:

- Xylazine inhibits normal intestinal motility. This may make xylazine sedation undesirable for upper gastro-intestinal radiographs, because it promotes filling of the stomach with gas and makes interpretation less certain.
- Brachycephalic dogs with airway disease or malfunction may develop life-threatening dyspnoea.
- The combination with other pre-anaesthetic agents or anaesthetic agents should be the subject of a benefit/risk assessment. This assessment should consider the composition of the products, their dose and the nature of the surgery. Recommended dosages are likely to vary according to the choice of the anaesthetic combination.

4.5 Special precautions for use

Special precautions for use in animals

- Keep the animals calm, because they may respond to external stimuli.
- Avoid intra-arterial administration.
- Tympany may occasionally occur in recumbent cattle and can be avoided by maintaining the animal in sternal recumbency.
- To avoid aspiration of saliva or food, lower the animal's head and neck. Fast the animals before use of the product.
- Older and exhausted animals are more sensitive to xylazine, whilst nervous or highly excitable animals may require a relatively high dose.
- In case of dehydration, xylazine should be used cautiously.

- Emesis is generally seen within 3-5 minutes after xylazine administration in cats and dogs. It is advisable to fast dogs and cats for 12 hours prior to surgery; they may have free access to drinking water.
- Pre-medication with atropine in cats and dogs may reduce salivation and bradycardia effects
- Do not exceed the recommended dosage.
- Following administration animals should be allowed to rest quietly until the full effect has been reached.
- It is advised to cool animals when the ambient temperature is above 25°C and to keep animals warm at low temperatures.
- For painful procedures, xylazine should always be used in combination with local or general anaesthesia.
- Xylazine produces a certain degree of ataxia; therefore, xylazine must be used cautiously in procedures involving the distal extremities and in standing castrations in the horse.
- Treated animals should be monitored until the effect has faded totally (e.g. cardiac and respiratory function, also in the post-operative phase) and should be segregated to avoid bullying.
- For use in young animals, see the age restriction mentioned in section 4.3. If the product is intended to be used in young animals below these age-limits, a benefit/risk assessment should be made by the veterinarian.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

This product is a sedative. Care should be taken to avoid accidental self-injection.

In the case of accidental oral intake or self-injection, seek medical advice immediately and show the package leaflet to the doctor, but DO NOT DRIVE, as sedation and changes in blood pressure may occur.

Avoid skin, eye or mucosal contact. In the case of accidental contact of the product with the skin or eyes, rinse with large amounts of fresh water. Remove contaminated clothes that are in direct contact with the skin. If symptoms occur, seek medical advice.

If pregnant women handle the product, special caution should be observed not to self-inject as uterine contractions and decreased foetal blood pressure may occur after accidental systemic exposure.

Advice to doctors

Xylazine is an α_2 -adrenoreceptor agonist. Symptoms after absorption may involve clinical effects including dose-dependent sedation, respiratory depression, bradycardia, hypotension, a dry mouth, and hyperglycaemia. Ventricular arrhythmias have also been reported. Respiratory and haemodynamic symptoms should be treated symptomatically.

4.6 Adverse reactions (frequency and seriousness)

In general, side effects, typical for an α_2 -adrenergic agonist, like bradycardia, reversible arrhythmia and hypotension can occur. Thermoregulation can be influenced and consequently body temperature can decrease or increase dependant on the ambient temperature. Depression of respiration and / or respiratory arrest can occur, especially in cats.

Cattle:

- Reversible local tissue irritation.
- In cattle xylazine may induce premature parturition, and it also reduces implantation of the ovum.
- Cattle, which have received high doses of xylazine sometimes suffer from loose faeces for 24 hours afterwards.
- Other adverse reactions include snoring, profound salivation, ruminal atony, atony of the tongue, regurgitation, bloating, nasal stridor, hypothermia, bradycardia, increased urination and reversible prolapse of the penis.
- In cattle, adverse effects are generally more pronounced after intramuscular administration compared to intravenous.

Horses:

- Reversible local tissue irritation.
- Horses often sweat as the effects of the sedation are wearing off.
- Severe bradycardia and reduced respiratory rate have been reported especially in horses.
- Following administration to horses, a transient rise followed by a fall in blood pressure usually occurs.
- More frequent urination has been reported
- Muscle tremors and movement in response to sharp auditory or physical stimuli are possible. Although rare, violent reactions have been reported in horses following the administration of xylazine.
- Ataxia and reversible prolapse of the penis may occur.
- In very rare cases xylazine may induce mild colic as the gut motility is depressed temporarily. As a preventive measure the horse should receive no feed after sedation until the effect has faded completely

Dogs and cats:

- Reversible local tissue irritation.
- Cats and dogs frequently vomit during the onset of the xylazine-induced sedation, especially when the animals have just been fed.
- Animals may show profound salivation following an injection with xylazine.
- Other adverse effects for dogs and cats include: muscle tremors, bradycardia with AV-block, hypotension, reduced respiratory rate, movement in response to strong auditory stimuli, hyperglycaemia and increased urination in cats.
- In cats xylazine causes uterine contractions and it may induce premature parturition.
- In dogs, adverse effects are generally more pronounced after subcutaneous administration compared to intramuscular and the effect (efficacy) can be less predictable.
- In susceptible dog breeds with a large chest (Great Dane, Irish Setter) rare cases of bloating have been reported.
- In anaesthetized animals, mainly during and after the recovery period, in very rare cases, cardio-respiratory disturbances (cardiac arrest, dyspnoea, bradypnea, pulmonary edema, hypotension) and neurological signs (seizures, prostration, pupillary disorders, tremors) were observed.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

4.7 Use during pregnancy, lactation and lay

Pregnancy

Although laboratory studies in rats have not shown any evidence of teratogenic or foetotoxic effects the use of the product during the first two thirds of pregnancy should only be made according to the benefit/risk assessment by the responsible veterinarian.

Do not use in the later stages of pregnancy (particularly in cattle and cats) except at parturition, because xylazine causes uterine contractions and it may induce premature labour.

Do not use in cattle receiving ovum transplants or in cattle at the time of implantation of the ovum as the increased uterine tone may reduce the chance of implantation of the ovum.

Lactation

The veterinary medicinal product can be used in lactating animals.

4.8 Interaction with other medicinal products and other forms of interaction

Other CNS depressant agents (barbiturates, narcotics, anaesthetics, tranquillizers, etc.) may cause additive CNS depression if used with xylazine. Dosages of these agents may need to be reduced. Xylazine should therefore be used cautiously in combination with neuroleptics or tranquillizers. Xylazine should not be used in combination with sympathomimetic drugs such as epinephrine as ventricular arrhythmia may follow.

The concurrent intravenous use of potentiated sulphonamides with alpha-2 agonists has been reported to cause cardiac arrhythmias which may be fatal. Whilst no such effects have been reported with this product, it is recommended that intravenous administration of Trimethoprim/Sulphonamide containing products should not be undertaken when horses have been sedated with xylazine.

4.9 Amounts to be administered and administration route

Cattle: intravenous use, intramuscular use.
Horses: intravenous use.
Dogs: intramuscular use.
Cats: intramuscular use, subcutaneous use.

To ensure a correct dosage body weight should be determined as accurately as possible.

The intravenous injection should be given slowly, especially in horses.

This veterinary medicinal product is for administration only by a veterinarian or under their supervision.

Cattle (IV,IM)

Dosage:

Dosage cattle			
Dose level	xylazine (mg/kg)	Xylexx 20 mg/ml (ml/100 kg)	Xylexx 20 mg/ml (ml/500 kg)
A. Intramuscular			
I	0.05	0.25	1.25
II	0.1	0.5	2.5
III	0.2	1	5
IV	0.3	1.5	7.5
B. Intravenous			
I	0.016-0.024	0.08-0.12	0.4-0.6
II	0.034-0.05	0.17-0.25	0.85-1.25
III	0.066-0.10	0.33-0.5	1.65-2.5

Dosage I: Sedation with slight reduction of muscle tone. The cattle are still able to stand.

Dosage II: Sedation with pronounced reduction of muscle tone and slight analgesia. The cattle mostly remain able to stand but may also lie down.

Dosage III: Deep sedation, further reduction in muscle tone, partial analgesia. The cattle lie down.

Dosage IV: Very deep sedation with a pronounced reduction in muscle tone, partial analgesia. The cattle lie down.

Horses (IV)

Dosage: single injection of 0.6-1 mg xylazine per kg body weight. (3-5 ml product per 100 kg body weight).

Dogs (IM)

Dosage: single injection of 0.5-3 mg xylazine per kg body weight. (0.25-1.5 ml product per 10 kg bodyweight).

Cats (IM, SC)

Dosage: single injection of 0.5-1 mg xylazine per kg body weight. (0.025-0.05 ml product per kg body weight).

The vial can be broached up to 30 times.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

In the event of an accidental overdose, cardiac arrhythmias, hypotension, and profound CNS and respiratory depression may occur. Seizures have also been reported after an overdose. Xylazine can be antagonized by α 2-adrenergic antagonists.

To treat the respiratory depressant effects of xylazine, mechanical respiratory support with or without respiratory stimulants (e.g. doxapram) can be recommended.

4.11 Withdrawal periods

Cattle:

Meat and offal: 1 day

Milk: zero hours

Horses:

Meat and offal: 1 day

Not authorised for use in mares producing milk for human consumption.

5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: psycholeptics, hypnotics and sedatives, xylazine.

ATCvet code: QN05CM92

5.1 Pharmacodynamic properties

Xylazine belongs to the α 2-adrenoceptor agonists.

- Xylazine is a α 2-adrenoceptor agonist, that acts by stimulation of central and peripheral α 2-adrenoceptors. Through its central stimulation of α 2-adrenoceptors, xylazine has potent antinociceptive activity. In addition to α 2-adrenergic activity, xylazine has α 1-adrenergic effects.
- Xylazine also produces skeletal muscle relaxation by inhibition of intraneuronal transmission of impulses at the central level of the central nervous system. The analgesic and skeletal muscle relaxation properties of xylazine show considerable interspecies variations. Sufficient analgesia generally will be attained in combination with other products only.
- In many species, administration of xylazine produces a short-lived arterial pressor effect followed by a longer period of hypotension and bradycardia. These contrasting actions upon the arterial pressure apparently are related to the α 2- and α 1-adrenergic actions of xylazine.
- Xylazine has several endocrine effects. Insulin (mediated by α 2-receptors in pancreatic β -cells which inhibit insulin release), ADH (decreased production of ADH, causing polyuria) and FSH (decreased) are reported to be influenced by xylazine.

5.2 Pharmacokinetic particulars

Absorption (and action) is rapid following intramuscular injection. Levels of drug peak rapidly (usually within 15 minutes) and then decline exponentially. Xylazine is a highly lipid soluble organic base and diffuses extensively and rapidly (V_d 1.9-2.7 L/kg bw). Within minutes after an intravenous injection, it can be found in a high concentration in the kidneys, the liver, the CNS, the hypophyses, and the diaphragm. So, there is a very rapid transfer from the blood vessels to the tissues. Intramuscular bioavailability is incomplete and variable ranging from 52-90% in the dog to 40-48% in the horse. Xylazine is metabolised extensively and eliminated rapidly (\pm 70% via the urine, while the

enteric elimination is $\pm 30\%$). The rapid elimination of xylazine is probably attributable to the extensive metabolism rather than to the renal excretion of unchanged xylazine.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Benzethonium chloride
Sodium hydroxide (for pH adjustment)
Hydrochloric acid, dilute (for pH adjustment)
Water for injections

6.2 Major incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

6.3 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.
Shelf life after first opening the immediate packaging: 28 days.

6.4 Special precautions for storage

Do not refrigerate or freeze.

6.5 Nature and composition of immediate packaging

Clear type II glass vials containing 30 ml product, closed with a bromobutyl rubber stopper and aluminium cap in a cardboard or polystyrene box.

Pack sizes:

Cardboard box with 1 vial of 30 ml
Cardboard box with 5 vials of 30 ml
Polystyrene box with 24 vials of 30 ml

Not all pack sizes may be marketed.

6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Alfasan Nederland B.V.
Kuipersweg 9
3449 JA Woerden
The Netherlands

8. MARKETING AUTHORISATION NUMBER

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

10. DATE OF REVISION OF THE TEXT

PROHIBITION OF SALE, SUPPLY AND/OR USE

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Cardboard box or polystyrene box

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Xylexx 20 mg/ml solution for injection
xylazine

2. STATEMENT OF ACTIVE SUBSTANCES

1 ml contains:

Xylazine 20.0 mg
(equivalent to 23.31 mg xylazine hydrochloride)

3. PHARMACEUTICAL FORM

Solution for injection

4. PACKAGE SIZE

30 ml
5 x 30 ml
24 x 30 ml

5. TARGET SPECIES

Cattle, horses, dogs and cats.

6. INDICATION(S)**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Cattle: intravenous use, intramuscular use
Horses: intravenous use
Dogs: intramuscular use
Cats: intramuscular use, subcutaneous use

Read the package leaflet before use.

8. WITHDRAWAL PERIODS

Withdrawal periods:
Cattle: Meat and offal: 1 day, Milk: zero hours.
Horses: Meat and offal: 1 day, Not authorised for use in mares producing milk for human consumption.

9. SPECIAL WARNING(S), IF NECESSARY

This product is harmful and causes sedation. Avoid accidental self-injection and skin, eye or mucosal contact. See package leaflet for full user warnings.

10. EXPIRY DATE

EXP {month/year}

Shelf life after first opening the container: 28 days

Once broached use by...

11. SPECIAL STORAGE CONDITIONS

Do not refrigerate or freeze.

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: read package leaflet.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only. To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Alfasan Nederland B.V.

Kuipersweg 9

3449 JA Woerden

The Netherlands


16. MARKETING AUTHORISATION NUMBER(S)**17. MANUFACTURER’S BATCH NUMBER**

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS
Glass vials of 30 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Xylexx 20 mg/ml solution for injection
xylazine



2. QUANTITY OF THE ACTIVE SUBSTANCE(S)

Xylazine 20.0 mg/ml (equivalent to 23.31 mg/ml xylazine hydrochloride)

3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

30 ml

4. ROUTE(S) OF ADMINISTRATION

IV, IM, SC

5. WITHDRAWAL PERIOD

Withdrawal periods:

Cattle: Meat and offal: 1 day, Milk: zero hours.

Horses: Meat and offal: 1 day, Not authorised for use in mares producing milk for human consumption.

6. BATCH NUMBER

Lot {number}

7. EXPIRY DATE

EXP {month/year}

Shelf life after first opening the container: 28 days

Once broached use by...

8. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

B. PACKAGE LEAFLET

PACKAGE LEAFLET:

Xylexx 20 mg/ml solution for injection for cattle, horses, dogs and cats (AT, BE, BG, CY, CZ, DE, EE, EL, ES, FR, HU, HR, IE, SI, IT, LT, LU, LV, NL, PL, PT, RO, SK, UK(NI))
Xylexx Vet 20 mg/ml solution for injection for cattle, horses, dogs and cats (DK, FI, IS, NO and SE)

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder and manufacturer responsible for batch release:

Alfasan Nederland B.V.
Kuipersweg 9
3449 JA Woerden
The Netherlands

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Xylexx 20 mg/ml solution for injection for cattle, horses, dogs and cats
xylazine

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

1 ml contains:

Active substance:

Xylazine 20.0 mg
(equivalent to 23.31 mg xylazine hydrochloride)

Excipient(s):

Benzethonium chloride 0.11 mg

Clear, colourless to almost colourless solution for injection, practically free from visible particles.

4. INDICATION(S)

In cattle, horses, dogs and cats:

- sedation;
- premedication in combination with an anaesthetic.

5. CONTRAINDICATIONS

Do not use in known cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in animals with gastrointestinal obstruction, as it is a muscle relaxant and the properties of the medicinal product appear to enhance the effects of an obstruction, and because of the risk of vomiting.

Do not use in cases of pulmonary disease (breathing deficiency) or cardiac disorders (especially in case of ventricular arrhythmia).

Do not use in cases of impaired liver or renal function. Do not use in cases of predetermined history of seizures. Do not use in cases of hypotension and shock.

Do not use in animals with diabetes mellitus.

Do not administer simultaneously with sympathomimetic amines (e.g. epinephrine).

Do not use in calves less than 1 week of age, foals less than 2 weeks of age or puppies and kittens under 6 weeks of age. Do not use during the last stage of pregnancy (danger of premature birth), except at parturition (see Special Warnings: Pregnancy and Lactation).

6. ADVERSE REACTIONS

In general, side effects, typical for an α_2 -adrenergic agonist, like bradycardia, reversible arrhythmia and hypotension can occur. Thermoregulation can be influenced and consequently body temperature can decrease or increase dependant on the ambient temperature. Depression of respiration and / or respiratory arrest can occur, especially in cats.

Cattle:

- Reversible local tissue irritation.
- In cattle xylazine may induce premature parturition, and it also reduces implantation of the ovum.
- Cattle, which have received high doses of xylazine sometimes suffer from loose faeces for 24 hours afterwards.
- Other adverse reactions include snoring, profound salivation, ruminal atony, atony of the tongue, regurgitation, bloating, nasal stridor, hypothermia, bradycardia, increased urination and reversible prolapse of the penis.
- In cattle, adverse effects are generally more pronounced after intramuscular administration compared to intravenous.

Horses:

- Reversible local tissue irritation.
- Horses often sweat as the effects of the sedation are wearing off.
- Severe bradycardia and reduced respiratory rate have been reported especially in horses.
- Following administration to horses, a transient rise followed by a fall in blood pressure usually occurs.
- More frequent urination has been reported
- Muscle tremors and movement in response to sharp auditory or physical stimuli are possible. Although rare, violent reactions have been reported in horses following the administration of xylazine.
- Ataxia and reversible prolapse of the penis may occur.
- In very rare cases xylazine may induce mild colic as the gut motility is depressed temporarily. As a preventive measure the horse should receive no feed after sedation until the effect has faded completely

Dogs and cats:

- Reversible local tissue irritation.
- Cats and dogs frequently vomit during the onset of the xylazine-induced sedation, especially when the animals have just been fed.
- Animals may show profound salivation following an injection with xylazine.
- Other adverse effects for dogs and cats include: muscle tremors, bradycardia with AV-block, hypotension, reduced respiratory rate, movement in response to strong auditory stimuli, hyperglycaemia and increased urination in cats.
- In cats xylazine causes uterine contractions and it may induce premature parturition.
- In dogs, adverse effects are generally more pronounced after subcutaneous administration compared to intramuscular and the effect (efficacy) can be less predictable.
- In susceptible dog breeds with a large chest (Great Dane, Irish Setter) rare cases of bloating have been reported.
- In anaesthetized animals, mainly during and after the recovery period, in very rare cases, cardio-respiratory disturbances (cardiac arrest, dyspnoea, bradypnea, pulmonary edema, hypotension) and neurological signs (seizures, prostration, pupillary disorders, tremors) were observed.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon. Alternatively, you can report via your national reporting system.

7. TARGET SPECIES

Cattle, horses, dogs and cats.



8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Cattle: intravenous use, intramuscular use.
Horses: intravenous use.
Dogs: intramuscular use.
Cats: intramuscular use, subcutaneous use.

To ensure a correct dosage body weight should be determined as accurately as possible.
The intravenous injection should be given slowly, especially in horses.
This veterinary medicinal product is for administration only by a veterinarian or under their supervision.

The vial can be broached up to 30 times.

Cattle (IV, IM)

Dosage:

Dosage cattle			
Dose level	xylazine (mg/kg)	Xylexx 20 mg/ml (ml/100 kg)	Xylexx 20 mg/ml (ml/500 kg)
A. Intramuscular			
I	0.05	0.25	1.25
II	0.1	0.5	2.5
III	0.2	1	5
IV	0.3	1.5	7.5
B. Intravenous			
I	0.016-0.024	0.08-0.12	0.4-0.6
II	0.034-0.05	0.17-0.25	0.85-1.25
III	0.066-0.10	0.33-0.5	1.65-2.5

Dosage I: Sedation with slight reduction of muscle tone. The cattle are still able to stand.

Dosage II: Sedation with pronounced reduction of the muscle tone and slight analgesia. The cattle mostly remain able to stand but may also lie down.

Dosage III: Deep sedation, further reduction in muscle tone, partial analgesia. The cattle lie down.

Dosage IV: Very deep sedation with a pronounced reduction in the muscle tone, partial analgesia. The cattle lie down.

Horses (IV)

Dosage: single injection of 0.6 -1 mg xylazine per kg body weight. (3-5 ml product per 100 kg bodyweight).

Dogs (IM)

Dosage: single injection of 0.5-3 mg xylazine per kg body weight. (0.25-1.5 ml product per 10 kg bodyweight).

Cats (IM, SC)

Dosage: single injection of 0.5-1 mg xylazine per kg body weight. (0.025-0.05 ml product per kg bodyweight).

9. ADVICE ON CORRECT ADMINISTRATION

10. WITHDRAWAL PERIODS

Cattle: Meat and offal: 1 day, Milk: zero hours

Horses: Meat and offal: 1 day, Not authorised for use in mares producing milk for human consumption.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Do not refrigerate or freeze. Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month.

Shelf life after first opening the container: 28 days.

12. SPECIAL WARNINGS

Special warnings for each target species

Cattle:

- Ruminants are highly susceptible to the effects of xylazine. Normally cattle remain standing at the lower doses, but some animals may lie down. At the highest recommended doses most animals will lie down and some animals may lapse into lateral recumbency.
- Reticulo-ruminal motor functions are depressed after injection of xylazine. This may result in bloat. It is advisable to withhold feed and water in adult cattle for several hours before administration of xylazine. Fasting in calves might be indicated but should only be done at the discretion of a benefit/risk assessment made by the responsible veterinarian.
- In cattle the ability to eructate, cough and swallow is retained but reduced during the period of sedation, therefore cattle must be closely watched during the recovery period: the animals should be maintained in sternal recumbency.
- In cattle life threatening effects may occur after intramuscular doses above 0.5 mg/kg body weight (respiratory and circulatory failure). Therefore, very precise dosing is required.
- The combination with other pre-anaesthetic agents or anaesthetic agents should be the subject of a benefit/risk assessment. This assessment should consider the composition of the products, their dose and the nature of the surgery. Recommended dosages are likely to vary according to the choice of the anaesthetic combination.

Horses:

- Xylazine inhibits the normal intestinal motility. Therefore, it should only be used in horses with colic that are not responsive to analgesics. The use of xylazine should be avoided in horses with caecal malfunction.

- After treatment of horses with xylazine, the animals are reluctant to walk, so whenever possible the drug should be administered in the place where the treatment/investigation is going to take place.
- Caution should be taken in the administration of the product to horses susceptible to laminitis.
- Horses with airway disease or malfunction may develop life-threatening dyspnoea.
- The dose should be kept as low as possible.
- The combination with other pre-anaesthetic agents or anaesthetic agents should be the subject of a benefit/risk assessment. This assessment should consider the composition of the products, their dose and the nature of the surgery. Recommended dosages are likely to vary according to the choice of the anaesthetic combination.

Dogs and cats:

- Xylazine inhibits normal intestinal motility. This may make xylazine sedation undesirable for upper gastro-intestinal radiographs, because it promotes filling of the stomach with gas and makes interpretation less certain.
- Brachycephalic dogs with airway disease or malfunction may develop life-threatening dyspnoea.
- The combination with other pre-anaesthetic agents or anaesthetic agents should be the subject of a benefit/risk assessment. This assessment should consider the composition of the products, their dose and the nature of the surgery. Recommended dosages are likely to vary according to the choice of the anaesthetic combination.

Special precautions for use in animals

- Keep the animals calm, because they may respond to external stimuli.
- Avoid intra-arterial administration.
- Tympany may occasionally occur in recumbent cattle and can be avoided by maintaining the animal in sternal recumbency.
- To avoid aspiration of saliva or food, lower the animal's head and neck. Fast the animals before use of the product.
- Older and exhausted animals are more sensitive to xylazine, whilst nervous or highly excitable animals may require a relatively high dose.
- In case of dehydration, xylazine should be used cautiously.
- Emesis is generally seen within 3-5 minutes after xylazine administration in cats and dogs. It is advisable to fast dogs and cats for 12 hours prior to surgery; they may have free access to drinking water.
- Pre-medication with atropine in cats and dogs may reduce salivation and bradycardia effects.
- Do not exceed the recommended dosage.
- Following administration animals should be allowed to rest quietly until the full effect has been reached.
- It is advised to cool animals when the ambient temperature is above 25°C and to keep animals warm at low temperatures.
- For painful procedures, xylazine should always be used in combination with local or general anaesthesia.
- Xylazine produces a certain degree of ataxia; therefore, xylazine must be used cautiously in procedures involving the distal extremities and in standing castrations in the horse.
- Treated animals should be monitored until the effect has faded totally (e.g. cardiac and respiratory function, also in the post-operative phase) and should be segregated to avoid bullying.
- For use in young animals, see the age restriction mentioned in section "Contraindications". If the product is intended to be used in young animals below these age-limits, a benefit/risk assessment should be made by the veterinarian.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

This product is a sedative. Care should be taken to avoid accidental self-injection.

In the case of accidental oral intake or self-injection, seek medical advice immediately and show the package leaflet to the doctor, but DO NOT DRIVE, as sedation and changes in blood pressure may occur.

Avoid skin, eye or mucosal contact. In the case of accidental contact of the product with the skin or eyes, rinse with large amounts of fresh water. Remove contaminated clothes that are in direct contact with the skin. If symptoms occur, seek medical advice.

If pregnant women handle the product, special caution should be observed not to self-inject as uterine contractions and decreased foetal blood pressure may occur after accidental systemic exposure.

Advice to doctors:

Xylazine is an α_2 -adrenoreceptor agonist. Symptoms after absorption may involve clinical effects including

dose-dependent sedation, respiratory depression, bradycardia, hypotension, a dry mouth, and hyperglycaemia. Ventricular arrhythmias have also been reported. Respiratory and haemodynamic symptoms should be treated symptomatically.

Pregnancy

Although laboratory studies in rats have not shown any evidence of teratogenic or foetotoxic effects the use of the product during the first two thirds of pregnancy should only be made according to the benefit/risk assessment by the responsible veterinarian.

Do not use in the later stages of pregnancy (particularly in cattle and cats) except at parturition, because xylazine causes uterine contractions and it may induce premature labour.

Do not use in cattle receiving ovum transplants or in cattle at the time of implantation of the ovum as the increased uterine tone may reduce the chance of implantation of the ovum.

Lactation

The veterinary medicinal product can be used in lactating animals.

Interaction with other medicinal products and other forms of interaction

Other CNS depressant agents (barbiturates, narcotics, anaesthetics, tranquillizers, etc.) may cause additive CNS depression if used with xylazine. Dosages of these agents may need to be reduced. Xylazine should therefore be used cautiously in combination with neuroleptics or tranquillizers. Xylazine should not be used in combination with sympathomimetic drugs such as epinephrine as ventricular arrhythmia may follow.

The concurrent intravenous use of potentiated sulphonamides with α_2 -agonists has been reported to cause cardiac arrhythmias which may be fatal. Whilst no such effects have been reported with this product, it is recommended that intravenous administration of Trimethoprim/Sulphonamide containing products should not be undertaken when horses have been sedated with xylazine.

Overdose (symptoms, emergency procedures, antidotes)

In the event of an accidental overdose, cardiac arrhythmias, hypotension, and profound CNS and respiratory depression may occur. Seizures have also been reported after an overdose. Xylazine can be antagonized by α_2 -adrenergic antagonists.

To treat the respiratory depressant effects of xylazine, mechanical respiratory support with or without respiratory stimulants (e.g. doxapram) can be recommended.

Incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Medicines should not be disposed of via wastewater <or household waste>. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

15. OTHER INFORMATION

Clear type II glass vials containing 30 ml product, closed with a bromobutyl rubber stopper and aluminium cap in a cardboard box.

Pack sizes:

Cardboard box with 1 vial of 30 ml

Cardboard box with 5 vials of 30 ml

Polystyrene box with 24 vials of 30 ml

Not all pack sizes may be marketed.