

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Exitel Plus Tablets For Dogs (IE, IS, BG, EE, NL, HU, PT, LV, SK, LT, SI, ES, EL, AT, CZ, NO)  
 Exitel Tablets for Dogs (DE)  
 Easimax Plus Tablets For Dogs. (UK)  
 Exitel Plus (DK, CY, RO, LU)  
 Exitel vet.150 mg/144 mg/50 mg Tablets for dogs (SE)  
 FRONTCONTROL WORMER Tablets for Dogs (IT, BE, FR)  
 Frontcontrol Vet Tabletti koiralle (FI)

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains:

#### Active substances:

Praziquantel	50 mg
Pyrantel	50 mg (equivalent to 144 mg pyrantel embonate)
Febantel	150 mg

#### Excipients:

Qualitative composition of excipients and other constituents
Lactose monohydrate
Microcrystalline cellulose
Magnesium stearate
Colloidal anhydrous silica
Croscarmellose sodium
Sodium laurilsulfate
Pork flavour

A pale yellow tablet with a cross breakline on one side.  
 The tablets can be divided into 2 or 4 equal parts.

### 3. CLINICAL INFORMATION

#### 3.1 Target species

Dogs.

#### 3.2 Indications for use for each target species

Treatment of mixed infections by nematodes and cestodes of the following species

#### Nematodes:

**Ascarids:** *Toxocara canis*, *Toxascaris leonina* (adult and late immature forms).

**Hookworms:** *Uncinaria stenocephala*, *Ancylostoma caninum* (adults).

**Whipworms:** *Trichuris vulpis* (adults).

#### Cestodes:

**Tapeworms:** *Echinococcus species* (*E. granulosus*, *E. multilocularis*), *Taenia species* (*T. hydatigena*, *T. pisiformis*, *T. taeniformis*), *Dipylidium caninum* (adult and immature forms).

### 3.3 Contraindications

Do not use simultaneously with piperazine compounds.

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

### 3.4 Special warnings

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*.

Tapeworm infestation is certain to reoccur unless control of intermediate hosts such as fleas, mice, etc. is undertaken.

Tapeworm infestation is unlikely in pups less than 6 weeks of age.

Parasite resistance to any particular class of anthelmintic may develop following frequent, repeated use of an anthelmintic of that class.

Unnecessary use of antiparasitics or use deviating from the instructions given in the SPC may increase the resistance selection pressure and lead to reduced efficacy. The decision to use the product should be based on confirmation of the parasitic species and burden, or of the risk of infection based on its epidemiological features, for each individual animal.

In the absence of risk of co-infection with nematodes or cestodes, a narrow spectrum product should be used.

The possibility that other animals in the same household can be a source of re-infection with nematodes and cestodes should be considered, and these should be treated as necessary with an appropriate product.

### 3.5 Special precautions for use

Special precautions for safe use in the target species:

Not applicable.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

In the interests of good hygiene, persons administering the tablets directly to the dog, or by adding them to the dog's food, should wash their hands afterwards.

Special precautions for the protection of the environment:

Not applicable.

Other precautions:

Echinococcosis represents a hazard for humans. As echinococcosis is a notifiable disease to the World Organisation for Animal Health (OIE), specific guidelines on the treatment and follow-up, and on the safeguard of persons, need to be obtained from the relevant competent authority.

### 3.6 Adverse events

Dogs:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Digestive tract disorders (diarrhoea, emesis)
-----------------------------------------------------------------------------------	-----------------------------------------------

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See also the last section of the package leaflet for respective contact details.

### **3.7 Use during pregnancy, lactation or lay**

#### Pregnancy:

Teratogenic effects attributed to high doses of febantel have been reported in sheep and rats. No studies have been performed in dogs during early pregnancy. Use of the veterinary medicinal product during pregnancy should be in accordance with a benefit risk assessment by the responsible veterinarian. The use is not recommended during the first 4 weeks of pregnancy in dogs. Do not exceed the stated dose when treating pregnant bitches.

### **3.8 Interaction with other medicinal products and other forms of interaction**

Do not use simultaneously with piperazine compounds as the anthelmintic effects of pyrantel and piperazine may be antagonised.

Concurrent use with other cholinergic compounds can lead to toxicity.

### **3.9 Administration routes and dosage**

Oral use.

To ensure a correct dosage, body weight should be determined as accurately as possible.

The recommended dose rates are: 15 mg/kg bodyweight febantel, 5 mg/kg pyrantel (equivalent to 14.4 mg/kg pyrantel embonate) and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 10 kg (22 lbs) bodyweight.

The tablets can be given directly to the dog or disguised in food. No starvation is needed before or after treatment.

Underdosing could result in ineffective use and may favour resistance development.

The advice of a veterinarian should be sought regarding the need for and the frequency of repeat administration.

### **3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)**

The combination of praziquantel, pyrantel embonate and febantel is well tolerated in dogs. In safety studies, a single dose of 5 times the recommended dose or greater gave rise to occasional vomiting.

### **3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance**

Not applicable.

### **3.12 Withdrawal periods**

Not applicable.

## **4. PHARMACOLOGICAL INFORMATION**

#### 4.1 ATCvet code : QP52AA51

#### 4.2 Pharmacodynamics

This veterinary medicinal product contains anthelmintics active against gastrointestinal roundworms and tapeworms. The veterinary medicinal product contains three active substances, as follows:

1. Febantel, a probenzimidazole
2. Pyrantel embonate (pamoate), a tetrahydropyrimidine derivative
3. Praziquantel, a partially hydrogenated pyrazinoisoquinoline derivative

In this fixed combination, pyrantel and febantel act against all relevant nematodes (ascarids, hookworms, and whipworms) in dogs. In particular, the activity spectrum covers *Toxocara canis*, *Toxascaris leonina*, *Uncinaria stenocephala*, *Ancylostoma caninum* and *Trichuris vulpis*. This combination shows synergistic activity in the case of hookworms and febantel is effective against *T. vulpis*.

The spectrum of activity of praziquantel covers all important cestode species in dogs, in particular *Taenia spp.*, *Dipylidium caninum*, *Echinococcus granulosus* and *Echinococcus multilocularis*. Praziquantel acts against all adult and immature forms of these parasites.

Praziquantel is very rapidly absorbed through the parasite's surface and distributed throughout the parasite. Both in vitro and in vivo studies have shown that praziquantel causes severe damage to the parasite integument, resulting in the contraction and paralysis of the parasites. There is an almost instantaneous tetanic contraction of the parasite musculature and a rapid vacuolisation of the syncytial tegument. This rapid contraction has been explained by changes in divalent cation fluxes, especially calcium.

Pyrantel acts as a cholinergic agonist. Its mode of action is to stimulate nicotinic cholinergic receptors of the parasite, induce spastic paralysis of the nematodes and thereby allow removal from the gastrointestinal system by peristalsis.

Within the mammalian system, febantel undergoes ring closure, forming fenbendazole and oxfendazole. It is these chemical entities which exert the anthelmintic effect by inhibition of tubulin polymerisation. Formation of microtubules is thereby prevented, resulting in disruption of structures vital to the normal functioning of the helminth. Glucose uptake in particular is affected, leading to a depletion in cell ATP. The parasite dies upon exhaustion of its energy reserves, which occurs 2 – 3 days later.

#### 4.3 Pharmacokinetics

Perorally administered praziquantel is absorbed almost completely from the intestinal tract. After absorption, the drug is distributed to all organs. Praziquantel is metabolised into inactive forms in the liver and secreted in bile. It is excreted within 24 hours to more than 95% of the administered dosage. Only traces of non-metabolised praziquantel are excreted.

Following administration of the veterinary medicinal product to dogs, peak plasma concentrations of praziquantel were achieved by approximately 2.5 hours.

The pamoate salt of pyrantel has low aqueous solubility, an attribute that reduces absorption from the gut and allows the drug to reach and be effective against parasites in the large intestine. Following absorption, pyrantel pamoate is quickly and almost completely metabolized into inactive metabolites that are excreted rapidly in the urine.

Febantel is absorbed relatively rapidly and metabolized to a number of metabolites including fenbendazole and oxfendazole, which have anthelmintic activity.

Following administration of the veterinary medicinal product to dogs, peak plasma concentrations of fenbendazole and oxfendazole were achieved by approximately 7-9 hours.

## **5. PHARMACEUTICAL PARTICULARS**

### **5.1 Major incompatibilities**

Not applicable.

### **5.2 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 5 years  
Discard any unused divided tablets immediately.

### **5.3 Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

### **5.4 Nature and composition of immediate packaging**

The veterinary medicinal product is presented in either:

Individual strips composed of aluminium foil 30 µm/30 gsm extruded polythene, containing 2, 4, 6, 8, 10, 12, 14, 16, 18 or 20 tablets.

or

Individual blisters composed of 45 µm, soft temper aluminium foil and 25 µm hard temper aluminium foil, containing 2 or 8 tablets.

The strips or blisters are packed into cartons containing either 2,4, 6, 8, 10, 12, 14, 16, 18, 20, 24, 28, 30, 32, 36, 40, 42, 44, 48, 50, 52, 56, 60, 70, 80, 84, 90, 98, 100, 104, 106, 120, 140, 150, 180, 200, 204, 206, 250, 280, 300, 500 or 1000 tablets.

Not all pack sizes may be marketed.

### **5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products**

Medicines should not be disposed of via wastewater.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

## **6. NAME OF THE MARKETING AUTHORISATION HOLDER**

## **7. MARKETING AUTHORISATION NUMBER(S)**

## **8. DATE OF FIRST AUTHORISATION**

**9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS**

{DD/MM/YYYY}

**10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

IE, UK, EE, RO, LU, IT, LT, LV, NL, BE: Veterinary medicinal product not subject to prescription.

AT, CY, CZ, DK, EL, SI, IS, DE, PT, HU, SK: Veterinary medicinal product subject to prescription.

SE,FR Veterinary medicinal product subject to veterinary prescription except for some pack sizes.

Detailed information on this veterinary medicinal product is available in the Union Product Database.

(<https://medicines.health.europa.eu/veterinary>).

**LABELLING****PARTICULARS TO APPEAR ON THE OUTER PACKAGE  
{ CARTON FOR ALL PACK SIZES }****1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Exitel Plus Tablets (IE, IS, BG, EE, NL, HU, PT, LV, SK, LT, SI, ES, EL, AT, CZ, NO)  
Exitel Tablets for Dogs (DE)  
Easimax Plus Tablets. (UK)  
Exitel Plus (DK, CY, RO, LU)  
Exitel vet.150 mg/144 mg/50 mg Tablets (SE)  
FRONTCONTROL WORMER Tablets for Dogs (IT, BE, FR)  
Frontcontrol Vet Tabletti koiralle (FI)

**2. STATEMENT OF ACTIVE SUBSTANCES**

50 mg/tablet Praziquantel, 50 mg/tablet Pyrantel (equivalent to 144 mg pyrantel embonate) and 150 mg/tablet Febantel.

**3. PACKAGE SIZE**

2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 24, 28, 30, 32, 36, 40, 42, 44, 48, 50, 52, 56, 60, 70, 80, 84, 90, 98, 100, 104, 106, 120, 140, 150, 180, 200, 204, 206, 250, 280, 300, 500 and 1000 tablets.

**4. TARGET SPECIES**

Dogs.

**5. INDICATIONS**

For products not subject to veterinary prescription:

Treatment of mixed infections by nematodes and cestodes.

**6. ROUTES OF ADMINISTRATION**

Oral use.

1 tablet per 10 kg bodyweight.

The tablets can be given directly to the dog or disguised in food.

**7. WITHDRAWAL PERIODS****8. EXPIRY DATE**

Exp. {mm/yyyy}

Discard any unused divided tablets immediately.

**9. SPECIAL STORAGE PRECAUTIONS**

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME OF THE MARKETING AUTHORISATION HOLDER**

**14. MARKETING AUTHORISATION NUMBERS**

**15. BATCH NUMBER**

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****{BLISTER FOIL TEXT}****1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Exitel Plus (IE, IS, BG, EE, DE, NL, HU, PT, LV, SK, LT, SI, ES, EL, AT, CZ, NO)


Easimax Plus (UK)

Exitel Plus (DK, CY, RO, LU)

Exitel vet. (SE)

FRONTCONTROL WORMER (IT, BE, FR)

Frontcontrol Vet (FI)

**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

Each tablet contains 50 mg Praziquantel, 50 mg Pyrantel (equivalent to 144 mg pyrantel embonate) and 150 mg Febantel.

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/yyyy}

## PACKAGE LEAFLET

### 1. Name of the veterinary medicinal product

Exitel Plus Tablets For Dogs (IE, IS, BG, EE, NL, HU, PT, LV, SK, LT, SI, ES, EL, AT, CZ, NO)  
Exitel Tablets for Dogs (DE)  
Easimax Plus Tablets For Dogs. (UK)  
Exitel Plus (DK, CY, RO, LU)  
Exitel vet.150 mg/144 mg/50 mg Tablets for dogs (SE)  
FRONTCONTROL WORMER Tablets for Dogs (IT, BE, FR)  
Frontcontrol Vet Tabletti koiralle (FI)

### 2. Composition

Each tablet contains Praziquantel 50 mg, 50 mg Pyrantel (equivalent to 144 mg Pyrantel Embonate) and Febantel 150 mg.

The veterinary medicinal product is a pale yellow tablet with a cross breakline on one side.

The tablets can be divided into 2 or 4 equal parts.

### 3. Target species

Dogs.

### 4. Indications for use

Treatment of mixed infections by nematodes and cestodes of the following species

#### Nematodes:

**Ascarids:** *Toxocara canis*, *Toxascaris leonina* (adult and late immature forms).

**Hookworms:** *Uncinaria stenocephala*, *Ancylostoma caninum* (adults).

**Whipworms:** *Trichuris vulpis* (adults).

#### Cestodes:

**Tapeworms:** *Echinococcus species* (*E. granulosus*, *E. multilocularis*), *Taenia species* (*T. hydatigena*, *T. pisiformis*, *T. taeniformis*), *Dipylidium caninum* (adult and immature forms).

### 5. Contraindications

Do not use simultaneously with piperazine compounds.

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

### 6. Special warnings

#### Special warnings:

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*.

Tapeworm infestation is certain to re-occur unless control of intermediate hosts such as fleas, mice etc. is undertaken.

Tapeworm infestation is unlikely in pups less than 6 weeks of age.

Parasite resistance to any particular class of anthelmintic may develop following frequent, repeated use of an anthelmintic of that class.

Unnecessary use of antiparasitics or use deviating from the instructions given in the SPC may increase the resistance selection pressure and lead to reduced efficacy. The decision to use the product should be based on confirmation of the parasitic species and burden, or of the risk of infection based on its epidemiological features, for each individual animal.

In the absence of risk of co-infection with nematodes or cestodes, a narrow spectrum product should be used.

The possibility that other animals in the same household can be a source of re-infection with nematodes and cestodes should be considered, and these should be treated as necessary with an appropriate product.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

In the interests of good hygiene, persons administering the tablets directly to the dog, or by adding them to the dog's food, should wash their hands afterwards.

For animal treatment only.

Pregnancy:

Teratogenic effects attributed to high doses of febantel have been reported in sheep and rats. No studies have been performed in dogs during early pregnancy. Use of the veterinary medicinal product during pregnancy should be in accordance with a benefit risk assessment by the responsible veterinarian. The use is not recommended during the first 4 weeks of pregnancy in dogs. Do not exceed the stated dose when treating pregnant bitches.

Interactions with other medicinal products and other forms of interaction:

Do not use simultaneously with piperazine compounds as the anthelmintic effects of pyrantel and piperazine may be antagonised.

Concurrent use with other cholinergic compounds (e.g. foxim) can lead to toxicity. If you are uncertain, and your dog uses other veterinary medicinal products, check with a veterinary surgeon or pharmacist.

Overdose:

The combination of praziquantel, pyrantel embonate and febantel is well tolerated in dogs. In safety studies, a single dose of 5 times the recommended dose or greater gave rise to occasional vomiting.

Other precautions:

Echinococcosis represents a hazard for humans. As echinococcosis is a notifiable disease to the World Organisation for Animal Health (OIE), specific guidelines on the treatment and follow-up, and on the safeguard of persons, need to be obtained from the relevant competent authority.

## **7. Adverse events**

Dogs:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Digestive tract disorders (diarrhoea, emesis)
--------------------------------------------------------------------------------	-----------------------------------------------

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system.

**8. Dosage for each species, routes and method of administration**

Oral use.

The recommended dose rates are: 15 mg/kg bodyweight febantel, 5 mg/kg pyrantel (equivalent to 14.4 mg/kg pyrantel embonate) and 5 mg/kg praziquantel.

1 tablet per 10 kg (22 lbs) bodyweight. The tablets can be given directly to the dog or disguised in food. No starvation is needed before, or after, treatment.

The tablet can be divided in two or four equal parts.

Dosage table:

Bodyweight (kg)	Tablets
0.5 - 2.5	¼
2.6-5.0	½
5.1-10.0	1
10.1-15.0	1½
15.1-20.0	2
20.1-25.0	2½
25.1-30.0	3
30.1-35.0	3½
35.1-40.0	4
>40.1	1 tablet per 10 kg

Underdosing could result in ineffective use and may favour resistance development.

The advice of a veterinarian should be sought regarding the need for and the frequency of repeat administration.

**9. Advice on correct administration**

To ensure a correct dosage, body weight should be determined as accurately as possible .

**10. Withdrawal periods**

Not applicable.

**11. Special storage precautions**

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month.

Discard any unused divided tablets immediately.

**12. Special precautions for disposal**

Medicines should not be disposed of via wastewater.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

**13. Classification of veterinary medicinal products**

IE, UK, EE, RO, LU, IT, LT, LV, NL, BE: Veterinary medicinal product not subject to prescription.  
AT, CY, CZ, DK, EL, SI, IS, DE, PT, HU, SK: Veterinary medicinal product subject to prescription.  
SE, FR: Veterinary medicinal product subject to veterinary prescription except for some pack sizes.

**14. Marketing authorisation numbers and pack sizes**

The veterinary medicinal product is presented in either:

Individual strips composed of aluminium foil 30 µm/30 gsm extruded polythene, containing 2, 4, 6, 8, 10, 12, 14, 16, 18 or 20 tablets.

or

Individual blisters composed of 45 µm, soft temper aluminium foil and 25 µm hard temper aluminium foil, containing 2 or 8 tablets.

The strips or blisters are packed into cartons containing either 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 24, 28, 30, 32, 36, 40, 42, 44, 48, 50, 52, 56, 60, 70, 80, 84, 90, 98, 100, 104, 106, 120, 140, 150, 180, 200, 204, 206, 250, 280, 300, 500 or 1000 tablets.

Not all pack sizes may be marketed.

**15. Date on which the package leaflet was last revised**

{DD/MM/YYYY}

Detailed information on this veterinary medicinal product is available in the Union Product Database. (<https://medicines.health.europa.eu/veterinary>).

**16. Contact details**

Marketing authorisation holder and contact details to report suspected adverse reactions:

Manufacturer responsible for batch release:

Chanelle Pharmaceuticals Manufacturing Ltd  
Loughrea  
Co. Galway  
Ireland

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

Local representatives and contact details to report suspected adverse reactions: