

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Felisecto Plus 15 mg/2.5 mg spot-on solution for cats \leq 2.5 kg
Felisecto Plus 30 mg/5 mg spot-on solution for cats > 2.5–5 kg
Felisecto Plus 60 mg/10 mg spot-on solution for cats > 5–10 kg

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each unit dose (pipette) delivers:

Active substances:

Felisecto Plus spot-on solution	unit dose (ml)	selamectin (mg)	sarolaner (mg)
Cats \leq 2.5 kg	0.25	15	2.5
Cats > 2.5–5 kg	0.5	30	5
Cats > 5–10 kg	1	60	10

Excipients:

Qualitative composition of excipients and other constituents	Quantitative composition if that information is essential for proper administration of the veterinary medicinal product
Butylhydroxytoluene	0.2 mg/ml
Dipropyleneglycol monomethyl ether	
Isopropyl alcohol	

Clear, colourless to yellow spot-on solution.

3. CLINICAL INFORMATION

3.1 Target species

Cats.

3.2 Indications for use for each target species

For cats with, or at risk from, mixed parasitic infestations by ticks and fleas, lice, mites, gastrointestinal nematodes or heartworm. The veterinary medicinal product is exclusively indicated when use against ticks and one or more of the other target parasites is indicated at the same time.

Ectoparasites:

- For the treatment and prevention of flea infestations (*Ctenocephalides* spp.). The veterinary medicinal product has immediate and persistent flea killing activity against new infestations for 5 weeks. The product kills adult fleas before they lay eggs for 5 weeks. Through its ovicidal and larvicidal action, the veterinary medicinal product may aid in the control of existing environmental flea infestations in areas to which the animal has access.
- The product can be used as part of a treatment strategy for flea allergy dermatitis (FAD).
- Treatment of tick infestations. The veterinary medicinal product has immediate and persistent acaricidal effect for 5 weeks against *Ixodes ricinus* and *Ixodes hexagonus*, and 4 weeks against *Dermacentor reticulatus* and *Rhipicephalus sanguineus*.
- Treatment of ear mites (*Otodectes cynotis*).
- Treatment of biting lice infestations (*Felicola subrostratus*).

Ticks must attach to the host and commence feeding in order to be exposed to sarolaner.

Nematodes:

- Treatment of adult roundworms (*Toxocara cati*) and adult intestinal hookworms (*Ancylostoma tubaeforme*).
- Prevention of heartworm disease caused by *Dirofilaria immitis* with monthly administration.

3.3 Contraindications

Do not use in cats that are suffering from concomitant disease, or are debilitated and underweight (for size and age).

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

3.4 Special warnings

It is recommended, in accordance with good veterinary practice, that all animals 6 months of age or more, living in countries where a vector exists should be tested for existing adult heartworm infections before beginning preventive use with the veterinary medicinal product.

This veterinary medicinal product is not effective against adult *D. immitis*. The administration to animals with adult heartworm infection did not pose safety concerns.

Whilst not routinely indicated, the potential benefits of performing periodic testing for heartworm infection in individual cases should be considered by the responsible veterinarian.

Ticks need to start feeding on the host to become exposed to sarolaner; therefore, the transmission of infectious tick-borne diseases cannot be excluded.

3.5 Special precautions for use

Special precautions for safe use in the target species:

Use of this veterinary medicinal product is indicated in cats aged at least 8 weeks old and weighing at least 1.25 kg bodyweight.

This veterinary medicinal product is to be applied to the skin surface only. Do not administer orally or parenterally.

Do not apply when the animal's hair coat is wet.

For ear mite treatment, do not apply directly to the ear canal.

It is important to apply the dose as indicated to prevent the animal from licking and ingesting the product. If significant ingestion occurs, transient gastrointestinal effects such as hypersalivation, emesis, soft faeces or reduced food consumption may be observed and should normally resolve without treatment.

Keep treated animals away from fires and other sources of ignition for at least 30 minutes or until the hair coat is dry.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

The veterinary medicinal product is harmful after ingestion. Keep the veterinary medicinal product in the original packaging until use, in order to prevent children from getting direct access to the veterinary medicinal product. Used pipettes should be disposed of immediately. In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

The veterinary medicinal product may cause eye irritation. Avoid eye contact including hand-to-eye-contact. Avoid direct contact with treated animals until the application area is dry. Wash hands after use and wash off any veterinary medicinal product in contact with the skin immediately with soap and water. In case of accidental eye exposure, flush the eyes immediately with water, seek medical advice immediately and show the package leaflet or the label to the physician.

Children are not allowed to play with treated cats for 4 hours after treatment. It is recommended to treat animals in the evening. On the day of treatment, treated animals should not be permitted to sleep in the same bed as their owner, especially children.

People with sensitive skin or known allergy to veterinary medicinal products of this type should handle the veterinary medicinal product with caution.

The veterinary medicinal product is highly flammable. Keep away from heat, sparks, open flame or other sources of ignition.

Special precautions for the protection of the environment:

The veterinary medicinal product should not enter water courses as this may be dangerous for fish and other aquatic organisms.

3.6 Adverse events

Cats:

Uncommon (1 to 10 animals / 1,000 animals treated):	Application site pruritus ¹ , application site alopecia ² Erythema ² Drooling ²
Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Convulsions ³ , ataxia ³ Vomiting ³ , diarrhoea ³

¹ mild and transient.

² mild to moderate.

³ mostly transient.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Pregnancy, lactation and fertility:

The safety of the veterinary medicinal product has not been established during pregnancy and lactation or in animals intended for breeding. However, selamectin is considered safe for use in breeding, pregnant and lactating cats. While the safety of sarolaner has not been evaluated in breeding, pregnant or lactating cats, laboratory studies with sarolaner in rats and rabbits have not produced any evidence of teratogenic effects. Use only according to the benefit-risk assessment by the responsible veterinarian.

3.8 Interaction with other medicinal products and other forms of interaction

During clinical field testing, no interactions between this veterinary medicinal product and routinely used veterinary medicinal products were observed.

3.9 Administration routes and dosage

Spot-on use.

The veterinary medicinal product should be administered as a single spot-on (topical) application in accordance with the following table (corresponding to a minimum of 6 mg/kg selamectin and 1 mg/kg sarolaner).

Bodyweight of cat (kg)	Pipette content (ml)	Strength and number of pipettes to be administered		
		15 mg/2.5 mg (yellow cap)	30 mg/5 mg (orange cap)	60 mg/10 mg (green cap)
≤ 2.5	0.25	1		
> 2.5–5	0.5		1	
> 5–10	1			1
> 10	Appropriate combination of pipettes			


Method and route of administration

Apply topically to the skin at the base of the neck in front of the shoulder blades. The pipette should be removed from the protective package immediately prior to administration.

Holding the pipette upright, firmly depress the cap to puncture the applicator seal, then remove the cap.

Part the hair at the base of the cat's neck in front of the shoulder blades to expose a small area of skin. Apply the tip of the pipette directly to the skin, without massaging.

Squeeze the pipette firmly 3–4 times to empty the contents in one spot. Avoid contact between the product and your fingers.



Transient cosmetic effects may occur at the application site such as temporary clumping or spiking of the hair, greasiness or dry white deposits, which normally resolve within 24 hours after product application. These effects do not affect the safety or efficacy of the veterinary medicinal product.

Treatment schedule

Fleas and ticks

For optimal control of tick and flea infestations, the veterinary medicinal product should be administered at monthly intervals and continued throughout the flea and/or tick season based on local epidemiological situations.

Following administration of the veterinary medicinal product, the adult fleas on the animal are killed within 24 hours, no viable eggs are produced, and larvae (found only in the environment) are also killed. This stops flea reproduction, breaks the flea lifecycle and may aid in the control of existing environmental flea infestations in areas to which the animal has access.

Prevention of heartworm disease

The veterinary medicinal product may be administered year-round or at least within one month of the animal's first exposure to mosquitoes and monthly thereafter until the end of the mosquito season. The final dose must be given within one month after the last exposure to mosquitoes. If a dose is missed and a monthly interval between dosing is exceeded, then immediate administration of the veterinary medicinal product and resumption of monthly dosing will minimise the opportunity for the development of adult heartworms. When replacing another heartworm preventive veterinary medicinal product in a heartworm disease prevention programme, the first dose of the veterinary medicinal product must be given within a month of the last dose of the former veterinary medicinal product.

Treatment of roundworm and hookworm infections

A single dose of the veterinary medicinal product should be administered. The need for and frequency of re-treatment should be in accordance with the advice of the prescribing veterinarian.

Treatment of biting lice

A single dose of the veterinary medicinal product should be administered.

Treatment of ear mites

A single dose of the veterinary medicinal product should be administered. Seek further veterinary examination 30 days after treatment to determine whether a second administration is necessary.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

No clinically significant adverse reactions were observed in kittens from 8 weeks of age treated with up to 5 times the maximum recommended dose of the veterinary medicinal product for up to 8 consecutive treatments at 28 day intervals, apart from a single cat administered 5 times the maximum dose that displayed transient hypersensitivity to touch, piloerection, mydriasis and mild tremor which resolved without treatment.

After accidental ingestion of a full product dose, transient gastrointestinal effects such as salivation, soft faeces, emesis, and reduced food consumption may occur; however, these should resolve without treatment.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Not applicable.

4. PHARMACOLOGICAL INFORMATION

4.1 ATCvet code: QP54AA55.

4.2 Pharmacodynamics

Selamectin is a semi-synthetic compound of the avermectin class. Selamectin paralyzes and/or kills a wide range of invertebrate parasites through interference with their chloride channel conductance

causing disruption of normal neurotransmission. This inhibits the electrical activity of nerve cells in nematodes and muscle cells in arthropods leading to their paralysis and/or death.

Selamectin has adulticidal, ovicidal and larvicidal activity against fleas. Therefore, it effectively breaks the flea life cycle by killing adults (on the animal), preventing the hatching of eggs (on the animal and in its environment) and by killing larvae (environment only). Debris from selamectin-treated pets kills flea eggs and larvae not previously exposed to selamectin and thus may aid in the control of existing environmental flea infestations in areas to which the animal has access. Selamectin is active against adult fleas (*Ctenocephalides* spp.) as well as mites (*Otodectes cynotis*), lice (*Felicola subrostratus*) and gastrointestinal nematodes (*Toxocara cati*, *Ancylostoma tubaeforme*). Activity has also been demonstrated against heartworm (*D. immitis*) larvae.

For fleas, the onset of efficacy is within 24 hours for 5 weeks after product application.

Sarolaner is an acaricide and insecticide belonging to the isoxazoline family. The primary target of action of sarolaner in insects and acarines is functional blockade of ligand-gated chloride channels (GABA-receptors and glutamate-receptors). Sarolaner blocks GABA- and glutamate-gated chloride channels in the central nervous system of insects and acarines. Disruption of these receptors by sarolaner prevents the uptake of chloride ions by GABA and glutamate gated ion channels, thus resulting in increased nerve stimulation and death of the target parasite. Sarolaner exhibits higher functional potency to block insect/acarine receptors compared to mammalian receptors. Sarolaner does not interact with known insecticidal binding sites of nicotinic or other GABAergic insecticides such as neonicotinoids, fiproles, milbemycins, avermectins, and cyclodienes. Sarolaner is active against adult fleas (*Ctenocephalides* spp.) as well as several tick species such as *Dermacentor reticulatus*, *Ixodes hexagonus*, *Ixodes ricinus*, and *Rhipicephalus sanguineus*.

For ticks (*I. ricinus*), the onset of efficacy is within 24 hours of attachment for one month after product application.

4.3 Pharmacokinetics

Following topical administration of the Felisecto Plus both selamectin and sarolaner are well absorbed with bioavailability mean values of 40.5% and 57.9%, respectively and distribute systemically. In cats, selamectin and sarolaner are low clearance compounds with long half-life values, 12.5 days and 41.5 days respectively, following topical administration.

In cats the primary route of selamectin elimination is in faeces and the majority is parent compound. Identification of selamectin metabolites in faeces indicated that metabolic clearance also contributes to the elimination. The primary route of elimination for sarolaner is biliary elimination of parent sarolaner, with contributions by metabolic clearance.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

None known.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 30 months.

5.3 Special precautions for storage

Store below 30 °C.

Do not remove the pipette from the blister until ready to use.

5.4 Nature and composition of immediate packaging

Translucent polypropylene unit dose pipettes individually packed in aluminium and aluminium/PVC blisters.

Felisecto Plus 15 mg/2.5 mg spot-on solution for cats ≤ 2.5 kg: 0.25 ml per pipette.

Felisecto Plus 30 mg/5 mg spot-on solution for cats > 2.5 –5 kg: 0.5 ml per pipette.

Felisecto Plus 60 mg/10 mg spot-on solution for cats > 5 –10 kg: 1 ml per pipette.

The veterinary medicinal product is available in packs of three pipettes (all pipette sizes) placed in cardboard boxes.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

Zoetis Belgium

7. MARKETING AUTHORISATION NUMBER(S)

EU/2/19/238/001-003

8. DATE OF FIRST AUTHORISATION

Date of first authorisation: 26/04/2019.

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

{DD/MM/YYYY}

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

ANNEX II

OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

None.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

CARDBOARD BOX

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Felisecto Plus 15 mg/2.5 mg spot-on solution \leq 2.5 kg
Felisecto Plus 30 mg/5 mg spot-on solution > 2.5–5 kg
Felisecto Plus 60 mg/10 mg spot-on solution > 5–10 kg

2. STATEMENT OF ACTIVE SUBSTANCES

15 mg selamectin/2.5 mg sarolaner/pipette
30 mg selamectin/5 mg sarolaner/pipette
60 mg selamectin/10 mg sarolaner/pipette

3. PACKAGE SIZE

3 x 0.25 ml
3 x 0.5 ml
3 x 1 ml


4. TARGET SPECIES

Cats.

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Spot-on use.



7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {mm/yyyy}

9. SPECIAL STORAGE PRECAUTIONS

Store below 30 °C.

Do not remove the pipette from the blister until ready to use.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Zoetis Belgium

14. MARKETING AUTHORISATION NUMBERS

EU/2/19/238/001 (3 x 0.25 ml)

EU/2/19/238/002 (3 x 0.5 ml)

EU/2/19/238/003 (3 x 1 ml)

15. BATCH NUMBER

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

BLISTER

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Felisecto Plus



2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

≤ 2.5 kg

> 2.5–5 kg

> 5–10 kg

selamectin/sarolaner

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

PIPETTE

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Felisecto Plus



2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

≤ 2.5 kg
> 2.5–5 kg
> 5–10 kg

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Felisecto Plus 15 mg/2.5 mg spot-on solution for cats \leq 2.5 kg
Felisecto Plus 30 mg/5 mg spot-on solution for cats > 2.5–5 kg
Felisecto Plus 60 mg/10 mg spot-on solution for cats > 5–10 kg

2. Composition

Each unit dose (pipette) delivers:

Active substances:

Felisecto Plus spot-on solution	unit dose (ml)	selamectin (mg)	sarolaner (mg)
Cats \leq 2.5 kg	0.25	15	2.5
Cats > 2.5–5 kg	0.5	30	5
Cats > 5–10 kg	1	60	10

Excipients:

Butylhydroxytoluene 0.2 mg/ml.

Clear, colourless to yellow spot-on solution.

3. Target species

Cats.

4. Indications for use

For cats with, or at risk from, mixed parasitic infestations by ticks and fleas, lice, mites, gastrointestinal nematodes or heartworm. The veterinary medicinal product is exclusively indicated when use against ticks and one or more of the other target parasites is indicated at the same time.

Ectoparasites:

- For the treatment and prevention of flea infestations (*Ctenocephalides* spp.). The veterinary medicinal product has immediate and persistent flea killing activity against new infestations for 5 weeks. The product kills adult fleas before they lay eggs for 5 weeks. Through its ovicidal and larvicidal action, the veterinary medicinal product may aid in the control of existing environmental flea infestations in areas to which the animal has access.
- The product can be used as part of a treatment strategy for flea allergy dermatitis (FAD).
- Treatment of tick infestations. The veterinary medicinal product has immediate and persistent acaricidal effect for 5 weeks against *Ixodes ricinus* and *Ixodes hexagonus*, and 4 weeks against *Dermacentor reticulatus* and *Rhipicephalus sanguineus*.
- Treatment of ear mites (*Otodectes cynotis*).
- Treatment of biting lice infestations (*Felicola subrostratus*).

Ticks must attach to the host and commence feeding in order to be exposed to sarolaner.

Nematodes:

- Treatment of adult roundworms (*Toxocara cati*) and adult intestinal hookworms (*Ancylostoma tubaeforme*).
- Prevention of heartworm disease caused by *Dirofilaria immitis* with monthly administration.

5. Contraindications

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

Do not use in cats that are suffering from concomitant disease, or are debilitated and underweight (for size and age).

6. Special warningsSpecial warnings:

It is recommended, in accordance with good veterinary practice, that all animals 6 months of age or more, living in countries where a vector exists should be tested for existing adult heartworm infections before beginning preventive use with the veterinary medicinal product.

This veterinary medicinal product is not effective against adult *D. immitis*. The administration to animals with adult heartworm infection did not pose safety concerns.

Whilst not routinely indicated, the potential benefits of performing periodic testing for heartworm infection in individual cases should be considered by the responsible veterinarian.

Ticks need to start feeding on the host to become exposed to sarolaner; therefore, the transmission of infectious tick-borne diseases cannot be excluded.

Special precautions for safe use in the target species:

Use of this veterinary medicinal product is indicated in cats aged at least 8 weeks old and weighing at least 1.25 kg bodyweight.

This veterinary medicinal product is to be applied to the skin surface only. Do not administer orally or parenterally.

Do not apply when the animal's hair coat is wet.

For ear mite treatment, do not apply directly to the ear canal.

It is important to apply the dose as indicated to prevent the animal from licking and ingesting the product. If significant ingestion occurs, transient gastrointestinal effects such as hypersalivation, emesis, soft faeces or reduced food consumption may be observed and should normally resolve without treatment.

Keep treated animals away from fires and other sources of ignition for at least 30 minutes or until the hair coat is dry.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

The veterinary medicinal product is harmful after ingestion. Keep the veterinary medicinal product in the original packaging until use, in order to prevent children from getting direct access to the veterinary medicinal product. Used pipettes should be disposed of immediately. In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

The veterinary medicinal product may cause eye irritation. Avoid eye contact including hand-to-eye-contact. Avoid direct contact with treated animals until the application area is dry. Wash hands after

use and wash off any veterinary medicinal product in contact with the skin immediately with soap and water. In case of accidental eye exposure, flush the eyes immediately with water, seek medical advice immediately and show the package leaflet or the label to the physician. People with sensitive skin or known allergy to veterinary medicinal products of this type should handle the veterinary medicinal product with caution.

Children are not allowed to play with treated cats for 4 hours after treatment. It is recommended to treat animals in the evening. On the day of treatment, treated animals should not be permitted to sleep in the same bed as their owner, especially children.

The veterinary medicinal product is highly flammable. Keep away from heat, sparks, open flame or other sources of ignition.

Special precautions for the protection of the environment:

The veterinary medicinal product should not enter water courses as this may be dangerous for fish and other aquatic organisms.

Pregnancy, lactation and fertility:

The safety of the veterinary medicinal product has not been established during pregnancy and lactation or in animals intended for breeding. However, selamectin is considered safe for use in breeding, pregnant and lactating cats. While the safety of sarolaner has not been evaluated in breeding, pregnant or lactating cats, laboratory studies with sarolaner in rats and rabbits have not produced any evidence of teratogenic effects. Use only according to the benefit-risk assessment by the responsible veterinarian.

Interaction with other medicinal products and other forms of interactions:

During clinical field testing, no interactions between this veterinary medicinal product and routinely used veterinary medicinal products were observed.

Overdose:

No clinically significant adverse reactions were observed in kittens from 8 weeks of age treated with up to 5 times the maximum recommended dose of the veterinary medicinal product for up to 8 consecutive treatments at 28 day intervals, apart from a single cat administered 5 times the maximum dose that displayed transient hypersensitivity to touch, piloerection, mydriasis and mild tremor which resolved without treatment.

After accidental ingestion of a full product dose, transient gastrointestinal effects such as salivation, soft faeces, emesis, and reduced food consumption may occur, however, these should resolve without treatment.

Major incompatibilities:

None known.

7. Adverse events

Cats:

Uncommon (1 to 10 animals / 1,000 animals treated):	Application site pruritus (itching) ¹ , application site alopecia (hair loss) ² Erythema (redness) ² Drooling ²
Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Convulsions ³ , ataxia (incoordination) ³ Vomiting ³ , diarrhoea ³

¹ mild and transient.

² mild to moderate.

³ mostly transient.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: {national system details}.

8. Dosage for each species, routes and method of administration

Spot-on use.

The veterinary medicinal product should be administered as a single spot-on (topical) application in accordance with the following table (corresponding to a minimum of 6 mg/kg selamectin and 1 mg/kg sarolaner).

Bodyweight of cat (kg)	Pipette content (ml)	Strength and number of pipettes to be administered		
		15 mg/2.5 mg (yellow cap)	30 mg/5 mg (orange cap)	60 mg/10 mg (green cap)
≤ 2.5	0.25	1		
> 2.5–5	0.5		1	
> 5–10	1			1
> 10	Appropriate combination of pipettes			

Fleas and ticks

For optimal control of tick and flea infestations, the veterinary medicinal product should be administered at monthly intervals and continued throughout the flea and/or tick season based on local epidemiological situations.

Following administration of the veterinary medicinal product, the adult fleas on the animal are killed within 24 hours, no viable eggs are produced, and larvae (found only in the environment) are also killed. This stops flea reproduction, breaks the flea lifecycle and may aid in the control of existing environmental flea infestations in areas to which the animal has access.

Prevention of heartworm disease

The veterinary medicinal product may be administered year-round or at least within one month of the animal's first exposure to mosquitoes and monthly thereafter until the end of the mosquito season. The final dose must be given within one month after the last exposure to mosquitoes. If a dose is missed and a monthly interval between dosing is exceeded, then immediate administration of the veterinary medicinal product and resumption of monthly dosing will minimise the opportunity for the development of adult heartworms. When replacing another heartworm preventive veterinary medicinal product in a heartworm disease prevention programme, the first dose of the veterinary medicinal product must be given within a month of the last dose of the former veterinary medicinal product.

Treatment of roundworm and hookworm infections

A single dose of the veterinary medicinal product should be administered. The need for and frequency of re-treatment should be in accordance with the advice of the prescribing veterinarian.

Treatment of biting lice

A single dose of the veterinary medicinal product should be administered.

Treatment of ear mites

A single dose of the veterinary medicinal product should be administered. Seek further veterinary examination 30 days after treatment to determine whether a second administration is necessary.

9. Advice on correct administration

This veterinary medicinal product is to be applied to the skin surface only. Do not administer orally or parenterally.

Do not apply when the animal's hair coat is wet.

For ear mite treatment, do not apply directly to the ear canal.

It is important to apply the dose as indicated to prevent the animal from licking and ingesting the product. If significant ingestion occurs, transient gastrointestinal effects such as hypersalivation, emesis, soft faeces or reduced food consumption may be observed and should normally resolve without treatment.

Apply topically to the skin at the base of the neck in front of the shoulder blades. The pipette should be removed from the protective package immediately prior to administration.

Holding the pipette upright, firmly depress the cap to puncture the applicator seal, then remove the cap.

Part the hair at the base of the cat's neck in front of the shoulder blades to expose a small area of skin. Apply the tip of the pipette directly to the skin without massaging.

Squeeze the pipette firmly 3–4 times to empty the contents in one spot. Avoid contact between the product and your fingers.



Transient cosmetic effects may occur at the application site such as temporary clumping or spiking of the hair, greasiness or dry white deposits, which normally resolve within 24 hours after product application. These effects do not affect the safety or efficacy of the veterinary medicinal product.

10. Withdrawal periods

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.

Store below 30 °C.

Do not remove the pipette from the blister until ready to use.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton, blister and pipette after Exp. The expiry date refers to the last day of that month.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack sizes

EU/2/19/238/001-003

The veterinary medicinal product is available in packs of three pipettes (all pipette sizes).

15. Date on which the package leaflet was last revised

{DD/MM/YYYY}

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

16. Contact details

Marketing authorisation holder and manufacturer responsible for batch release:

Zoetis Belgium
Rue Laid Burniat 1
1348 Louvain-La-Neuve
Belgium

Local representatives and contact details to report suspected adverse reactions:

België/Belgique/Belgien
Zoetis Belgium
Mercuriusstraat 20
BE-1930 Zaventem
Tél/Tel: +32 (0) 800 99 189

Lietuva
Zoetis Belgium
Mercuriusstraat 20
1930 Zaventem
Belgija
Tel: +370 610 05088

Република България

Zoetis Belgium
Rue Laid Burniat 1
1348 Louvain-La-Neuve
Белгия
Тел: +359 888 51 30 30

Česká republika

Zoetis Česká republika, s.r.o.
náměstí 14. října 642/17
CZ 150 00 Praha
Tel: +420 257 101 111

Danmark

Zoetis Animal Health ApS
Øster Alle 48
DK-2100 København
Tlf: +45 70 20 73 05
adr.scandinavia@zoetis.com

Deutschland

Zoetis Deutschland GmbH
Schellingstr. 1
DE-10785 Berlin
Tel: +49 30 2020 0049
tierarzneimittelsicherheit@zoetis.com

Eesti

Zoetis Belgium
Mercuriusstraat 20
1930 Zaventem
Belgia
Tel: +370 610 05088

Ελλάδα

Zoetis Hellas S.A.
Φραγκοκκλησιάς 7, Μαρούσι
EL-15125 Αττική
Τηλ: +30 210 6791900

España

Zoetis Spain, S.L.
Parque Empresarial Vía Norte Edificio nº1,
c/ Quintanavides nº13
ES-28050 Madrid
Tel: +34 91 4191900

France

Zoetis France
10 rue Raymond David
FR-92240 Malakoff
Tél: +33 (0)800 73 00 65

Luxembourg/Luxemburg

Zoetis Belgium
Mercuriusstraat 20
1930 Zaventem
Belsch
Tél/Tel: +32 (2) 746 80 11

Magyarország

Zoetis Hungary Kft.
Csörsz u. 41.
HU-1124 Budapest
Tel.: +36 1 224 5200

Malta

Agrimed Limited
Mdina Road, Zebbug ZBG 9016,
MT
Tel: +356 21 465 797

Nederland

Zoetis B.V.
Rivium Westlaan 74
NL-2909 LD Capelle aan den IJssel
Tel: +31 (0)10 714 0900

Norge

Zoetis Animal Health ApS
Øster Alle 48
DK-2100 København
Danmark
Tlf: +47 23 29 86 80
adr.scandinavia@zoetis.com

Österreich

Zoetis Österreich GmbH
Floridsdorfer Hauptstr. 1
AT-1210 Wien
Tel: +43 (0)1 2701100 100
tierarzneimittelsicherheit@zoetis.com

Polska

Zoetis Polska Sp. z o.o.
ul. Postępu 17B
PL - 02-676 Warszawa
Tel.: +48 22 2234800

Portugal

Zoetis Portugal Lda.
Lagoas Park, Edifício 10
PT-2740-271 Porto Salvo
Tel: +351 21 042 72 00

Hrvatska

Zoetis B.V.
Podružnica Zagreb za promidžbu
Petra Hektorovića 2
HR-10000 Zagreb
Tel: +385 1 6441 462

Ireland

Zoetis Belgium S.A. (Irish Branch)
2nd Floor, Building 10,
Cherrywood Business Park,
Loughlinstown,
Co. Dublin,
IE – Dublin D18 T3Y1
Tel: +353 (0) 1 256 9800

Ísland

Zoetis Animal Health ApS
Øster Alle 48
DK-2100 København
Danmörku
Sími: +45 70 20 73 05
adr.scandinavia@zoetis.com

Italia

Zoetis Italia S.r.l.
Via Andrea Doria 41M,
IT-00192 Roma
Tel: +39 06 3366 8111

Κύπρος

Zoetis Hellas S.A.
Φραγκοκκλησιάς 7, Μαρούσι
15125, Αττική
Ελλάδα
Τηλ: +30 210 6791900

Latvija

Zoetis Belgium
Mercuriusstraat 20
1930 Zaventem
Belģija
Tel: +370 610 05088

România

Zoetis România S.R.L.
Expo Business Park, 54A Aviator Popișteanu,
Clădirea 2, Etaj 1-3, Sector 1,
București, 012095 - RO
Tel: +40785019479

Slovenija

Zoetis B.V.
Podružnica Zagreb za promidžbu
Petra Hektorovića 2,
10000 Zagreb,
Hrvaška
Tel: +385 1 6441 462

Slovenská republika

Zoetis Česká republika, s.r.o.
náměstí 14. října 642/17
150 00 Praha
Česká republika
Tel: +420 257 101 111

Suomi/Finland

Zoetis Finland Oy
Bulevardi 21 / SPACES
FI-00180 Helsinki/Helsingfors
Suomi/Finland
Puh/Tel: +358 10 336 7000
laaketurva@zoetis.com

Sverige

Zoetis Animal Health ApS
Øster Alle 48
DK-2100 Köpenhamn
Danmark
Tel: +46 (0) 76 760 0677
adr.scandinavia@zoetis.com

United Kingdom (Northern Ireland)

Zoetis Belgium S.A. (Irish Branch)
2nd Floor, Building 10,
Cherrywood Business Park,
Loughlinstown,
Co. Dublin,
IE – Dublin D18 T3Y1
Tel: +353 (0) 1 256 9800

17. Other information

Selamectin is a semi-synthetic compound of the avermectin class. Selamectin has adulticidal, ovicidal and larvicidal activity against fleas. Therefore, it effectively breaks the flea life cycle by killing adults (on the animal), preventing the hatching of eggs (on the animal and in its environment) and by killing larvae (environment only). Debris from selamectin-treated pets kills flea eggs and larvae not

previously exposed to selamectin and thus may aid in the control of existing environmental flea infestations in areas to which the animal has access. Selamectin is active against adult fleas (*Ctenocephalides* spp.) as well as mites (*Otodectes cynotis*), lice (*Felicola subrostratus*) and gastrointestinal nematodes (*Toxocara cati*, *Ancylostoma tubaeforme*). Activity has also been demonstrated against heartworm (*D. immitis*) larvae.

For fleas, the onset of efficacy is within 24 hours for 5 weeks after product application.

Sarolaner is an acaricide and insecticide belonging to the isoxazoline family. Sarolaner is active against adult fleas (*Ctenocephalides* spp.) as well as several tick species such as *Dermacentor reticulatus*, *Ixodes hexagonus*, *Ixodes ricinus*, and *Rhipicephalus sanguineus*.

For ticks (*I. ricinus*), the onset of efficacy is within 24 hours of attachment for one month after product application.