

*[Version 9,03/2022] corr. 11/2022*

## **ANNEX I**

### **SUMMARY OF PRODUCT CHARACTERISTICS**

October 2023

1b-spc-pl

## 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Buprelab Vet 0.3 mg/ml solution for injection for dogs and cats  
SE, FI  
Buprelab 0.3 mg/ml solution for injection for dogs and cats  
ES, LV, LT, LU, BE, CY, FR, HU, RO  
Buprelab  
EE

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml contains:

### Active substance:

Buprenorphine	0.3 mg
(equivalent to buprenorphine hydrochloride)	0.324 mg)

### Excipients:

Qualitative composition of excipients and other constituents	Quantitative composition if that information is essential for proper administration of the veterinary medicinal product
Chlorocresol	1.35 mg
Glucose anhydrous	
Hydrochloric acid	
Water for injections	

Clear and colourless solution for injection.

## 3. CLINICAL INFORMATION

### 3.1 Target species

Dogs and cats.

### 3.2 Indications for use for each target species

Dogs:

- Post-operative analgesia.
- Potentiation of the sedative effects of centrally acting agents.

Cats:

- Post-operative analgesia.

### 3.3 Contraindications

Do not administer by the intrathecal or peridural route.

Do not use pre-operatively for Caesarian section (see Section 3.7).

Do not use in animals with known hypersensitivity to the active substance or to any of the excipients.

**3.4 Special warnings**

None.

**3.5 Special precautions for use**

Special precautions for safe use in the target species:

Buprenorphine may cause respiratory depression and as with other opioid drugs, care should be taken when treating animals with impaired respiratory function or animals that are receiving drugs that can cause respiratory depression.

In case of renal, cardiac or hepatic dysfunction or shock, there may be greater risk associated with the use of the product. The benefit/risk assessment for using the product should be made by the attending vet. Safety has not been fully evaluated in clinically compromised cats.

Buprenorphine should be used with caution in animals with impaired liver function, especially biliary tract disease, as the substance is metabolised by the liver and its intensity and duration of action may be affected in such animals.

The safety of buprenorphine has not been demonstrated in animals less than 7 weeks of age, therefore, use in such animals should be based on the benefit/risk assessment of the veterinarian.

Repeat administration earlier than the recommended repeat interval suggested in Section 3.9 is not recommended.

Long-term safety of buprenorphine in cats has not been investigated beyond 5 consecutive days of administration.

The effect of an opioid on head injury is dependent on the type and severity of the injury and the respiratory support supplied. The veterinary medicinal product should be used in accordance with the benefit/risk assessment of the attending veterinarian.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Wash hands/affected area thoroughly after any accidental spillage. As buprenorphine has opioid-like activity, care should be taken to avoid self-injection. In case of accidental self-injection or ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Following eye contamination or skin contact, wash thoroughly with cold running water. Seek medical advice if irritation persists.

Special precautions for the protection of the environment:

Not applicable.

**3.6 Adverse events**

Dogs:

Uncommon (1 to 10 animals / 1,000 animals treated):	Ptyalism, bradycardia, hypothermia, agitation, dehydration and miosis. Respiratory depression. <sup>1</sup>
Rare (1 to 10 animals / 10,000 animals	Hypertension, tachycardia.

October 2023

1b-spc-pl

treated):	Sedation <sup>2</sup>
Very rare ( $<1$ animal / 10,000 animals treated, including isolated reports):	Discomfort , injection site pain. <sup>3</sup>

1- Refer to section 3.5 Special precautions for safe use in the target species.

2- When used to provide analgesia, sedation may occur at dose levels higher than those recommended.

3- Resulting in vocalisation.

Cats:

Common (1 to 10 animals / 100 animals treated):	Mydriasis. Behavioural disorders (restlessness, purring and excessive rubbing). <sup>4</sup>
Uncommon (1 to 10 animals / 1,000 animals treated):	Respiratory depression. <sup>1</sup>
Rare (1 to 10 animals / 10,000 animals treated):	Sedation. <sup>2</sup>
Very rare ( $<1$ animal / 10,000 animals treated, including isolated reports):	Discomfort , injection site pain. <sup>3</sup>

1- Refer to section 3.5 Special precautions for safe use in the target species.

2- When used to provide analgesia, sedation may occur at dose levels higher than those recommended.

3- Resulting in vocalisation.

4- Will usually resolve within 24 hours.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

### 3.7 Use during pregnancy, lactation or lay

#### Pregnancy:

Laboratory studies in rats have not produced any evidence of a teratogenic effect. However, these studies have shown post-implantation losses and early foetal deaths. These may have resulted from a reduction in parental body condition during gestation and in post-natal care owing to sedation of the mothers.

**The safety of the veterinary medicinal product has not been established during pregnancy. Use only according to the benefit/risk assessment by the responsible veterinarian.**

The veterinary medicinal product should not be used pre-operatively in cases of Caesarean section, due to the risk of respiratory depression in the offspring periparturiently, and should only be used post-operatively with special care (see below).

Lactation:

Studies in lactating rats have shown that, after intramuscular administration of buprenorphine, concentrations of unchanged buprenorphine in the milk equalled or exceeded that in the plasma. As it is likely that buprenorphine will be excreted in the milk of other species, use is not recommended during lactation.

**3.8 Interaction with other medicinal products and other forms of interaction**

Buprenorphine may cause some drowsiness, which may be potentiated by other centrally acting agents, including tranquillisers, sedatives and hypnotics.

There is evidence in humans to indicate that therapeutic doses of buprenorphine do not reduce the analgesic efficacy of standard doses of an opioid agonist, and that when buprenorphine is employed within the normal therapeutic range, standard doses of opioid agonist may be administered before the effects of the former have ended without compromising analgesia. However, it is recommended that buprenorphine is not used in conjunction with morphine or other opioid-type analgesics, e.g. etorphine, fentanyl, pethidine, methadone, papaveretum or butorphanol.

Buprenorphine has been used with acepromazine, alphaxalone/alphadalone, atropine, dexmedetomidine, halothane, isoflurane, ketamine, medetomidine, propofol, sevoflurane, thiopentone and xylazine. When used in combination with sedatives, depressive effects on heart rate and respiration may be augmented.

**3.9 Administration routes and dosage**

Intramuscular or intravenous use

Species	Administration routes	Post-operative analgesia	Potential of the sedative
Dogs	intramuscular or intravenous	10 - 20 micrograms buprenorphine/kg of b.w. (equivalent to 0.03-0.06 ml of veterinary medicinal product per kg de b.w.).  Repeat, if necessary, after 3-4 hours with 10 micrograms of buprenorphine per kg de b.w. or after 5-6 hours with 20 micrograms of buprenorphine per kg de b.w.	10-20 micrograms of buprenorphine per kg of b.w. (equivalent to 0.03-0.06 ml of veterinary medicinal product per kg de b-w.)
Cats	intramuscular or intravenous	10 - 20 micrograms buprenorphine/kg of b.w. (equivalent to 0.03-0.06 ml of veterinary medicinal product per kg de b.w.), repeated, if	-----

	necessary, once, after 1-2 hours.	
--	-----------------------------------	--

While sedative effects are present by 15 minutes after administration, analgesic activity becomes apparent after approximately 30 minutes. To ensure that analgesia is present during surgery and immediately on recovery, the product should be administered preoperatively as part of premedication.

When administered for potentiation of sedation or as part of premedication, the dose of other centrally-acting agents, such as acepromazine or medetomidine, should be reduced. The reduction will depend on the degree of sedation required, the individual animal, the type of other agents included in premedication and how anaesthesia is to be induced and maintained. It may also be possible to reduce the amount of inhalational anaesthetic used.

Animals administered opioids possessing sedative and analgesic properties may show variable responses. Therefore, the response of individual animals should be monitored, and subsequent doses should be adjusted accordingly. In some cases, repeat doses may fail to provide additional analgesia. In these cases, consideration should be given to using a suitable injectable NSAID.

An appropriately graduated syringe must be used to allow accurate dosing. The stopper cannot be breached more than 44 times.

### 3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

In cases of overdosage, supportive measures should be instituted, and, if appropriate, naloxone or respiratory stimulants may be used.

When administered at overdose to dogs, buprenorphine may cause lethargy. At very high doses, bradycardia and miosis may be observed.

Naloxone may be of benefit in reversing reduced respiratory rate and respiratory stimulants such as Doxapram are also effective in man. Because of the prolonged duration of effect of buprenorphine in comparison to such drugs, they may need to be administered repeatedly or by continuous infusion. Volunteer studies in man have indicated that opiate antagonists may not fully reverse the effects of buprenorphine.

In toxicological studies of buprenorphine hydrochloride in dogs, biliary hyperplasia was observed after oral administration for one year at dose levels of 3.5mg/kg/day and above. Biliary hyperplasia was not observed following daily intramuscular injection of dose levels up to 2.5mg/kg/day for 3 months. This is well in excess of any clinical dose regimen in the dog.

Please also refer to Sections 3.5 and 3.6 of this SPC.

### 3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

To be completed at the national phase.

### 3.12 Withdrawal periods

Not applicable.

October 2023

1b-spc-pl

## 4. PHARMACOLOGICAL INFORMATION

### 4.1 ATCvet code: QN02AE01

### 4.2 Pharmacodynamics

In summary, buprenorphine is a potent, long-acting analgesic acting at opiate receptors in the central nervous system. Buprenorphine can potentiate the effects of other centrally acting agents, but unlike most opiates, buprenorphine has at clinical doses, only a limited sedative effect of its own.

Buprenorphine exerts its analgesic effect via high affinity binding to various subclasses of opiate receptors, particularly  $\mu$ , in the central nervous system. At clinical dose levels for analgesia, buprenorphine binds to opiate receptors with high affinity and high receptor avidity, such that its dissociation from the receptor site is slow, as demonstrated in *in vitro* studies. This unique property of buprenorphine could account for its longer duration of activity when compared to morphine. In circumstances where excessive opiate agonist is already bound to opiate receptors, buprenorphine can exert a narcotic antagonistic activity as a consequence of its high-affinity opiate receptor binding, such that an antagonistic effect on morphine equivalent to naloxone has been demonstrated.

Buprenorphine has little effect on gastro-intestinal motility.

### 4.3 Pharmacokinetics

When given parenterally, the veterinary medicinal product may be administered by intramuscular or intravenous injection.

Buprenorphine is rapidly absorbed after intramuscular injection in various animal species and man. The substance is highly lipophilic and the volume of distribution in body compartments is large. Pharmacological effects (e.g. mydriasis) may occur within minutes of administration and signs of sedation normally appear by 15 minutes. Analgesic effects appear around 30 minutes with peak effects usually being observed at about 1 – 1.5 hours.

Following intravenous administration to dogs at a 20 $\mu$ g/kg dose, the mean terminal half-life was 9 hours and the mean clearance was 24 ml/kg/min, however, there is considerable inter-dog variability in pharmacokinetic parameters.

Following intramuscular administration to cats, the mean terminal half-life was 6.3 hours, and the clearance was 23 mL/kg/min; however, there was considerable inter-cat variability in pharmacokinetic parameters.

Combined pharmacokinetic and pharmacodynamic studies in cats have demonstrated a marked hysteresis between plasma concentration and analgesic effect. Plasma concentrations of buprenorphine should not be used to formulate individual animal dosage regimens, which should be determined by monitoring the patient's response.

The major route of excretion in all species except the rabbit (where urinary excretion predominates) is the faeces. Buprenorphine undergoes N-dealkylation and glucuronide conjugation by the intestinal wall and the liver and its metabolites are excreted via the bile into the gastro-intestinal tract.

October 2023

1b-spc-pl

In tissue distribution studies carried out in rats and rhesus monkeys the highest concentrations of drug-related material were observed in liver, lung and brain. Peak levels occurred rapidly and declined to low levels by 24 hours after dosing.

Protein binding studies in rats have shown that buprenorphine is highly bound to plasma proteins, principally to alpha and beta globulins.

## **5. PHARMACEUTICAL PARTICULARS**

### **5.1 Major incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

### **5.2 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

Shelf life after first opening the immediate packaging: 28 days

### **5.3 Special precautions for storage**

Keep the vial in the outer carton in order to protect from light.

### **5.4 Nature and composition of immediate packaging**

Amber glass vial type I with chlorobutyl stopper and aluminium cap.

#### Pack sizes:

Box with 1 vial of 10 ml

Box with 5 vials of 10 ml

Box with 10 vials of 10 ml

Not all pack sizes may be marketed.

### **5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products**

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

## **6. NAME OF THE MARKETING AUTHORISATION HOLDER**

Labiana Life Sciences, S.A.

## **7. MARKETING AUTHORISATION NUMBER(S)**

XXXX



**October 2023**

**1b-spc-pl**

**8. DATE OF FIRST AUTHORISATION**

Date of first authorisation: {DD/MM/YYYY}.

**9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT  
CHARACTERISTICS**

{MM/YYYY}

**10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the [Union Product Database](https://medicines.health.europa.eu/veterinary)  
(<https://medicines.health.europa.eu/veterinary>).