

1. NAME OF THE VETERINARY MEDICINAL PRODUCT
For all CMSs except Spain

KENOCIDIN
Chlorhexidine digluconate 5mg/g, Teat dip solution for cattle (dairy)

For Spain

KENOCIDIN
5mg/g, Teat dip solution for cattle (dairy)
Chlorhexidine digluconate

2. STATEMENT OF THE ACTIVE SUBSTANCE AND OTHER INGREDIENT
Chlorhexidine digluconate, 5 mg/g
(Equivalent to chlorhexidine 2.815 mg/g)
Glycerol 51 mg/g
Allantoin 1mg/g

3. ROUTE OF ADMINISTRATION
Teat use

FOR ANIMAL TREATMENT ONLY.

4. PACKAGE SIZES
1L, 5L, 10L, 20L, 25L, 60L, 200L. Not all pack sizes may be marketed.

5. MARKETING AUTHORISATION NUMBER

UK: Vm 22136/4002 - For UK transport emergencies only phone 01865407333
IE: VPA number: 10792/003/001

6. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND
OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR
BATCH RELEASE, IF DIFFERENT

CIDLINES NV
Waterpoortstraat 2
8900 Ieper
Belgium
Tel. +32 (0) 57 21 78 77
Fax. +32 (0) 57 21 78 79
Mail: info@cidlines.com

7. TARGET SPECIES
Cattle (dairy).

8. DISTRIBUTION CATEGORY
UK: AMV-GSL

IE: LM

KEEP OUT OF THE REACH AND SIGHT OF CHILDREN.

9. SPECIAL STORAGE PRECAUTIONS

Keep container tightly closed.

Protect from frost

If the veterinary medicinal product has frozen, thaw in a warm place and shake well before use

Protect from light

Do not use after the expiry date stated on the label

The 200 litre container should not be returned for re-filling.

10. WITHDRAWAL PERIOD

Meat & offal: zero days

Milk: zero hours

11. ADVERSE REACTIONS

None known.

12. SPECIAL WARNING(S)

Ensure udder and teats are clean and dry before the next milking.

Can be used during pregnancy and lactation.

Chlorhexidine can be inactivated by anionic and nonionic surfactants (eg soaps, even natural) or inorganic anions, so do not mix with tap water, other chemicals, disinfectants and other products for the teat and udder care

This veterinary medicinal product is for teat use, significant absorption does not occur.

Special precautions for use in animals

- **For external use only.**
- **Use for the treatment of teats with cutaneous lesions may delay the wound healing process. It is recommended to discontinue the treatment until the lesions are healed.**
- **The presence of organic matter (pus, blood, etc.) may limit the action of the disinfectant chlorhexidine “**
- **If the temperature is below freezing, allow teats to air dry before letting cows outside**
- **If signs of disease appear, consult a veterinary surgeon**
- **Allow product to dry before exposing the cows to wet (rainy), cold or windy conditions.**

Operator Warnings

- **Avoid contact with eyes. If splashed in the eye, rinse with clean running water and seek medical advice.**
- **In case of ingestion, drink large quantities of water and seek medical attention immediately.**
- **Keep away from food and animal feed.**
- **Wash hands after use.**
 - People with known hypersensitivity to chlorhexidine should avoid contact with the veterinary medicinal product.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIAL, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements. The product should not enter water courses as this may be dangerous for fish and other aquatic organisms.

UK Only: Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority. To dispose of unused product to land you must have an authorisation under the Groundwater Regulations 1998.

14. INDICATION

Teat disinfection as a part of a prevention strategy for mastitis in lactating dairy cows. For the maintenance of good teat skin and teat end condition.

15. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

The product is ready to use as a post-milking teat dip, applied up to two times per day.

Use at least 5ml per cow per application.

Dip the teats immediately after milking each cow. Ensure that the teat is completely covered to three quarters of its length.

The dip cup should be replenished as necessary.

If a common dip cup is used for application, a fresh solution should always be used at each milking. The dip cup should be emptied, cleaned and rinsed after each milking session or when the cup becomes contaminated during milking. Do not pour the remaining solution from the dip cup back into the original container. Do not use the product for cleaning and/or sanitizing milking equipment.

16. CONTRAINDICATIONS

Do not use in case of known hypersensitivity to chlorhexidine or any of the excipients.

Shelf-life after first opening the immediate packaging: 6months.

Discard date: ____ / ____ / ____

UK:

