

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

GALLIMUNE 302 ND+IB+EDS emulsion for injection

GALLIMUNE ND+IB+EDS emulsion for injection (Belgium, Denmark, Luxembourg, Sweden)

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 0.3 ml dose of vaccine contains:

Active substances:

Inactivated Newcastle disease virus, Ulster 2C strain, at least..... 50 PD₅₀¹

Inactivated infectious bronchitis virus, Mass41 strain, at least18 HI.U

Inactivated egg drop syndrome virus (EDS76), V127 strain, at least180 HI.U

The concentrations are expressed by the antibody titre obtained during the potency test. One unit (U) corresponding to an antibody titre of 1.

HI: haemagglutination inhibiting

¹ Minimum protective dose according to monograph 0870 of Ph. Eur.

Adjuvant:

Paraffin oil.....170 to 186 mg

Excipients:

Qualitative composition of excipients and other constituents	Quantitative composition if that information is essential for proper administration of the veterinary medicinal product
Thiomersal	30 µg
Formaldehyde	at most 43.2 µg
Ester of fatty acids and of polyols	
Ester of fatty acids and of ethoxylated polyols	
Water for injections	

Whitish homogeneous emulsion.

3. CLINICAL INFORMATION

3.1 Target species

Chickens (breeder and layer pullets).

3.2 Indications for use for each target species

Booster immunisation of breeder and layer pullets after vaccination with live vaccines against:

- Newcastle disease virus in order to reduce egg drop linked to Newcastle disease infection,
- Infectious bronchitis virus in order to reduce egg drop linked to infectious bronchitis infection caused by the Mass41 strain,

Active immunisation of breeder and layer pullets in order to reduce egg drop linked to infection with egg drop syndrome virus (EDS76) without priming.

Onset of immunity: 4 weeks after vaccination.
Duration of immunity: one laying period.

3.3 Contraindications

None.

3.4 Special warnings

Vaccinate healthy animals only.

3.5 Special precautions for use

Special precautions for safe use in the target species:

Not applicable.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

To the user:

This veterinary medicinal product contains mineral oil. Accidental injection/self-injection may result in severe pain and swelling, particularly if injected into a joint or finger, and in rare cases could result in the loss of the affected finger if prompt medical attention is not given. If you are accidentally injected with this product, seek prompt medical advice even if only a very small amount is injected and take the package insert with you. If pain persists for more than 12 hours after medical examination, seek medical advice again.

To the physician:

This veterinary medicinal product contains mineral oil. Even if small amounts have been injected, accidental injection with this product can cause intense swelling, which may, for example, result in ischaemic necrosis and even the loss of a digit. Expert, PROMPT, surgical attention is required and may necessitate early incision and irrigation of the injected area, especially where there is involvement of finger pulp or tendon.

Special precautions for the protection of the environment:

Not applicable.

3.6 Adverse events

Chickens:

Very common (> 1 animal / 10 animals treated):	Abnormal histology ¹
---	---------------------------------

¹ In clinical studies, lesions at the injection site linked to the oily adjuvant were observed histologically three weeks after injections in 87 % of cases, e.g. small quantities of oily residues and occasional aseptic micro-abscesses. No palpable reactions were observed.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Do not use in birds in lay and within 2 weeks before the start of the laying period.

3.8 Interaction with other medicinal products and other forms of interaction

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

3.9 Administration routes and dosage

Administer one dose (0,3 ml) by intramuscular route from the age of 18 weeks and at least 4 weeks after the priming with live vaccines against Newcastle disease (strain Hitchner B1 or VG/GA-AVINEW) and infectious bronchitis (strain Mass H120).

Shake well before use.

Apply usual aseptic procedures.

Do not use syringes with natural rubber or butyl elastomer pistons.

Equipment including needles and syringes must be sterile before use.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

In addition to the adverse events mentioned in paragraph “Adverse events”, transitory apathy and slight oedema at injection site may occur after the administration of a double dose of vaccine.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

[To be completed nationally]

<Any person intending to manufacture, import, possess, distribute, sell, supply and use this veterinary medicinal product must consult the relevant Member State’s Competent Authority on the current vaccination policies as these activities may be prohibited in a Member State on the whole or part of its territory pursuant to national legislation.>

<Official control authority batch release is required for this product.>

3.12 Withdrawal periods

Zero days.

4. IMMUNOLOGICAL INFORMATION

4.1 ATCvet code: QI01AA13

Inactivated vaccine in oily adjuvant against Newcastle disease, Infectious bronchitis and egg drop syndrome (EDS76).

The vaccine stimulates active immunity of breeder and layer pullets against egg drop syndrome (EDS76) (without priming), Newcastle disease and infectious bronchitis, subsequent to priming with live vaccines against these diseases.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

Do not mix with any other veterinary medicinal product.

5.2 Shelf life

Shelf-life of the veterinary medicinal product as packaged for sale: 18 months.

Shelf life after first opening the immediate packaging: use immediately.

5.3 Special precautions for storage

Store and transport refrigerated (2 °C – 8 °C).

Protect from light.

Do not freeze.

5.4 Nature and composition of immediate packaging

Nature of primary packaging elements:

- Polypropylene bottle
- Nitrile elastomer closure
- Aluminium cap

Sales presentations:

- 150-ml (500-dose) bottle.
- 150-ml (500-dose) bottle, box of 10 bottles.
- 300-ml (1 000-dose) bottle.
- 300-ml (1 000-dose) bottle, box of 10 bottles.

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

[To be completed nationally]

7. MARKETING AUTHORISATION NUMBER(S)

[To be completed nationally]

8. DATE OF FIRST AUTHORISATION

[To be completed nationally]

{DD/MM/YYYY}

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

[To be completed nationally]
{MM/YYYY}

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

500 doses, Bottle of 150 ml
10 x 500 doses, 10 Bottles of 150 ml
1000 doses, Bottle of 300 ml
10 x 1000 doses, 10 Bottles of 300 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

GALLIMUNE 302 ND+IB+EDS emulsion for injection
GALLIMUNE ND+IB+EDS emulsion for injection (Belgium, Denmark, Luxembourg, Sweden)

2. STATEMENT OF ACTIVE SUBSTANCES

Each 0.3 ml dose contains:

Inactivated Newcastle disease virus, Ulster 2C strain..... ≥ 50 PD₅₀
Inactivated infectious bronchitis virus, Mass41 strain ≥ 18 HI.U
Inactivated egg drop syndrome virus (EDS76), V127 strain..... ≥ 180 HI.U

3. PACKAGE SIZE

0.3 ml/d
500 doses (150 ml)
10 x 500 doses (10 x 150 ml)
1 000 doses (300 ml)
10 x 1 000 doses (10 x 300 ml)

4. TARGET SPECIES

Chickens (breeder and layer pullets).

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

For intramuscular use.
Shake well before use.

7. WITHDRAWAL PERIODS

Withdrawal period: zero days.

8. EXPIRY DATE

Exp.: {dd/mm/yyyy}

Once opened use immediately.

9. SPECIAL STORAGE PRECAUTIONS

Store and transport refrigerated.

Protect from light.

Do not freeze.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

[To be completed nationally]

14. MARKETING AUTHORISATION NUMBERS

[To be completed nationally]

15. BATCH NUMBER

Lot {number}

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

500 doses, 150 ml
1000 doses, 300 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

GALLIMUNE 302 ND+IB+EDS emulsion for injection
GALLIMUNE ND+IB+EDS emulsion for injection (Belgium, Denmark, Luxembourg, Sweden)

2. STATEMENT OF ACTIVE SUBSTANCES

Per dose:

Inactivated:

- Newcastle disease virus, Ulster 2C strain ≥ 50 PD₅₀,
- infectious bronchitis virus, Mass41 strain..... ≥ 18 HI.U,
- egg drop syndrome virus (EDS76), V127 strain..... ≥ 180 HI.U,

0.3 ml/d

500 doses

1 000 doses

3. TARGET SPECIES

Chickens (breeder and layer pullets)

4. ROUTES OF ADMINISTRATION

For intramuscular use.

Shake well before use.

Read the package leaflet before use.

5. WITHDRAWAL PERIODS

Withdrawal period: zero days.

6. EXPIRY DATE

Exp. {dd/mm/yyyy}

Once opened use immediately.

7. SPECIAL STORAGE PRECAUTIONS

Store and transport refrigerated.

Protect from light.

Do not freeze.

8. NAME OF THE MARKETING AUTHORISATION HOLDER

[To be completed nationally]

9. BATCH NUMBER

Lot {number}

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

GALLIMUNE 302 ND+IB+EDS emulsion for injection

GALLIMUNE ND+IB+EDS emulsion for injection (Belgium, Denmark, Luxembourg, Sweden)

2. Composition

Each 0.3 ml dose contains:

Inactivated Newcastle disease virus, Ulster 2C strain..... $\geq 50 \text{ PD}_{50}$ ¹

Inactivated infectious bronchitis virus, Mass41 strain $\geq 18 \text{ HI.U}$

Inactivated egg drop syndrome virus (EDS76), V127 strain..... $\geq 180 \text{ HI.U}$

Thiomersal..... $30 \mu\text{g}$

Formaldehyde..... $\leq 43.2 \mu\text{g}$

Paraffin oil (as adjuvant) 170 to 186 mg

The concentrations are expressed by the antibody titre obtained during the potency test. One unit (U) corresponding to an antibody titre of 1.

HI: haemagglutination inhibiting

¹ Minimum protective dose according to monograph 0870 of Ph. Eur.

Whitish homogeneous emulsion.

3. Target species

Chickens (breeder and layer pullets).

4. Indications for use

Booster immunisation of breeder and layer pullets after vaccination with live vaccines against:

- Newcastle disease virus
- infectious bronchitis virus

Active immunisation of breeder and layer pullets in order to reduce egg drop linked to infection with egg drop syndrome virus (EDS76) without priming.

Onset of immunity: 4 weeks after vaccination.

Duration of immunity: one laying period.

5. Contraindications

None.

6. Special warning(s)

Vaccinate healthy animals only.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

To the user:

This veterinary medicinal product contains mineral oil. Accidental injection/self injection may result in severe pain and swelling, particularly if injected into a joint or finger, and in rare cases could result in the loss of the affected finger if prompt medical attention is not given.

If you are accidentally injected with this product, seek prompt medical advice even if only a very small amount is injected and take the package insert with you.

If pain persists for more than 12 hours after medical examination, seek medical advice again.

To the physician:

This veterinary medicinal product contains mineral oil. Even if small amounts have been injected, accidental injection with this product can cause intense swelling, which may, for example, result in ischaemic necrosis and even the loss of a digit. Expert, PROMPT, surgical attention is required and may necessitate early incision and irrigation of the injected area, especially where there is involvement of finger pulp or tendon.

Laying birds:

Do not use in birds in lay and within 2 weeks before the start of the laying period.

Interaction with other medicinal products and other forms of interaction:

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Overdose:

In addition to the adverse events mentioned in paragraph “Adverse events”, transitory apathy and slight oedema at injection site may occur after the administration of a double dose of vaccine.

Major incompatibilities:

Do not mix with any other veterinary medicinal product.

7. Adverse events

Chickens:

Very common (> 1 animal / 10 animals treated):

Abnormal histology¹

¹ In clinical studies, lesions at the injection site linked to the oily adjuvant were observed histologically three weeks after injections in 87 % of cases, e.g. small quantities of oily residues and occasional aseptic micro-abscesses. No palpable reactions were observed.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: {national system details}

8. Dosage for each species, routes and method of administration

Administer one dose (0.3 ml) by intramuscular route from the age of 18 weeks and at least 4 weeks after the priming with live vaccines against Newcastle disease (strain Hitchner B1 or VG/GA-AVINEW) and infectious bronchitis (strain Mass H120).

9. Advice on correct administration

- Shake well before use.
- Apply usual aseptic procedures.
- Do not use syringes with natural rubber or butyl elastomer pistons.
- Equipment including needles and syringes must be sterile before use.

10. Withdrawal periods

Zero days.

11. Special storage precautions

Keep out of the reach and sight of children.

Store and transport refrigerated (2 °C – 8 °C).

Protect from light.

Do not freeze.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after “Exp.”.

Shelf life after first opening the immediate packaging: use immediately.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

14. Marketing authorisation numbers and pack sizes

To be completed nationally.

Pack sizes:

150-ml (500-dose) bottle.

150-ml (500-dose) bottle, box of 10 bottles.

300-ml (1 000-dose) bottle.

300-ml (1 000-dose) bottle, box of 10 bottles.

Not all pack sizes may be marketed.

15. Date on which the package leaflet was last approved

{MM/YYYY}

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

16. Contact details

Marketing authorisation holder and contact details to report suspected adverse reactions:
To be completed nationally.

Manufacturer responsible for the batch release:
Boehringer Ingelheim Animal Health France SCS
Laboratoire Porte des Alpes
Rue de l'Aviation
69800 Saint-Priest
France

Local representatives and contact details to report suspected adverse reactions:
To be completed nationally.

17. Other information

Inactivated vaccine in oily adjuvant against Newcastle disease, infectious bronchitis and egg drop syndrome (EDS76).