

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Anthelmin Plus Flavour Tablets for dogs (BE, IE, IT, NL, UK(NI))
Dehinel Plus Flavour Tablets for dogs (BG)
Zikyall Sabor Tablets for dogs (ES, PT)
Anthelmin vet 150 mg/144 mg/50 mg tablets for dogs (FI)

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains:

Active substances:

Praziquantel	50 mg
Pyrantel embonate	144 mg
Febantel	150 mg

Excipients:

Qualitative composition of excipients and other constituents
Lactose monohydrate
Maize starch
Povidone K-30
Sodium laurilsulfate
Microcrystalline cellulose (E460)
Colloidal anhydrous silica
Magnesium stearate (E572)
Meat flavour

Yellow coloured, round, biconvex tablets with visible darker spots, cross-scored on one side.

3. CLINICAL INFORMATION

3.1 Target species

Dogs (weighing less than 17.5 kg).

3.2 Indications for use for each target species

For the treatment of mixed infections with the following roundworms and tapeworms in adult dogs and puppies:

Nematodes

Ascarids: *Toxocara canis*, *Toxascaris leonina* (late immature forms and mature forms)

Hookworms: *Uncinaria stenocephala*, *Ancylostoma caninum* (adults)

Cestodes

Tapeworms: *Taenia* spp., *Dipylidium caninum*

3.3 Contraindications

Do not use simultaneously with piperazine compounds.

Do not exceed the stated dosage when treating pregnant bitches.

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.
Do not use in dogs younger than 2 weeks of age and/or weighing less than 2 kg.

3.4 Special warnings

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*. Tapeworm infestation is certain to re-occur unless control of intermediate hosts such as fleas, mice etc. is undertaken.

Parasite resistance to any particular class of anthelmintic may develop following frequent, repeated use of an anthelmintic of that class.

3.5 Special precautions for use

Special precautions for safe use in the target species:

Any part-used tablets should be discarded.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

In the interests of good hygiene, persons administering the tablet directly to a dog or by adding it to the dog's food, should wash their hands afterwards.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Special precautions for the protection of the environment:

Not applicable.

3.6 Adverse events

Dogs:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Loose stool, diarrhoea, vomiting.*
---	------------------------------------

*Transient.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Pregnancy:

Do not use in bitches during the first two-thirds of pregnancy.

Consult a veterinary surgeon before treating pregnant animals for roundworms.

Lactation:

Can be used during lactation (see Sections 3.3 and 3.9).

3.8 Interaction with other medicinal products and other forms of interaction

Do not combine with piperazine as the anthelmintic effects of pyrantel and piperazine (used in many worming products for dogs) may be antagonized.

Concurrent use with other cholinergic compounds can lead to toxicity.

3.9 Administration routes and dosage

For oral use.

Dosage

The recommended dose rates are: 15 mg/kg bodyweight febantel, 14.4 mg/kg pyrantel and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 10 kg bodyweight.

Tablets may be halved/quartered to allow accuracy of dosing.

Administration and Duration of Treatment

The tablet(s) can be given directly to the dog or disguised in food. No restriction of access to food is required either before or after administration of the veterinary medicinal product.

To ensure a correct dosage, body weight should be determined as accurately as possible.

Puppies may be wormed with this veterinary medicinal product from 2 weeks of age and every 2 weeks until 12 weeks of age. Thereafter they should be treated at 3 monthly intervals until 6 months of age.

For the control of *Toxocara*, nursing bitches should be dosed 2 weeks after giving birth and every 2 weeks until weaning.

For adult dogs, a single dose should be used. The advice of a veterinarian should be sought regarding the need for and frequency of repeat treatment.

In the event of a heavy roundworm infestation, a repeat dose should be given after 14 days.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

Benzimidazoles possess wide safety margin. Pyrantel is not absorbed systematically to any extent. Praziquantel also has a wide safety margin, of up to five times the recommended dose.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Not applicable.

4. PHARMACOLOGICAL INFORMATION

4.1 ATCvet code: QP52AC55

4.2 Pharmacodynamics

The veterinary medicinal product contains anthelmintics active against roundworms and tapeworms. The veterinary medicinal product contains three active substances: febantel, pyrantel embonate (pamoate) and praziquantel, a partially hydrogenated pyrazino-isoquinoline derivative used widely as an anthelmintic for both human and veterinary use. Pyrantel acts as a cholinergic agonist. Its mode of action is to stimulate nicotinic cholinergic receptors of the parasite, induce spastic paralysis and thereby allow removal from the gastro-intestinal (GI) system by peristalsis.

With the mammalian system febantel undergoes ring closure forming fenbendazole and oxfendazole. It is these chemical entities which exert the anthelmintic effect by inhibition of tubulin polymerization.

Formation of microtubules is thereby prevented, resulting in disruption to structures vital to the normal functioning of the helminth. Glucose uptake, in particular, is affected, leading to depletion in cell ATP. The parasite dies upon exhaustion of its energy reserves, which occurs 2 – 3 days later.

Praziquantel is very rapidly absorbed and distributed throughout the parasite. Both *in vitro* and *in vivo* studies have shown that praziquantel causes severe damage to the parasite integument, resulting in contraction and paralysis. There is an almost instantaneous tetanic contraction of the parasite musculature and a rapid vacuolisation of the syncytial tegument. This rapid contraction has been explained by changes in divalent cation fluxes, especially calcium.

In this fixed combination veterinary medicinal product pyrantel and febantel act synergistically against all relevant nematodes (ascarids and hookworms) in dogs. In particular, the activity spectrum covers *Toxocara canis*, *Toxascaris leonina*, *Uncinaria stenocephala* and *Ancylostoma caninum*. The spectrum of activity of praziquantel covers also cestode species in dogs, in particular all *Taenia* spp. and *Dipylidium caninum*. Praziquantel acts against adult and immature forms of these parasites.

4.3 Pharmacokinetics

Perorally administered praziquantel is absorbed almost completely from the intestinal tract. After absorption, the drug is distributed to all organs. Praziquantel is metabolized into inactive forms in the liver and secreted in bile. It is excreted within 24 hours to more than 95% of the administered dosage. Only traces of non-metabolised praziquantel are excreted.

The pamoate salt of pyrantel has low aqueous solubility, an attribute that reduces absorption from the gut and allows the drug to reach and be effective against parasites in the large intestine. Because of the low systemic absorption of pyrantel pamoate, there is very little danger of adverse reactions/toxicity in the host. Following absorption, pyrantel pamoate is quickly and almost completely metabolised into inactive metabolites that are excreted rapidly in the urine.

Febantel is absorbed relatively rapidly and metabolised to a number of metabolites including fenbendazole and oxfendazole, which have anthelmintic activity.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

Not applicable.

5.2 Shelf life

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.

5.3 Special precautions for storage

This medicinal product does not require any special storage conditions.

5.4 Nature and composition of immediate packaging

OPA/Al/PVC-Al blister: 2 tablets (1 blister with 2 tablets), in a box.

OPA/Al/PVC-Al blister: 4 tablets (2 blisters with 2 tablets), in a box.

OPA/Al/PVC-Al blister: 10 tablets (1 blister with 10 tablets), in a box.

OPA/Al/PVC-Al blister: 30 tablets (3 blisters with 10 tablets), in a box.

OPA/Al/PVC-Al blister: 50 tablets (5 blisters with 10 tablets), in a box.

OPA/Al/PVC-Al blister: 100 tablets (10 blisters with 10 tablets), in a box.

OPA/Al/PVC-Al blister: 300 tablets (30 blisters with 10 tablets), in a box.

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

KRKA d.d, Novo mesto

7. MARKETING AUTHORISATION NUMBER(S)

8. DATE OF FIRST AUTHORISATION

Date of first authorisation: DD/MM/YYYY

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

{MM/YYYY}

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription. (PT, ES)

Veterinary medicinal product not subject to prescription. (BE, BG, IE, IT, NL,UK(NI))

Veterinary medicinal product subject to prescription except for some pack sizes. FI (packaging for 24, 30,50, 60, 100, 102 tbl)

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

ANNEX III

LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Cardboard box

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Anthelmin Plus Flavour Tablets f (BE, IE, IT, NL, UK(NI))
Dehinel Plus Flavour Tablets (BG)
Zikyall Sabor Tablets (ES, PT)
Anthelmin vet tablets (FI)

2. STATEMENT OF ACTIVE SUBSTANCES

Each tablet contains:

Praziquantel	50 mg
Pyrantel embonate	144 mg
Febantel	150 mg

3. PACKAGE SIZE

2 tbl
4 tbl
10 tbl
30 tbl
50 tbl
100 tbl
300 tbl

4. TARGET SPECIES

For dogs (weighing less than 17.5 kg)

5. INDICATIONS**6. ROUTES OF ADMINISTRATION**

Oral use.

The tablets can be divided into 2 or 4 equal parts.

Dosage:

Body weight	Tablets
2.0 - 2.5 kg	¼
2.6 - 5 kg	½
5.1 - 7.5 kg	¾
7.6 - 10 kg	1

10.1 - 12.5 kg	1 ¼
12.6 - 15 kg	1 ½
15.1 - 17.5 kg	1 ¾
17.6 - 20 kg	2
20.1 - 22.5 kg	2 ¼
22.6 - 25 kg	2 ½
25.1 - 27.5 kg	2 ¾
27.6 - 30 kg	3

Not for use in dogs younger than 2 weeks of age and/or weighing less than 2 kg.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {mm/yyyy}

9. SPECIAL STORAGE PRECAUTIONS

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

KRKA, d.d., Novo mesto

14. MARKETING AUTHORISATION NUMBERS

15. BATCH NUMBER

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON *SMALL IMMEDIATE PACKAGING UNITS*

Blister

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Anthelmin Plus Flavour (BE, IE, IT, NL, UK(NI))
Dehinel Plus Flavour (BG)
Zikyall Sabor (ES, PT)
Anthelmin vet (FI)



2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

50 mg/144 mg/150 mg

3. BATCH NUMBER

Lot {number}

3. EXPIRY DATE

Exp. {mm/yyyy}

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Anthelmin Plus Flavour Tablets for dogs (BE, IE, IT, NL, UK (NI))

Dehinel Plus Flavour Tablets for dogs (BG)

Zikyall Sabor Tablets for dogs (ES, PT)

Anthelmin vet 150 mg/144 mg/50 mg tablets for dogs (FI)

2. Composition

Each tablet contains:

Praziquantel	50 mg
Pyrantel embonate	144 mg
Febantel	150 mg

Yellow coloured, round, biconvex tablets with visible darker spots, cross-scored on one side. The tablets can be divided into 2 or 4 equal parts.

3. Target species

Dogs (weighing less than 17.5 kg).



4. Indications for use

For the treatment of mixed infections with the following roundworms and tapeworms in adult dogs and puppies:

Nematodes

Ascarids: *Toxocara canis*, *Toxascaris leonina* (late immature forms and mature forms)

Hookworms: *Uncinaria stenocephala*, *Ancylostoma caninum* (adults)

Cestodes

Tapeworms: *Taenia* spp., *Dipylidium caninum*,

5. Contraindications

Do not use simultaneously with piperazine compounds.

Do not exceed the stated dosage when treating pregnant bitches.

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

Do not use in dogs younger than 2 weeks of age and/or weighing less than 2 kg.

6. Special warnings

Special warnings:

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*.

Tapeworm infestation is certain to re-occur unless control of intermediate hosts such as fleas, mice etc. is undertaken.

Parasite resistance to any particular class of anthelmintic may develop following frequent, repeated use of an anthelmintic of that class.

Special precautions for safe use in the target species:

Any part-used tablets should be discarded.

Not for use in dogs younger than 2 weeks and/or weighing less than 2 kg.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In the interests of good hygiene, persons administering the tablet directly to a dog or by adding it to the dog's food, should wash their hands afterwards.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or label to the physician.

Pregnancy:

Do not use in bitches during the first two-thirds of pregnancy.

Consult a veterinary surgeon before treating pregnant animals for roundworms.

Lactation:

Can be used during lactation (see “Contraindications” and “Advice on correct administration”).

Interaction with other medicinal products and other forms of interaction:

Do not use simultaneously with piperazine as the anthelmintic effects of pyrantel and piperazine (used in many worming veterinary medicinal products for dogs) may be antagonized.

Concurrent use with other cholinergic compounds can lead to toxicity.

Overdose:

Benzimidazoles possess wide safety margin. Pyrantel is not absorbed systematically to any extent. Praziquantel also has a wide safety margin, of up to five times the recommended dose.

Major incompatibilities:

Not applicable.

7. Adverse events

Dogs:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Loose stool, diarrhoea, Vomiting.*
---	------------------------------------

*Transient.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: {national system details}

8. Dosage for each species, routes and method of administration

For oral use.

The recommended dose rates are: 15 mg/kg bodyweight febantel, 14.4 mg/kg pyrantel and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 10 kg bodyweight.

Tablets may be halved/quartered to allow accuracy of dosing.

The tablet(s) can be given directly to the dog or disguised in food. No restriction of access to food is required either before or after administration of the veterinary medicinal product.

9. Advice on correct administration

To ensure administration of a correct dosage, body weight should be determined as accurately as possible.

Puppies may be wormed with this veterinary medicinal product from 2 weeks of age and every 2 weeks until 12 weeks of age. Thereafter they should be treated at 3 monthly intervals until 6 months of age.

For the control of *Toxocara*, nursing bitches should be dosed 2 weeks after giving birth and every 2 weeks until weaning.

For adult dogs, a single dose should be used. The advice of a veterinarian should be sought regarding the need for and frequency of repeat treatment.

In the event of a heavy roundworm infestation, a repeat dose should be given after 14 days.

10. Withdrawal periods

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.

This medicinal product does not require any special storage conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton after {Exp.}. The expiry date refers to the last day of that month.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription. (PT, ES)

Veterinary medicinal product not subject to prescription. (BE, BG, IE, IT, NL,UK(NI))

Veterinary medicinal product subject to prescription except for some pack sizes. FI (packaging for 24, 30,50, 60, 100, 102 tbl)

14. Marketing authorisation numbers and pack sizes

OPA/Al/PVC-Al blister: 2 tablets (1 blister with 2 tablets), in a box.

OPA/Al/PVC-Al blister: 4 tablets (2 blisters with 2 tablets), in a box.

OPA/Al/PVC-Al blister: 10 tablets (1 blister with 10 tablets), in a box.

OPA/Al/PVC-Al blister: 30 tablets (3 blisters with 10 tablets), in a box.

OPA/Al/PVC-Al blister: 50 tablets (5 blisters with 10 tablets), in a box.

OPA/Al/PVC-Al blister: 100 tablets (10 blisters with 10 tablets), in a box.

OPA/Al/PVC-Al blister: 300 tablets (30 blisters with 10 tablets), in a box.

Not all pack sizes may be marketed.

15. Date on which the package leaflet was last revised

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

16. Contact details

Marketing authorisation holder and contact details to report suspected adverse reactions:

KRKA, d.d., Novo mesto

Šmarješka cesta 6

8501 Novo mesto

Slovenia

Telephone number, if applicable

Manufacturer responsible for batch release:

KRKA d.d, Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

KRKA - FARMA d.o.o., V. Holjevca 20/E, 10450 Jastrebarsko, Croatia

Local representatives and contact details to report suspected adverse reactions:

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

17. Other information