

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Tsefalen 500 mg film-coated tablets for dogs (AT, BE, BG, CY, CZ, DE, DK, EL, ES, FI, FR, HU, IE, IT, LT, LU, LV, MT, NL, PL, PT, RO, SE, SI, SK, UK (NI))  
Falakef 500 mg film-coated tablets for dogs (EE, NO)

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains:

### Active substance:

Cefalexin 500 mg  
(equivalent to cefalexin monohydrate 525.9 mg)

### Excipient:

Qualitative composition of excipients and other constituents	Quantitative composition if that information is essential for proper administration of the veterinary medicinal product
Titanium dioxide (E171)	-
Iron oxide yellow (E172)	0.13 mg
Iron oxide red (E172)	0.02 mg
Povidone K-90	-
Sodium Starch Glycolate Type A	-
Magnesium stearate	-
Glycerol	-
Talc	-
Hypromellose	-

Orange coloured oblong film-coated tablets, with a break-line on one side. Engraved with GP4 on the other side.

## 3. CLINICAL INFORMATION

### 3.1 Target species

Dogs.

### 3.2 Indications for use for each target species

For the treatment of infections of the respiratory system, urogenital system and skin, localised infections in soft tissue and gastrointestinal infections caused by cefalexin-sensitive bacteria.

### 3.3 Contraindications

Do not use in cases of hypersensitivity to the active substance, to other cephalosporins, to other substances of the beta lactam group or to any of the excipients.  
Do not use in rabbits, gerbils, guinea pigs and hamsters.

### 3.4 Special warnings

None.

### 3.5 Special precautions for use

#### Special precautions for safe use in the target species:

Wherever possible, use of the veterinary medicinal product should be based on susceptibility testing of the bacteria isolated from the animal and take into account official and local antimicrobial policies.

Deviating from the instructions given in the SPC when using the veterinary medicinal product may increase the prevalence of bacteria resistant to cefalexin and may also decrease the effectiveness of other beta-lactam antimicrobial treatments, due to the potential for cross-resistance. Therefore, deviation from the instructions must only be undertaken according to a risk/benefit assessment by the responsible veterinarian.

Do not administer in cases of known resistance to cephalosporin and penicillin.

As with other antibiotics which are excreted mainly by the kidneys, systemic accumulation may occur when renal function is impaired. In case of known renal insufficiency the dose should be reduced, and antimicrobials known to be nephrotoxic should not be administered concurrently.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Penicillins and cephalosporins may cause sensitisation (allergy) following injection, inhalation, ingestion, or skin contact. Sensitivity to penicillins may lead to cross sensitivity to cephalosporins and *vice versa*. Allergic reactions to these substances may occasionally be serious. Do not handle this veterinary medicinal product if you know you are sensitised or if you have been advised not to be in contact with such substances.

Handle this veterinary medicinal product with great care to avoid exposure, taking all recommended precautions. If you develop symptoms following exposure such as skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty breathing are more serious symptoms and require urgent medical attention.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Wash hands after use.

#### Special precautions for the protection of the environment:

Not applicable.

### 3.6 Adverse events

Dogs:

Rare (1 to 10 animals / 10,000 animals treated):	Hypersensitivity reaction <sup>1</sup> .
Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Nausea, vomiting, diarrhoea.

<sup>1</sup> In cases of hypersensitivity reactions the treatment should be discontinued.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

### 3.7 Use during pregnancy, lactation or lay

The safety of the veterinary medicinal product has not been established during pregnancy and lactation.

#### Pregnancy and lactation:

Use only according to the benefit-risk assessment by the responsible veterinarian. Laboratory studies in rats and mice have not produced any evidence of teratogenic, foetotoxic, or maternotoxic effects.

### 3.8 Interaction with other medicinal products and other forms of interaction

In order to ensure efficacy, the veterinary medicinal product should not be used in combination with bacteriostatic antibiotics.

Concurrent use of first generation cephalosporins with polypeptide antibiotics, aminoglycosides or some diuretics such as furosemide can enhance nephrotoxicity risks.

### 3.9 Administration routes and dosage

Oral use.

The recommended dose is 15 mg of cefalexin per kg of body weight twice a day (i.e. equivalent to 1 tablet twice a day for a dog weighing 33 kg). In severe or acute conditions, the dose may be doubled to 30 mg/kg twice daily.

The following is a guide for the use of the veterinary medicinal product:

<b>Bodyweight min kg</b>	<b>Bodyweight max kg</b>	<b>Number of tablets per dose*</b>
10.0	16.5	0.5
16.6	33.0	1
33.1	40.0	1.5

*\*Dose to be given twice per day*

The veterinary medicinal product must be administered for a minimum of 5 days.

- 14 days in cases of urinary tract infection,
- At least 15 days in cases of superficial infectious dermatitis,
- At least 28 days in cases of deep infectious dermatitis.

Any increase in dose or duration of treatment should be accordingly to a benefit/risk assessment by the responsible veterinarian (e.g. chronic pyoderma).

To ensure a correct dosage, body weight should be determined as accurately as possible.

The veterinary medicinal product can be given as whole tablets or crushed and added to food if necessary.

### 3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

Concerning acute toxicity, an LD50 > 0.5 g/kg has been recorded following oral administration in dogs. The administration of cefalexin has been shown to produce no serious side effects at several times the recommended dose rate.

### 3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

### 3.12 Withdrawal periods

Not applicable.

## 4. PHARMACOLOGICAL INFORMATION

### 4.1 ATCvet code: QJ01DB01

### 4.2 Pharmacodynamics

Cefalexin is a broad spectrum cephalosporin antibiotic with bactericidal activity against a wide range of Gram-positive and Gram-negative bacteria.

Cefalexin is a semi-synthetic bactericidal broad spectrum antibiotic belonging to the cephalosporin group which acts by interfering with bacterial cell wall formation. This bactericidal activity is mediated by drug binding to bacterial enzymes known as penicillin binding proteins (PBPs). Such enzymes are located on the inner membrane of the cell wall and their transpeptidase activity is required for the terminal stages of assembling this essential structure of the bacterial cell. Inactivation of PBPs interferes with the cross-linkage of peptidoglycan chains necessary for bacterial cell wall strength and rigidity. The bactericidal effect of cefalexin is mainly "time dependent".

Cefalexin is resistant to the action of staphylococcal penicillinase and is therefore active against the strains of *Staphylococcus aureus* that are not sensitive to penicillin (or related antibiotics such as ampicillin or amoxicillin) because of production of penicillinase.

Cefalexin is also active against the majority of ampicillin-resistant *E.coli*.

The following micro-organisms have been shown to be susceptible to Cefalexin *in vitro*: *Corynebacterium* spp, *Staphylococcus* spp (including penicillin-resistant strains), *Streptococcus* spp, *Escherichia coli*, *Moraxella* spp, *Pasteurella multocida*.

MIC data collected for cefalexin in canine isolates from the European Union (EU) (Stegmann *et al.* 2006)

Bacterial species/group and origin	No. isolates	MIC50	MIC90
<i>Staphylococcus pseudintermedius</i> (EU)	270	1	2
<i>Staphylococcus aureus</i> (EU)	36	2	8
Coagulase-negative staphylococci (EU)	21	1	8
Coagulase-positive staphylococci (EU)	24	1	2
$\beta$ -haemolytic streptococci (EU)	86	<0.5	2
<i>Enterococcus spp.</i> (EU)	331	>64	>64

<i>Pasteurella multocida</i> (EU)	193	4	4
<i>Escherichia coli</i> (EU)	260	8	16
<i>Proteus spp.</i> (EU)	71	16	16
<i>Klebsiella spp.</i> (EU)	11	4	4
<i>Enterobacter spp.</i> (EU)	39	8	>64

The three basic mechanisms of resistance to cephalosporins result from reduced permeability, enzymatic inactivation, or absence of specific penicillin-binding proteins.

### 4.3 Pharmacokinetics

Cefalexin is rapidly and almost completely absorbed in the gastrointestinal tract following oral administration. Cefalexin binds to a limited extent (10-20%) to plasma proteins. After oral administration of 15 mg/kg in tablets, peak blood concentration ( $C_{max}=15 \mu\text{g/ml}$ ) is usually reached between 1 and 2 hours ( $T_{max}=90 \text{ min}$ ).

Bioavailability is nearly 100% of the administered dose ( $AUC 6279 \mu\text{g min / ml}$ ). Cefalexin does not undergo biotransformation processes that are of pharmacokinetic significance.

The elimination half-life of cefalexin is about 1.5 hours ( $t_{1/2} = 90 \text{ min}$ ).

Elimination of the microbiologically active form is almost entirely via the kidneys by tubular excretion and glomerular filtration.

## 5. PHARMACEUTICAL PARTICULARS

### 5.1 Major incompatibilities

None known.

### 5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.

Shell life afterfirst opening the immediate packaging: 48 hours.

### 5.3 Special precautions for storage

This veterinary medicinal product does not require any special temperature conditions.

Return any halved tablet to the blister pack.

### 5.4 Nature and composition of immediate packaging

Carton box containing 1 PVC/Aluminium blister pack of 12 tablets.

Carton box containing 3 PVC/Aluminium blister pack of 12 tablets, with a total of 36 tablets.

Carton box containing 9 PVC/Aluminium blister pack of 12 tablets, with a total of 108 tablets.

Not all pack sizes may be marketed.

### 5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

**6. NAME OF THE MARKETING AUTHORISATION HOLDER**

NEXTMUNE Italy S.R.L.

**7. MARKETING AUTHORISATION NUMBER(S)**

**8. DATE OF FIRST AUTHORISATION**

Date of first authorisation:

**9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS**

**10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).



**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

{CARTON BOX}

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Tsefalen 500 mg film-coated tablets (AT, BE, BG, CY, CZ, DE, DK, EL, ES, FI, FR, HU, IE, IT, LT, LU, LV, MT, NL, PL, PT, RO, SE, SI, SK, UK (NI))  
Falakef 500 mg film-coated tablets (EE, NO)

**2. STATEMENT OF ACTIVE SUBSTANCES**

Cefalexin 500 mg

**3. PACKAGE SIZE**

12 tablets  
36 tablets  
108 tablets

**4. TARGET SPECIES**

Dogs.

**5. INDICATIONS**

**6. ROUTES OF ADMINISTRATION**

Oral use.

**7. WITHDRAWAL PERIODS**

**8. EXPIRY DATE**

EXP. {mm/yy }

Once opened, use within 48 hours.

**9. SPECIAL STORAGE PRECAUTIONS**

Return any halved tablet to the blister pack and use within 48 hours.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

User warnings  
Penicillins and cephalosporins may occasionally cause severe allergic reactions. See package leaflet for user warnings.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME OF THE MARKETING AUTHORISATION HOLDER**

NEXTMUNE Italy S.R.L.

**14. MARKETING AUTHORISATION NUMBERS**

**15. BATCH NUMBER**

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

{ BLISTER }

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Tsefalen (AT, BE, BG, CY, CZ, DE, DK, EL, ES, FI, FR, HU, IE, IT, LT, LU, LV, MT, NL, PL, PT, RO, SE, SI, SK, UK (NI))  
Falakef (EE, NO)

**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

Cefalexin 500 mg.

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/yyyy}  
Once opened, use within 48 hours.

**B. PACKAGE LEAFLET**

## PACKAGE LEAFLET

### 1. Name of the veterinary medicinal product

Tsefalen 500 mg film-coated tablets for dogs (AT, BE, BG, CY, CZ, DE, DK, EL, ES, FI, FR, HU, IE, IT, LT, LU, LV, MT, NL, PL, PT, RO, SE, SI, SK, UK (NI))

Falakef 500 mg film-coated tablets for dogs (EE, NO)

### 2. Composition

Each film-coated tablet contains:

#### Active substance:

Cefalexin 500 mg

(equivalent to 525.9 mg cefalexin monohydrate)

#### Excipients:

Iron oxide yellow (E172) 0.13 mg

Iron oxide red (E172) 0.02 mg

Orange coloured oblong film-coated tablets, with a break-line on one side. Engraved with GP4 on the other side.

### 3. Target species

Dogs.

### 4. Indications for use

For the treatment of infections of the respiratory system, urogenital system and skin, localised infections in soft tissue and gastrointestinal infections caused by cefalexin-sensitive bacteria.

### 5. Contraindications

Do not use in cases of hypersensitivity to the active substance, to other cephalosporins, to other substances of the beta lactam group or to any of the excipients.

Do not use in rabbits, gerbils, guinea pigs and hamsters.

### 6. Special warnings

#### Special warnings:

None.

#### Special precautions for safe use in the target species:

Wherever possible, use of the veterinary medicinal product should be based on susceptibility testing of the bacteria isolated from the animal and take into account official and local antimicrobial policies.

Deviating from the instructions given in the SPC when using the veterinary medicinal product may increase the prevalence of bacteria resistant to cefalexin and may also decrease the effectiveness of other beta-lactam antimicrobial treatments, due to the potential for cross-resistance. Therefore, deviation from the instructions must only be undertaken according to a risk/benefit assessment by the responsible veterinarian.

Do not administer in cases of known resistance to cephalosporin and penicillin.

As with other antibiotics which are excreted mainly by the kidneys, systemic accumulation may occur when renal function is impaired. In case of known renal insufficiency the dose should be reduced, and antimicrobials known to be nephrotoxic should not be administered concurrently.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Penicillins and cephalosporins may cause sensitisation (allergy) following injection, inhalation, ingestion, or skin contact. Sensitivity to penicillins may lead to cross sensitivity to cephalosporins and *vice versa*. Allergic reactions to these substances may occasionally be serious. Do not handle this veterinary medicinal product if you know you are sensitised or if you have been advised not to be in contact with such substances.

Handle this veterinary medicinal product with great care to avoid exposure, taking all recommended precautions. If you develop symptoms following exposure such as skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty breathing are more serious symptoms and require urgent medical attention.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Wash hands after use.

Special precautions for the protection of the environment:

Not applicable.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy and lactation. Use only accordingly to the benefit/risk assessment by the responsible veterinarian. Laboratory studies in rats and mice have not produced any evidence of teratogenic, foetotoxic, maternotoxic effects.

Interaction with other medicinal products and other forms of interaction:

In order to ensure efficacy, the veterinary medicinal product should not be used in combination with bacteriostatic antibiotics.

Concurrent use of first generation cephalosporins with polypeptide antibiotics, aminoglycosides or some diuretics such as furosemide can enhance nephrotoxicity risks.

Overdose:

Concerning acute toxicity, an LD50 > 0.5 g/kg has been recorded following oral administration in dogs. The administration of cefalexin has been shown to produce no serious side effects at several times the recommended dose rate.

Special restrictions for use and special conditions for use:

Not applicable.

Major incompatibilities:

None.

**7. Adverse events**

Dogs:

Rare (1 to 10 animals / 10,000 animals treated): Hypersensitivity reaction <sup>1</sup> .
Very rare (<1 animal / 10,000 animals treated, including isolated reports): Nausea, vomiting, diarrhoea.

<sup>1</sup> In cases of hypersensitivity reactions the treatment should be discontinued.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system:

## **8. Dosage for each species, routes and method of administration**

Oral use.

The recommended dose is 15 mg of cefalexin per kg of body weight twice a day (i.e. equivalent to 1 tablet twice a day for a dog weighing 33 kg). In severe or acute conditions, the dose may be doubled to 30 mg/kg twice daily.

The following is a guide for the use of the veterinary medicinal product:

<b>Bodyweight min kg</b>	<b>Bodyweight max kg</b>	<b>Number of tablets per dose*</b>
10.0	16.5	0.5
16.6	33.0	1
33.1	40.0	1.5

*\*Dose to be given twice per day*

The veterinary medicinal product must be administered for a minimum of 5 days.

- 14 days in cases of urinary tract infection,
- At least 15 days in cases of superficial infectious dermatitis,
- At least 28 days in cases of deep infectious dermatitis.

Any increase in dose or duration of treatment should be accordingly to a benefit/risk assessment by the responsible veterinarian (e.g. chronic pyoderma).

## **9. Advice on correct administration**

To ensure a correct dosage, body weight should be determined as accurately as possible.

The veterinary medicinal product can be given as whole tablets or crushed and added to food if necessary.

## **10. Withdrawal periods**

Not applicable.

## **11. Special storage precautions**

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special temperature conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton after Exp. The expiry date refers to the last day of that month. Shelf life after first opening the immediate packaging: 48 hours.

Return any halved tablet to the blister pack.

## **12. Special precautions for disposal**

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

## **13. Classification of veterinary medicinal products**

Veterinary medicinal product subject to prescription.

## **14. Marketing authorisation numbers and pack sizes**

Carton box containing 1 PVC/Aluminium blister pack of 12 tablets.

Carton box containing 3 PVC/Aluminium blister pack of 12 tablets, with a total of 36 tablets.

Carton box containing 9 PVC/Aluminium blister pack of 12 tablets, with a total of 108 tablets.

Not all pack sizes may be marketed.

## **15. Date on which the package leaflet was last revised**

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

## **16. Contact details**

Marketing authorisation holder and contact details to report suspected adverse reactions:

NEXTMUNE ITALY S.R.L.  
Via G.B. Benzoni, 50  
26020 Palazzo Pignano (CR)  
Italy  
Phone +39.0373.982024  
E-mail: info.it@nextmune.com

Manufacturer responsible for batch release:

ACS DOBFAR S.p.A.  
via Laurentina Km 24,730  
00071 Pomezia (RM)  
Italy

## **17. Other information**