

*[Version 9.1,11/2024]*

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Eliminall 50 mg spot-on solution for cats (DE, GR, PT)  
Exproline vet 50 mg spot-on solution for cats (FI)

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 0.5 ml pipette contains:

### Active substances:

Fipronil 50 mg

### Excipients:

Qualitative composition of excipients and other constituents	Quantitative composition if that information is essential for proper administration of the veterinary medicinal product
Butylhydroxyanisol (E320)	0.10 mg
Butylhydroxytoluene (E321)	0.05 mg
Polysorbate 80	
Povidone K25	
Dimethyl sulfoxide	

Light yellow to yellow, clear liquid.

## 3. CLINICAL INFORMATION

### 3.1 Target species

Cats

### 3.2 Indications for use for each target species

Treatment and prevention of flea (*Ctenocephalides* spp.) and tick (*Dermacentor reticulatus*, *Ixodes ricinus*) infestations in cats.

The veterinary medicinal product has a persistent insecticidal efficacy for up to 4 weeks against fleas (*Ctenocephalides* spp.) and acaricidal efficacy for up to 4 weeks against *Ixodes ricinus* and for up to 1 week against *Dermacentor reticulatus* and *Rhipicephalus sanguineus*. If *Rhipicephalus sanguineus* ticks are present when the veterinary medicinal product is applied, all the ticks may not be killed within the first 48 hours but they may be killed within a week.

The veterinary medicinal product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD), where this has been previously diagnosed by a veterinary surgeon.

### 3.3 Contraindications

In the absence of available data, the veterinary medicinal product should not be used on kittens less than 2 months old and/or weighing less than 1 kg.

Do not use on sick (e.g. systemic diseases, fever...) or convalescent animals. Do not use in rabbits, as adverse reactions and even death could occur.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

### **3.4 Special warnings**

Fleas from pets often infest the animal's basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated, in case of massive infestation and at the beginning of the control measures, with a suitable insecticide and vacuumed regularly.

The veterinary medicinal product does not prevent ticks from attaching to the animals. If the animal has been treated prior to exposure to the ticks, the tick will be killed in the first 24-48 hours after attachment. This will usually be prior to engorgement, minimising but not excluding the risk of transmission of diseases. Once dead, ticks will often drop off the animal, but any remaining ticks may be removed with a gentle pull.

No data on the effect of bathing/shampooing on the efficacy of the veterinary medicinal product in cats are available. However, based on information available for dogs, weekly immersion in water for one minute reduces the period of persistent insecticidal efficacy against fleas by one week.

For optimum control of flea problems in a multi-pet household, all dogs and cats in the household should be treated with a suitable insecticide.

When used as part of a strategy for the treatment of Flea Allergy Dermatitis, monthly applications to the allergic patient and to other cats in the household are recommended.

### **3.5 Special precautions for use**

#### Special precautions for safe use in the target species:

Avoid contact with the animal's eyes. In the case of accidental eye contact, immediately and thoroughly flush the eyes with water.

Do not apply the veterinary medicinal product on wounds or damaged skin.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product can cause mucous membrane and eye irritation. Therefore, contact between the veterinary medicinal product and the mouth or eyes should be avoided.

In case of accidental eye contact immediately and thoroughly flush the eyes with water. If eye irritation persists seek medical advice and show the package leaflet or the label to the physician. Avoid contents coming into contact with the fingers. If this occurs, wash off immediately with soap and water.

Wash hands after use.

Do not smoke, drink or eat during application.

People with a known hypersensitivity to fipronil or dimethyl sulfoxide or other excipients should avoid contact with the veterinary medicinal product.

Treated animals should not be handled until the application site is dry, and children should not be allowed to play with treated animals until the application site is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals should not be allowed to sleep with owners, especially children.

Keep pipettes in the original packaging and dispose of used pipette immediately.

#### Special precautions for the protection of the environment:

Not applicable.

### 3.6 Adverse events

Cats:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Application site skin squamosis <sup>1</sup> , Application site alopecia <sup>1</sup> , Application site pruritus <sup>1</sup> , Application site erythema <sup>1</sup> Generalised itching, Alopecia general Hypersalivation, Vomiting Neurological signs <sup>3</sup> , Hyperaesthesia <sup>3</sup> Depression <sup>3</sup>
Undetermined (cannot be estimated from the available data)	Hypersalivation <sup>2</sup>

<sup>1</sup>Transient.

<sup>2</sup>May be observed for a brief period in the case of licking the administration site.

<sup>3</sup>Reversible.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

### 3.7 Use during pregnancy, lactation or lay

Pregnancy and lactation:

Laboratory studies have not produced any evidence of teratogenic or embryotoxic effects. Studies have not been carried out with this veterinary medicinal product in pregnant and lactating queens. Use only according to the benefit-risk assessment by the responsible veterinarian.

### 3.8 Interaction with other medicinal products and other forms of interaction

None known.

### 3.9 Administration routes and dosage

Route of administration and dosage:


Spot-on use. External use only.

Administer by topical application to the skin 1 pipette of 0.5 ml per animal.

To ensure a correct dosage, body weight should be determined as accurately as possible.

Method of administration:

Remove the pipette from the packaging. Hold the pipette in an upright position, twist and pull the cap off. Turn the cap around and place the other end of the cap back on the pipette. Push and twist the cap to break the seal, then remove the cap from the pipette. Spread the hairs of the animal between the shoulder blades to make the skin visible. Place the tip of the pipette on the skin and squeeze gently to empty its contents onto the skin, preferably at two spots, one at the base of the skull and a second 2-3 cms further back.



It is important to make sure that the veterinary medicinal product is applied to an area where the animal cannot lick it off, and to make sure that animals do not lick each other following treatment. Temporary changes to the coat (clumped/greasy hair and/or deposits on the hair) may be noted at the application site which normally disappears within 24 hours.

Treatment schedule:

For optimal control of flea and/or tick infestation the treatment schedule can be based on the local epidemiological situation.

In the absence of safety studies, the minimum treatment interval is 4 weeks.

**3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)**

No adverse effects were observed in target animal safety studies in cats and kittens aged 2 months and older and weighing about 1 kg treated once a month at five times the recommended dose for three consecutive months. The risk of experiencing adverse effects may however increase with overdosing (see section 3.6).

**3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance**

Not applicable.

**3.12 Withdrawal periods**

Not applicable.

**4. PHARMACOLOGICAL INFORMATION**

**4.1 ATCvet code : QP53AX15**

**4.2 Pharmacodynamics**

Fipronil is an insecticide and acaricide belonging to the phenylpyrazole family. It acts by inhibiting the GABA complex, binding to the chloride channel and thereby blocking pre- and post-synaptic transfer of chloride ions across cell membranes. This results in uncontrolled activity of the central nervous system and death of insects or acarids.

Fipronil exhibits an insecticidal and acaricidal activity against fleas (*Ctenocephalides* spp.) and ticks (*Rhipicephalus* spp., *Dermacentor* spp., *Ixodes* spp. including *Ixodes ricinus*) in the cat.

Fleas will be killed within 24 hours. Ticks will usually be killed within 48 h after contact with fipronil, however if *Rhipicephalus sanguineus* ticks are present when the veterinary medicinal product is applied, all of the ticks may not be killed within the first 48 hours, but may be killed within a week.

**4.3 Pharmacokinetics**

*In vitro*, fipronil is mainly metabolised with subcellular liver fractions to its sulfone derivative. However, this may be of limited relevance *in vivo* as fipronil is poorly absorbed in the cat. The concentrations of fipronil on the hair decrease with time.

## **5. PHARMACEUTICAL PARTICULARS**

### **5.1 Major incompatibilities**

None known.

### **5.2 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.

### **5.3 Special precautions for storage**

Store in the original container in order to protect from light and moisture.

The veterinary medicinal product should be maintained at room temperature (above 14°C) for approximately one hour prior to administration.

### **5.4 Nature and composition of immediate packaging**

White polypropylene pipette closed with either a polyethylene or polyoxymethylene cap. Each 0.5 ml pipette is packed in a polyethylene terephthalate/aluminium/low density polyethylene triplex bag.

#### Pack sizes:

Box containing 1, 3, 6, 10, 20 or 30 pipettes.

Not all pack sizes may be marketed.

### **5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products**

The veterinary medicinal product should not enter water courses as Fipronil may be dangerous for fish and other aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty container.

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

## **6. NAME OF THE MARKETING AUTHORISATION HOLDER**

KRKA, d.d., Novo mesto

## **7. MARKETING AUTHORISATION NUMBER(S)**

## **8. DATE OF FIRST AUTHORISATION**

Date of first authorisation:

**9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS**

<{MM/YYYY}>

<{DD/MM/YYYY}>

<{DD month YYYY}>

**10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription (GR).

Veterinary medicinal product not subject to prescription (FI, DE, PT).

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**BOX**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Eliminall 50 mg spot-on solution

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each 0.5 ml pipette contains:

fipronil                      50 mg

**3. PACKAGE SIZE**

- 1 x 0.5 ml
- 3 x 0.5 ml
- 6 x 0.5 ml
- 10 x 0.5 ml
- 20 x 0.5 ml
- 30 x 0.5 ml

**4. TARGET SPECIES**

Cats




**5. INDICATIONS**

Treatment of fleas and ticks.

*For products not subject to veterinary prescription.*

**6. ROUTES OF ADMINISTRATION**

Spot-on use.



**7. WITHDRAWAL PERIODS**

**8. EXPIRY DATE**

Exp.

**9. SPECIAL STORAGE PRECAUTIONS**

Store in the original container in order to protect from light and moisture.  
The veterinary medicinal product should be maintained at room temperature (above 14°C) for approximately one hour prior to administration.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME OF THE MARKETING AUTHORISATION HOLDER**

KRKA, d.d., Novo mesto

**14. MARKETING AUTHORISATION NUMBERS**

**15. BATCH NUMBER**

Lot

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**BAG**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Eliminall



**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

fipronil                      50 mg


**3. BATCH NUMBER**

Lot

**4. EXPIRY DATE**

Exp.

**KRKA**



**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**PIPETTE**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Eliminall



**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

50 mg

**3. BATCH NUMBER**

Lot

**4. EXPIRY DATE**

Exp.

## **B. PACKAGE LEAFLET**

## PACKAGE LEAFLET

### 1. Name of the veterinary medicinal product

Eliminall 50 mg spot-on solution for cats

### 2. Composition

Each 0.5 ml pipette contains:

#### Active substance:

Fipronil 50 mg

#### Excipients:

Butylhydroxyanisol (E320) 0.10 mg

Butylhydroxytoluene (E321) 0.05 mg

Light yellow to yellow, clear liquid.

### 3. Target species

Cats



### 4. Indications for use

Treatment and prevention of flea (*Ctenocephalides* spp.) and tick (*Dermacentor reticulatus*, *Ixodes ricinus*) infestations in cats.

The veterinary medicinal product has a persistent insecticidal efficacy for up to 4 weeks against fleas (*Ctenocephalides* spp.) and acaricidal efficacy for up to 4 weeks against *Ixodes ricinus* and for up to 1 week against *Dermacentor reticulatus* and *Rhipicephalus sanguineus*. If *Rhipicephalus sanguineus* ticks are present when the veterinary medicinal product is applied, all the ticks may not be killed within the first 48 hours but they may be killed within a week.

The veterinary medicinal product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD), where this has been previously diagnosed by a veterinary surgeon.

### 5. Contraindications

In the absence of available data, the veterinary medicinal product should not be used on kittens less than 2 months old and/or weighing less than 1 kg.

Do not use on sick (e.g. systemic diseases, fever...) or convalescent animals. Do not use in rabbits, as adverse reactions and even death could occur.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

### 6. Special warnings

#### Special warnings:

Fleas from pets often infest the animal's basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated, in case of massive infestation and at the beginning of the control measures, with a suitable insecticide and vacuumed regularly.

The veterinary medicinal product does not prevent ticks from attaching to the animals. If the animal has been treated prior to exposure to the ticks, the tick will be killed in the first 24-48 hours after attachment. This will usually be prior to engorgement, minimising but not excluding the risk of transmission of diseases. Once dead, ticks will often drop off the animal, but any remaining ticks may be removed with a gentle pull.

No data on the effect of bathing/shampooing on the efficacy of the veterinary medicinal product in cats are available. However, based on information available for dogs, weekly immersion in water for one minute reduces the period of persistent insecticidal efficacy against fleas by one week.

For optimum control of flea problems in a multi-pet household, all dogs and cats in the household should be treated with a suitable insecticide.

When used as part of a strategy for the treatment of Flea Allergy Dermatitis, monthly applications to the allergic patient and to other cats in the household are recommended.

#### Special precautions for safe use in the target species:

Avoid contact with the animal's eyes. In the case of accidental eye contact, immediately and thoroughly flush the eyes with water.

Do not apply the veterinary medicinal product on wounds or damaged skin.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product can cause mucous membrane and eye irritation. Therefore, contact between the veterinary medicinal product and the mouth or eyes should be avoided.

In case of accidental eye contact immediately and thoroughly flush the eyes with water. If eye irritation persists seek medical advice and show the package leaflet or the label to the physician.

Avoid contents coming into contact with the fingers. If this occurs, wash off immediately with soap and water.

Wash hands after use.

Do not smoke, drink or eat during application.

People with a known hypersensitivity to fipronil or dimethyl sulfoxide or other excipients should avoid contact with the veterinary medicinal product.

Treated animals should not be handled until the application site is dry, and children should not be allowed to play with treated animals until the application site is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals should not be allowed to sleep with owners, especially children.

Keep pipettes in the original packaging and dispose of used pipette immediately.

#### Pregnancy and lactation:

Laboratory studies using fipronil have not shown any evidence of teratogenic or embryotoxic effects. Studies have not been carried out with this veterinary medicinal product in pregnant and lactating queens. Use only according to the benefit-risk assessment by the responsible veterinarian.

#### Interaction with other medicinal products and other forms of interaction:

None known.

#### Overdose:

No adverse effects were observed in target animal safety studies in cats and kittens aged 2 months and older and weighing about 1 kg treated once a month at five times the recommended dose for three consecutive months. The risk of experiencing adverse effects may however increase with overdosing (see section Adverse events).

#### Major incompatibilities:

None known.

## 7. Adverse events

Cats:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Application site skin squamosis (scaling) <sup>1</sup> , Application site alopecia (hair loss) <sup>1</sup> , Application site pruritus (itching) <sup>1</sup> , Application site erythema (reddening) <sup>1</sup>  Generalised itching, Alopecia general (General hair loss)  Hypersalivation, Vomiting  Neurological signs <sup>3</sup> , Hyperaesthesia <sup>3</sup>  Depression <sup>3</sup>
Undetermined (cannot be estimated from the available data)	Hypersalivation <sup>2</sup>

<sup>1</sup>Transient.

<sup>2</sup>May be observed for a brief period in the case of licking the administration site.

<sup>3</sup>Reversible.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: {national system details}.

## 8. Dosage for each species, routes and method of administration

### Route of administration and dosage:


Spot-on use. External use only.

Administer by topical application to the skin 1 pipette of 0.5 ml per animal.

To ensure a correct dosage, body weight should be determined as accurately as possible.

### Method of administration:

Remove the pipette from the packaging. Hold the pipette in an upright position, twist and pull the cap off. Turn the cap around and place the other end of the cap back on the pipette. Push and twist the cap to break the seal, then remove the cap from the pipette. Spread the hairs of the animal between the shoulder blades to make the skin visible. Place the tip of the pipette on the skin and squeeze gently to empty its contents onto the skin, preferably at two spots, one at the base of the skull and a second 2-3 cms further back.



## **9. Advice on correct administration**

It is important to make sure that the veterinary medicinal product is applied to an area where the animal cannot lick it off, and to make sure that animals do not lick each other following treatment. Temporary changes to the coat (clumped/greasy hair and/or deposits on the hair) may be noted at the application site.

For optimal control of flea and/or tick infestation the treatment schedule can be based on the local epidemiological situation.

In the absence of safety studies, the minimum treatment interval is 4 weeks.

## **10. Withdrawal periods**

Not applicable.

## **11. Special storage precautions**

Keep out of the sight and reach of children.

Store in the original container in order to protect from light and moisture.

The veterinary medicinal product should be maintained at room temperature (above 14°C) for approximately one hour prior to administration.

Do not use this veterinary medicinal product after the expiry date which is stated on the labels and carton after Exp. The expiry date refers to the last day of that month.

## **12. Special precautions for disposal**

This veterinary medicinal product should not enter water courses as Fipronil may be dangerous for fish and other aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty container.

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

## **13. Classification of veterinary medicinal products**

Veterinary medicinal product subject to prescription (GR).

Veterinary medicinal product not subject to prescription (FI, DE, PT).

## **14. Marketing authorisation numbers and pack sizes**

White polypropylene pipette closed with either a polyethylene or polyoxymethylene cap. Each 0.5 ml pipette is packed in a polyethylene terephthalate/aluminium/low density polyethylene triplex bag.

Pack sizes:

Box containing 1, 3, 6, 10, 20 or 30 pipettes.

Not all pack sizes may be marketed.

**15. Date on which the package leaflet was last revised**

<{MM/YYYY}>

<{DD/MM/YYYY}>

<{DD month YYYY}>

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

**16. Contact details**

Marketing authorisation holder and manufacturer responsible for batch release and contact details to report suspected adverse reactions:

KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

Local representatives and contact details to report suspected adverse reactions:

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.