

*[Version 9,10/2021]*

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Suigen APP 2,9,11 emulsion for injection for pigs

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One dose (1 ml) of the vaccine contains:

### Active substances:

Inactivated strains of:

<i>Actinobacillus pleuropneumoniae</i> , serovar 2, strain WSLB 3012, inactivated	RP $\geq$ 1*
<i>Actinobacillus pleuropneumoniae</i> , serovar 9, strain WSLB 3013 and serovar 11, strain WSLB 3057, inactivated **	RP $\geq$ 1*
<i>Actinobacillus pleuropneumoniae</i> , APX I toxoid	RP $\geq$ 1*
<i>Actinobacillus pleuropneumoniae</i> , APX II toxoid	RP $\geq$ 1*
<i>Actinobacillus pleuropneumoniae</i> , APX III toxoid	RP $\geq$ 1*

\* RP = Relative potency (determined by ELISA method) in comparison with the reference serum obtained after vaccination of mice with a vaccine batch that has successfully passed the challenge test on the target species.

\*\* The serovars 9 and 11 are determined together as one value because the potency test is not able to distinguish between these 2 antigen variants.

### Adjuvant:

Montanide ISA 35 VG

0.2 ml

### Excipients:

Qualitative composition of excipients and other constituents	Quantitative composition if that information is essential for proper administration of the veterinary medicinal product
Thiomersal	0.1 mg
Sodium chloride	
Water for injection	

The ready to use vaccine has a milky liquid of light grey to white colour and may contain a small amount of sediment which easily disperses after shaking.

## 3. CLINICAL INFORMATION

### 3.1 Target species

Pigs

### 3.2 Indications for use for each target species

For active immunisation of pigs from 6 weeks of age onwards to reduce lung lesions and to reduce colonisation of the respiratory tract caused by pleuropneumonia due to *Actinobacillus pleuropneumoniae* serovars expressing the APX toxins I, II and III.

Onset of immunity: 3 weeks after the second dose

Duration of immunity: 20 weeks after the second dose

### 3.3 Contraindications

None

### 3.4 Special warnings

Vaccinate healthy animals only

### 3.5 Special precautions for use

Special precautions for safe use in the target species:

Not applicable

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Special precautions for the protection of the environment:

Not applicable.

### 3.6 Adverse events

Pigs

Very common (>1 animal / 10 animals treated):	- Injection site induration
Common (1 to 10 animals / 100 animals treated):	- Injection site swelling* - Injection site reddening - Elevated temperature**

\* with a diameter of 10 cm which spontaneously subsides within 3 to 14 days

\*\* up to 0.8°C for 1 or 2 days after injection

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

### 3.7 Use during pregnancy, lactation or lay

The safety of the veterinary medicinal product has not been established during pregnancy and lactation.

### 3.8 Interaction with other medicinal products and other forms of interaction

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case-by-case basis.

### 3.9 Administration routes and dosage

Before use, allow the vaccine to reach room temperature of 15 to 25 °C and shake well.

Administer intramuscularly (preferably to the paraauricular area) one dose (1ml) of the veterinary medicinal product according to the following regimen of vaccination.

From an age of 6 weeks, administer 2 doses at an interval of 3 weeks

### **3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)**

No other symptoms were observed after an overdose administration (2 doses) of the veterinary medicinal product than those described in section 3.6, except for a temporary elevation of body temperature up to 1.5 °C in some of the animals.

### **3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance**

Not applicable.

### **3.12 Withdrawal periods**

Zero days.

## **4. IMMUNOLOGICAL INFORMATION**

### **4.1 ATCvet code:**

QI09AB07

Immunological properties

The vaccine contains inactivated whole-cell antigens of *Actinobacillus pleuropneumoniae* s.2, s.9 and s.11 and toxoids APX I, APX II and APX III. The vaccination with these antigens induce an active immunisation to protect against the consequences of field infection by *Actinobacillus pleuropneumoniae*.

## **5. PHARMACEUTICAL PARTICULARS**

### **5.1 Major incompatibilities**

Do not mix with any other veterinary medicinal product

### **5.2 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 2 years

Shelf life after first opening the immediate packaging: 10 hours

### **5.3 Special precautions for storage**

Store and transport refrigerated (2 °C – 8 °C).

Do not freeze.

Protect from light.

### **5.4 Nature and composition of immediate packaging**

High density polyethylene vial of 15, 60, 120 or 250 ml volume, glass vial (hydrolytic glass class I) of 10 ml or glass vial (hydrolytic glass class II) of 50 and 100 ml sealed with a chlorobutyl rubber stopper for perforation and an aluminium cap or flip-off cap, in a cardboard or plastic box with 10 wells.

Package Leaflet is a part of each packaging.

Package sizes:

Cardboard box:

1 x 10 ml (1 x 10 doses) in 10 ml glass vial (class I),  
1 x 10 ml (1 x 10 doses) in 15 ml HDPE plastic vial,  
1 x 50 ml (1 x 50 doses) in 50 ml glass vial (class II),  
1 x 50 ml (1 x 50 doses) in 60 ml HDPE plastic vial,  
1 x 100 ml (1 x 100 doses) in 100 ml glass vial (class II),  
1 x 100 ml (1 x 100 doses) in 120 ml HDPE plastic vial,  
1 x 250 ml (1 x 250 doses) in 250 ml HDPE plastic vial

Plastic box:

10 x 10 ml (10 x 10 doses) in 10 ml glass vial (class I)

Not all pack sizes may be marketed.

### **5.5 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Medicines should not be disposed of via wastewater or household waste.

## **6. NAME OF THE MARKETING AUTHORISATION HOLDER**

VIRBAC

## **7. MARKETING AUTHORISATION NUMBER(S)**

## **8. DATE OF FIRST AUTHORISATION**

DD/MM/YYYY

## **9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS**

## **10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

## **ANNEX II**

Not applicable

## **OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

None

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**



**PARTICULARS TO APPEAR ON THE OUTER PACKAGE****CARTON BOX****PLASTIC BOX WITH COVER****1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Suigen APP 2,9,11 emulsion for injection

**2. STATEMENT OF ACTIVE SUBSTANCES**

One dose (1 ml) of the vaccine contains:

**Active substances:**

Inactivated strains of:

*Actinobacillus pleuropneumoniae*, serovar 2, strain WSLB 3012, inactivated RP  $\geq 1^*$ *Actinobacillus pleuropneumoniae*, serovar 9, strain WSLB 3013 and serovar 11, strain WSLB 3057, inactivated \*\* RP  $\geq 1^*$ *Actinobacillus pleuropneumoniae*, APX I toxoid RP  $\geq 1^*$ *Actinobacillus pleuropneumoniae*, APX II toxoid RP  $\geq 1^*$ *Actinobacillus pleuropneumoniae*, APX III toxoid RP  $\geq 1^*$ 

\* RP = Relative potency (determined by ELISA method) in comparison with the reference serum obtained after vaccination of mice with a vaccine batch that has successfully passed the challenge test in the target species.

\*\* The serovars 9 and 11 are determined together as one value because the potency test is not able to distinguish between these 2 antigen variants.

**3. PACKAGE SIZE**

10 ml

50 ml,

100 ml,

250 ml

10×10 ml

**4. TARGET SPECIES**

Pigs

**5. INDICATIONS****6. ROUTES OF ADMINISTRATION**

Intramuscular route

**7. WITHDRAWAL PERIODS**

Withdrawal period: zero days

**8. EXPIRY DATE**

Exp. {mm/yyyy}  
Once opened use within 10 hours

<b>9. SPECIAL STORAGE PRECAUTIONS</b>
---------------------------------------

Store and transport refrigerated.  
Do not freeze.  
Protect from light.

<b>10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”</b>
--

Read the package leaflet before use.

<b>11. THE WORDS “FOR ANIMAL TREATMENT ONLY”</b>
--

For animal treatment only.

<b>12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”</b>
--

Keep out of the sight and reach of children.

<b>13. NAME OF THE MARKETING AUTHORISATION HOLDER</b>
---

MAH: VIRBAC

Manufacturer: BIOVETA

<b>14. MARKETING AUTHORISATION NUMBERS</b>
--

To be completed nationally

<b>15. BATCH NUMBER</b>
-------------------------

Lot {number}

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE****PAPER ETIQUETTE 100 ml (250 ml)****1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Suigen APP 2,9,11 emulsion for injection

**2. STATEMENT OF ACTIVE SUBSTANCES**

One dose (1 ml) of the vaccine contains:

**Active substances:**

Inactivated strains of:

<i>Actinobacillus pleuropneumoniae</i> , serovar 2, strain WSLB 3012, inactivated	RP $\geq$ 1*
<i>Actinobacillus pleuropneumoniae</i> , serovar 9, strain WSLB 3013 and serovar 11, strain WSLB 3057, inactivated **	RP $\geq$ 1*
<i>Actinobacillus pleuropneumoniae</i> , APX I toxoid	RP $\geq$ 1*
<i>Actinobacillus pleuropneumoniae</i> , APX II toxoid	RP $\geq$ 1*
<i>Actinobacillus pleuropneumoniae</i> , APX III toxoid	RP $\geq$ 1*

\* RP = Relative potency (determined by ELISA method) in comparison with the reference serum obtained after vaccination of mice with a vaccine batch that has successfully passed the challenge test in the target species.

\*\* The serovars 9 and 11 are determined together as one value.

**3. TARGET SPECIES**

Pigs

**4. ROUTES OF ADMINISTRATION**

Intramuscular route.

Read the package leaflet before use.

**5. WITHDRAWAL PERIODS**

Withdrawal period: zero days

**6. EXPIRY DATE**

Exp. {mm/yyyy}

Once opened use within 10 hours

**7. SPECIAL STORAGE PRECAUTIONS**

Store and transport refrigerated.

Do not freeze.

Protect from light.

**8. NAME OF THE MARKETING AUTHORISATION HOLDER**

VIRBAC

Manufacturer: BIOVETA

<b>9. BATCH NUMBER</b>
------------------------

Lot {number}

<b>MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS</b>
---

<b>PAPER ETIQUETTE 10 ml (50ml)</b>
-------------------------------------

<b>1. NAME OF THE VETERINARY MEDICINAL PRODUCT</b>
--

Suigen APP 2,9,11

<b>2. QUALITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES</b>
--

Inactivated strains of:

*Actinobacillus pleuropneumoniae*, serovar 2, strain WSLB 3012, inactivated

*Actinobacillus pleuropneumoniae*, serovar 9, strain WSLB 3013 and serovar 11, strain WSLB 3057, inactivated \*\*

*Actinobacillus pleuropneumoniae*, APX I toxoid

*Actinobacillus pleuropneumoniae*, APX II toxoid

*Actinobacillus pleuropneumoniae*, APX III toxoid

<b>3. BATCH NUMBER</b>
------------------------

Lot {number}

<b>4. EXPIRY DATE</b>
-----------------------

Exp. {mm/yyyy}

Once opened use within 10 hours

## **B. PACKAGE LEAFLET**

## PACKAGE LEAFLET

### 1. Name of the veterinary medicinal product

Suigen APP 2,9,11 emulsion for injection for pigs

### 2. Composition

One dose (1 ml) of the vaccine contains:

#### Active substances:

Inactivated strains of:

<i>Actinobacillus pleuropneumoniae</i> , serovar 2, strain WSLB 3012, inactivated	RP $\geq$ 1*
<i>Actinobacillus pleuropneumoniae</i> , serovar 9, strain WSLB 3013 and serovar 11, strain WSLB 3057, inactivated **	RP $\geq$ 1*
<i>Actinobacillus pleuropneumoniae</i> , APX I toxoid	RP $\geq$ 1*
<i>Actinobacillus pleuropneumoniae</i> , APX II toxoid	RP $\geq$ 1*
<i>Actinobacillus pleuropneumoniae</i> , APX III toxoid	RP $\geq$ 1*

\* RP = Relative potency (determined by ELISA method) in comparison with the reference serum obtained after vaccination of mice with a vaccine batch that has successfully passed the challenge test in the target species.

\*\* The serovars 9 and 11 are determined together as one value because the potency test is not able to distinguish between these 2 antigen variants.

#### Adjuvant:

Montanide ISA 35 VG	0.2 ml
---------------------	--------

#### Excipients:

Thiomersal	0.1 mg
Sodium chloride	
Water for injection	

The ready to use vaccine has a milky liquid of light grey to white colour and may contain a small amount of sediment which easily disperses after shaking.

### 3. Target species

Pigs

### 4. Indications for use

For active immunisation of pigs from 6 weeks of age onwards to reduce lung lesions and to reduce colonisation of the respiratory tract caused by pleuropneumonia due to *Actinobacillus pleuropneumoniae* serovars expressing the APX toxins I, II and III.

Onset of immunity: 3 weeks after the second dose

Duration of immunity: 20 weeks after the second dose

### 5. Contraindications

None

## 6. Special warnings

### Special warnings:

Vaccinate healthy animals only

### Special precautions for safe use in the target species:

Not applicable

### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

### Special precautions for the protection of the environment:

Not applicable

### Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy and lactation.

### Interactions with other medicinal products and other forms of interaction:

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case-by-case basis.

### Overdose:

No other symptoms were observed after an overdose administration (2 doses) of the veterinary medicinal product than those described in section "Adverse events", except for a temporary elevation of body temperature up to 1.5 °C in some of the animals.

Special restrictions for use and special conditions for use:

Not applicable.

### Major incompatibilities:

Do not mix with any other veterinary medicinal product

## 7. Adverse events

Pigs

Very common (>1 animal / 10 animals treated):	- Injection site induration
Common (1 to 10 animals / 100 animals treated):	- Injection site swelling* - Injection site reddening. - Elevated temperature**

\* with a diameter of 10 cm which spontaneously subside within 3 to 14 days.

\*\* up to 0.8°C for 1 or 2 days after injection



Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system.

#### **8. Dosage for each species, routes and method of administration**

Vaccination: Piglets from the age of 6 weeks are vaccinated with 2 doses of 1.0 ml, 3 weeks apart.

Administration route: Intramuscular, preferably to the parauricular area.

#### **9. Advice on correct administration**

Before use, allow the vaccine to reach room temperature of 15 to 25 °C and shake well.

#### **10. Withdrawal periods**

Zero days

#### **11. Special storage precautions**

Keep out of the sight and reach of children.

Store and transport refrigerated (2 °C – 8 °C).

Do not freeze.

Protect from light.

Do not use this veterinary medicinal product after the expiry date which is stated on the label EXP.

The expiry date refers to the last day of that month.

Shelf life after first opening the container: 10 hours.

#### **12. Special precautions for disposal**

Medicines should not be disposed of via wastewater or household waste.

#### **13. Classification of veterinary medicinal products**

Veterinary medicinal product subject to prescription.

#### **14. Marketing authorisation numbers and pack sizes**

Marketing authorisation numbers

Cardboard box:

1 x 10 ml (1 x 10 doses) in 10 ml glass vial (class I),  
1 x 10 ml (1 x 10 doses) in 15 ml HDPE plastic vial,  
10 x 10 ml (10 x 10 doses) in 10 ml glass vial (class I),  
1 x 50 ml (1 x 50 doses) in 50 ml glass vial (class II),  
1 x 50 ml (1 x 50 doses) in 60 ml HDPE plastic vial,  
1 x 100 ml (1 x 100 doses) in 100 ml glass vial (class II),  
1 x 100 ml (1 x 100 doses) in 120 ml HDPE plastic vial,  
1 x 250 ml (1 x 250 doses) in 250 ml HDPE plastic vial

Plastic box:

10 x 10 ml (10 x 10 doses) in 10 ml glass vial (class I)

Not all pack sizes may be marketed.

#### **15. Date on which the package leaflet was last revised**

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

#### **16. Contact details**

Marketing authorisation holder:

VIRBAC

1ère avenue – 2065 m – L.I.D.

06516 Carros Cedex

France

Telephone: +33 492 08 73 04

Manufacturer responsible for batch release:

Bioveta a.s.

Komenského 212/12

683 23 Ivanovice na Hané

Czechia

Telephone: 00420 517 318 500

Local representative and contact details to report suspected adverse reactions:

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.