

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Noromectin 10 mg/ml Solution for Injection for Cattle and Pigs [IE] [EL] [FR] [IS][IT]
[NL] [PT] [ES]

Noromectin Injection 10mg/ml Solution for Injection for Cattle and Pigs. [DE]

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml contains:

Active substance:

Ivermectin 10 mg

Excipients:

| Qualitative composition of excipients and other constituents |
|--|
| Glycerol Formal |
| Polyethylene Glycol 200 |

A clear, colourless solution.

3. CLINICAL INFORMATION

3.1 Target species

Cattle and pigs.

3.2 Indications for use for each target species

Cattle

Treatment of infections by the following parasites:

Gastrointestinal roundworms (adults and fourth stage larvae):

Ostertagia ostertagi (including inhibited *O. ostertagi*),

Ostertagia lyrata,

Haemonchus placei,

Trichostrongylus axei,

Trichostrongylus colubriformis,

Cooperia oncophora,

Cooperia punctata,

Cooperia pectinata,

Oesophagostomum radiatum,

Nematodirus helvetianus (adult).

Lungworms (adult and fourth stage larvae):

Dictyocaulus viviparus.

Warbles:

Hypoderma bovis,
Hypoderma lineatum.

Sucking Lice:

Linognathus vituli,
Haematopinus eurysternus,

Mange Mites:

Psoroptes communis var *bovis*, *Sarcoptes scabiei* var *bovis*.

The veterinary medicinal product may also be used to reduce infection of the mange mite *Chorioptes bovis*, but complete elimination may not occur.

Pigs

Treatment of infections by the following parasites:

Gastrointestinal roundworms:

Ascaris suum (adults and fourth-stage larvae),
Hyoststrongylus rubidus (adults and fourth-stage larvae),
Oesophagostomum spp. (adults and fourth-stage larvae),
Strongyloides ransomi (adults).

Lungworms:

Metastrongylus spp (adults).

Lice:

Haematopinus suis.

Mange mites:

Sarcoptes scabiei var *suis*.

3.3 Contraindications

Do not use in dogs or cats as severe adverse reactions may occur.

The product is not for intravenous or intramuscular use.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

3.4 Special warnings

In cattle: to avoid secondary reactions due to the death of *Hypoderma* larvae in the oesophagus or in the spine it is recommended to administer the veterinary medicinal product at the end of warble fly activity and before the larvae reach their resting sites. Consult your veterinarian on the correct timing of treatment.

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.
- Underdosing, which may be due to underestimation of body weight, misadministration of the product, or lack of calibration of the dosing device.

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, anthelmintic belonging to another pharmacological class and having a different mode of action should be used.

Resistance to ivermectin has been reported in *Ostertagia ostertagi* in cattle. Therefore, the use of this veterinary medicinal product should be based on local (regional, farm) epidemiological information about susceptibility of this helminth species and recommendations on how to limit further selection for resistance to anthelmintics

3.5 Special precautions for use

Special precautions for safe use in the target species:

Since ivermectin is highly bound to plasma proteins, special care should be taken in cases of sick animals or in nutritional conditions associated with low plasma protein levels.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Do not smoke or eat while handling the product.

Direct contact of the product with the skin should be avoided.

Wash hands after use.

Take care to avoid self-injection. Inadvertent self-injection may result in local irritation and/or pain at the injection site.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Special precautions for the protection of the environment:

Ivermectin is extremely dangerous to fish and aquatic life.

Other precautions:

Avermectins may not be well tolerated in all non-target species (cases of intolerance with fatal outcome are reported in dogs – especially collies, old English sheepdogs and related breeds or crosses, and also in turtles/tortoises).

3.6 Adverse events

Cattle:

| | |
|--|--------------------------------------|
| Common (1 to 10 animals / 100 animals treated): | Injection site swelling ¹ |
|--|--------------------------------------|

| | |
|--|-------------------------|
| Very rare (<1 animal / 10,000 animals treated, including isolated reports): | Discomfort ² |
|--|-------------------------|

¹ Soft tissue swellings which disappear without treatment.

² Transient.

Pigs:

| | |
|--|--|
| Very rare (<1 animal / 10,000 animals treated, including isolated reports): | Pain ¹ Injection site swelling ¹ |
|--|--|

¹ Mild and transient. Disappears without treatment.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Pregnancy and lactation:

Can be used during pregnancy and lactation in sows and in cattle provided that the milk is not intended for human consumption. See also section 3.12.

3.8 Interaction with other medicinal products and other forms of interaction

The effects of GABA agonists are increased by ivermectin.

3.9 Administration routes and dosage

Subcutaneous use.

For single administration only.

To ensure administration of a correct dose, body weight should be determined as accurately as possible. Accuracy of the dosing device should be checked.

If animals are to be treated collectively rather than individually, they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- or over-dosing.

Cattle

Ivermectin should be administered at a dosage rate of 200 micrograms per kg bodyweight (1 ml/50 kg). It should be injected subcutaneously in front of or behind the shoulder using aseptic technique. A sterile 17-gauge, 12.7 mm needle (equivalent to half inch needle) is

recommended. Use of a draw-off needle is recommended to avoid excess broaching of the stopper.

Pigs

The veterinary medicinal product should be administered at a dosage rate of 300 micrograms per kg bodyweight (1 ml/33 kg). It should be injected subcutaneously into the neck using aseptic technique. A sterile 17-gauge, 12.7 mm needle (equivalent to half inch needle) is recommended. Exact dosing is important especially in pigs with low bodyweight, therefore a syringe capable of dosing in 0.1 ml steps should be used.

The treatment schedule should be based on the local epidemiological situation.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

In the case of overdose a symptomatic treatment should be given. The symptoms of overdose can be trembling convulsions and coma.

In cattle, single dose of 4.0 mg ivermectin per kg (20 times the use level) given subcutaneously resulted in ataxia and depression.

No systemic or local signs of toxic effects were reported at 3 times the recommended dose in both species.

Ivermectin has a recognised wide safety margin in swine. A dose of 30 mg ivermectin per kg (100 x the recommended dose of 0.3 mg per kg) injected subcutaneously to pigs caused lethargy, ataxia, bilateral mydriasis, intermittent tremors, laboured breathing and lateral recumbency.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Cattle

Meat and offal: 49 days.

Not authorised for use in animals producing milk for human consumption. Do not use in non-lactating dairy cows including pregnant dairy heifers within 60 days of calving.

Pigs

Meat and offal: 18 days.

4. PHARMACOLOGICAL INFORMATION

4.1 ATC vet code:

QP54AA01

4.2 Pharmacodynamics

Ivermectin is a macrocyclic lactone derivative and acts by inhibiting nerve impulses. It binds selectively and with high affinity to glutamate-gated chloride ion channels which occur in invertebrate nerve and muscle cells. This leads to an increase in the permeability of the cell membrane to chloride ions with hyperpolarisation of the nerve or muscle cell, resulting in paralysis and death of the relevant parasites. Compounds of this class may also interact with other ligand gated chloride channels, such as those gated by the neurotransmitter gamma-aminobutyric acid (GABA). The margin of safety for compounds of this class is attributable to the fact that mammals do not have glutamate-gated chloride channels. The macrocyclic lactones have a low affinity for other mammalian ligand-gated chloride channels, and they do not readily cross the blood-brain barrier.

4.3 Pharmacokinetics

After subcutaneous administration of the recommended dose of the product to cattle (200 µg/kg), the following parameters were observed: C_{max} of 37 ng/ml and AUC of 7558 ng/ml.h. After subcutaneous administration of the recommended dose of the product to pigs (300 µg/kg), the following parameters were observed: C_{max} of 14 ng/ml, and AUC of 1887 ng/ml.h. Ivermectin is only partially metabolised. In cattle, only about 1-2% is excreted in the urine the remainder is excreted in the faeces, approximately 60% of which is excreted as unaltered drug. The remainder is excreted as metabolites or degradation products. Biliary excretion, followed by elimination in faeces is probably the major route of ivermectin excretion in pigs.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.
Shelf life after first opening of the immediate packaging: 28 days.

5.3 Special precautions for storage

Store below 25 °C.
Protect from direct sunlight.

5.4 Nature and composition of immediate packaging

The product will be supplied in 50 ml, 100 ml, 250 ml, 500 ml and 1 L volumes, presented in high-density polyethylene vials with bromobutyl bungs and aluminium caps.

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

The veterinary medicinal product should not enter water courses as ivermectin is extremely dangerous for fish and other aquatic organisms. Do not contaminate surface waters or ditches with product or used containers.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

Norbrook Laboratories (Ireland) Limited

7. MARKETING AUTHORISATION NUMBER(S)

8. DATE OF THE FIRST AUTHORISATION

Date of first authorisation: 13 March 2000.

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

{DD/MM/YYYY}

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the United Product Database (<https://medicines.health.europa.eu/veterinary>).

ANNEX III

LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

{CARTON}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Noromectin 10 mg/ml Solution for Injection [IE] [EL] [FR] [IS][IT]
[NL] [PT] [ES]

Noromectin Injection 10mg/ml Solution for Injection [DE]

2. STATEMENT OF ACTIVE SUBSTANCES

Each ml contains:

Active substance:

Ivermectin 10 mg

3. PACKAGE SIZE

50 ml,
100 ml,
250 ml,
500 ml,
1000 ml

4. TARGET SPECIES

Cattle and pigs.

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Subcutaneous use.

For single administration only.

To ensure administration of a correct dose, body weight should be determined as accurately as possible. Accuracy of the dosing device should be checked.

If animals are to be treated collectively rather than individually, they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- or over- dosing.

Cattle:

The veterinary medicinal product should be given only by subcutaneous injection, in front of or behind the shoulder using aseptic technique. at the recommended dosage level of 1 ml per 50 kg bodyweight. A sterile 17-gauge, 12.7 mm needle (equivalent to half inch needle) is recommended. Use of a draw-off needle is recommended to avoid excess broaching of the stopper.

The table below indicates the recommended dose volumes for various bodyweights and the number of doses/pack, for example:

| Bodyweight (kg) | Dose Volume (ml) |
|---|-------------------------|
| Up to 50 | 1 |
| 51 – 100 | 2 |
| 101 – 150 | 3 |
| 151 – 200 | 4 |
| 201 – 250 | 5 |
| 251 – 300 | 6 |
| Over 300 kg bodyweight give 1 ml per 50 kg bodyweight | |

Pigs:

The veterinary medicinal product should be given only by subcutaneous injection in the neck at the recommended dosage level of 1 ml per 33 kg bodyweight. The use of a sterile 17-gauge, 12.7 mm needle (equivalent to half inch needle) is recommended. Exact dosing is important especially in pigs with low bodyweight, therefore a syringe capable of dosing in 0.1 ml steps should be used.

The table below indicates the recommended dose volumes for various bodyweights and the number of doses/pack. In pigs, especially those under 16 kg for which less than 0.5 ml of veterinary medicinal product is indicated, dosing accuracy is important. For piglets weighing less than 16 kg give at 0.1 ml per 3 kg bodyweight.

| Bodyweight (kg) | Dose Volume (ml) |
|--|-------------------------|
| 16 | 0.5 |
| 33 | 1.0 |
| 50 | 1.5 |
| 66 | 2.0 |
| 99 | 3.0 |
| 133 | 4.0 |
| 166 | 5.0 |
| 200 | 6.0 |
| Over 200 kg bodyweight give 1 ml per 33 kg bodyweight. | |

Exact dosing is important especially in pigs with low bodyweight, therefore a syringe capable of dosing in 0.1 ml steps should be used.

The treatment schedule should be based on the local epidemiological situation.

| |
|------------------------------|
| 7. WITHDRAWAL PERIODS |
|------------------------------|

Withdrawal period:

Cattle:

Meat and offal: 49 days.

Not authorised for use in animals producing milk for human consumption. Do not use in non-lactating dairy cows including pregnant heifers within 60 days of calving.

Pigs:

Meat and offal: 18 days.

| |
|-----------------------|
| 8. EXPIRY DATE |
|-----------------------|

Exp. {mm/yyyy}

Once opened use within 28 days.

| |
|---------------------------------------|
| 9. SPECIAL STORAGE PRECAUTIONS |
|---------------------------------------|

Store below 25 °C.

Protect from direct sunlight.

| |
|--|
| 10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE” |
|--|

Read the package leaflet before use.

| |
|--|
| 11. THE WORDS “FOR ANIMAL TREATMENT ONLY” |
|--|

For animal treatment only.

| |
|--|
| 12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN” |
|--|

Keep out of the sight and reach of children.

| |
|---|
| 13. NAME OF THE MARKETING AUTHORISATION HOLDER |
|---|

Norbrook Laboratories (Ireland) Limited

| |
|--|
| 14. MARKETING AUTHORISATION NUMBERS |
|--|

| |
|-------------------------|
| 15. BATCH NUMBER |
|-------------------------|

Lot {number}

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

{VIAL}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Noromectin 10 mg/ml Solution for Injection [IE] [EL] [FR] [IS][IT]
[NL] [PT] [ES]

Noromectin Injection 10mg/ml Solution for Injection [DE]

2. STATEMENT OF ACTIVE SUBSTANCES

Each ml contains:

Active substance:

Ivermectin 10 mg

3. TARGET SPECIES

Cattle and pigs.

4. ROUTES OF ADMINISTRATION

Subcutaneous use.

Read the package leaflet before use.

5. WITHDRAWAL PERIODS

Withdrawal period:

Cattle:

Meat and offal: 49 days.

Not authorised for use in animals producing milk for human consumption. Do not use in non-lactating dairy cows including pregnant heifers within 60 days of calving.

Pigs:

Meat and offal: 18 days.

6. EXPIRY DATE

Exp. {mm/yyyy}

Once opened use within 28 days.

7. SPECIAL STORAGE PRECAUTIONS

Store below 25 °C.
Protect from direct sunlight.

| |
|--|
| 8. NAME OF THE MARKETING AUTHORISATION HOLDER |
|--|

Norbrook Laboratories (Ireland) Limited

| |
|------------------------|
| 9. BATCH NUMBER |
|------------------------|

Lot {number}

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Noromectin 10 mg/ml Solution for Injection for Cattle and Pigs [IE] [EL] [FR] [IS][IT] [NL] [PT] [ES]

Noromectin Injection 10mg/ml Solution for Injection for Cattle and Pigs. [DE]

2. Composition

Each ml contains:

Active substance:

Ivermectin 10 mg

A clear, colourless solution.

3. Target Species

Cattle and Pigs.

4. Indications for use

Cattle:

Treatment of infections by the following parasites:

Gastrointestinal roundworms (adults and fourth stage larvae):

Ostertagia ostertagi (including inhibited *O. ostertagi*), *Ostertagia lyrata*, *Haemonchus placei*, *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Cooperia oncophora*, *Cooperia punctata*, *Cooperia pectinata*, *Oesophagostomum radiatum*, *Nematodirus helvetianus* (adult).

Lungworms (adults and fourth stage larvae):

Dictyocaulus viviparus.

Warbles (parasitic stages):

Hypoderma bovis, *Hypoderma lineatum*.

Sucking lice:

Linognathus vituli, *Haematopinus eurysternus*.

Mange mites:

Psoroptes communis var *bovis*, *Sarcoptes scabiei* var *bovis*.

The veterinary medicinal product may also be used to reduce infection of the mange mites *Chorioptes bovis* but complete elimination may not occur.

Pigs:

Treatment of infections by the following parasites:

Gastrointestinal roundworms:

Ascaris suum (adults and fourth stage larvae)

Hyoststrongylus rubidus (adults and fourth stage larvae)

Oesophagostomum spp (adults and fourth stage larvae)

Strongyloides ransomi (adults)

Lungworms:

Metastrongylus spp (adults)

Lice:

Haematopinus suis

Mange mites:

Sarcoptes scabiei var *suis*

5. Contraindications

Do not use in dogs or cats as severe adverse reactions may occur.

The product is not for intravenous or intramuscular use.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

6. Special Warnings

Special warnings:

In cattle: to avoid secondary reactions due to the death of *Hypoderma* larvae in the oesophagus or in the spine it is recommended to administer the veterinary medicinal product at the end of warble fly activity and before the larvae reach their resting sites. Consult your veterinarian on the correct timing of treatment.

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.
- Underdosing, which may be due to underestimation of body weight, misadministration of the product, or lack of calibration of the dosing device.

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, anthelmintic belonging to another pharmacological class and having a different mode of action should be used.

Resistance to ivermectin has been reported in *Ostertagia ostertagi* in cattle. Therefore, the use of this veterinary medicinal product should be based on local (regional, farm)

epidemiological information about susceptibility of this helminth species and recommendations on how to limit further selection for resistance to anthelmintics.

Special precautions for safe use in the target species:

Since ivermectin is highly bound to plasma proteins, special care should be taken in cases of sick animals or in nutritional conditions associated with low plasma protein levels.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Do not smoke or eat while handling the product.

Direct contact of the product with the skin should be avoided.

Wash hands after use.

Take care to avoid self-injection. Inadvertent self-injection may result in local irritation and/or pain at the injection site.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Special precautions for the protection of the environment:

Ivermectin is extremely dangerous to fish and aquatic life.

Other precautions:

Avermectins may not be well tolerated in all non-target species (cases of intolerance with fatal outcome are reported in dogs – especially collies, old English sheepdogs and related breeds or crosses, and also in turtles/tortoises).

Pregnancy and lactation:

Can be used during pregnancy and lactation in sows and in cattle provided that the milk is not intended for human consumption. See also section ‘Withdrawal periods’.

Interaction with other medicinal products and other forms of interaction:

The effects of GABA agonists are increased by ivermectin.

Overdose:

In the case of overdose a symptomatic treatment should be given. The symptoms of overdose can be trembling convulsions and coma.

In cattle, single dose of 4.0 mg ivermectin per kg (20 times the use level) given subcutaneously resulted in ataxia and depression.

No systemic or local signs of toxic effects were reported at 3 times the recommended dose in both species.

Ivermectin has a recognised wide safety margin in swine. A dose of 30 mg ivermectin per kg (100 x the recommended dose of 0.3 mg per kg) injected subcutaneously to pigs caused lethargy, ataxia, bilateral mydriasis, intermittent tremors, laboured breathing and lateral recumbency.

Major incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

7. Adverse events

Cattle:

| | |
|---|--------------------------------------|
| Common (1 to 10 animals / 100 animals treated): | Injection site swelling ¹ |
| Very rare (<1 animal / 10,000 animals treated, including isolated reports): | Discomfort ² |

¹ Soft tissue swellings which disappear without treatment.

² Transient.

Pigs:

| | |
|---|--|
| Very rare (<1 animal / 10,000 animals treated, including isolated reports): | Pain ¹ Injection site swelling ¹ |
|---|--|

¹ Mild and transient. Disappears without treatment.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. If you notice any side effects, even those not already listed in this package leaflet, or if you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via the national reporting system: {national system details}

8. Dosage for each species, routes and method of administration

Subcutaneous use.

For single administration only.

To ensure administration of a correct dose, body weight should be determined as accurately as possible. Accuracy of the dosing device should be checked.

If animals are to be treated collectively rather than individually, they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- or over- dosing.

Cattle:

The veterinary medicinal product should be administered at a dosage rate of 200 microgram/kg bodyweight (1 ml per 50 kg). It should be injected subcutaneously in front of or behind the shoulder using aseptic technique. A sterile 17-gauge, 12.7 mm needle (equivalent to half inch needle) is recommended. Use of a draw-off needle is recommended to avoid excess broaching of the stopper.

For example:

| Bodyweight (kg) | Dose Volume (ml) |
|---|------------------|
| Up to 50 | 1 |
| 51 – 100 | 2 |
| 101 – 150 | 3 |
| 151 – 200 | 4 |
| 201 – 250 | 5 |
| 251 – 300 | 6 |
| Over 300 kg bodyweight give 1 ml per 50 kg bodyweight | |

Pigs:

The veterinary medicinal product should be administered at a dosage rate of 300 microgram/kg bodyweight (1 ml per 33 kg bodyweight). It should be injected subcutaneously into the neck using aseptic technique. A sterile 17-gauge, 12.7 mm needle (equivalent to half inch needle) is recommended.

| Bodyweight (kg) | Dose Volume (ml) |
|---|------------------|
| 16 | 0.5 |
| 33 | 1.0 |
| 50 | 1.5 |
| 66 | 2.0 |
| 99 | 3.0 |
| 133 | 4.0 |
| 166 | 5.0 |
| 200 | 6.0 |
| Over 200 kg bodyweight give 1 ml per 33 kg bodyweight | |

9. Advice on correct administration

The treatment schedule should be based on the local epidemiological situation and according to veterinary advice.

Cattle:

Warbles: The best time to treat is in *late* autumn or early winter, before the small migrating larvae have time to cause serious damage.

Pigs:

For effective mange control, care must be taken to prevent reinfestation from exposure to untreated animals or contaminated facilities.

Since louse eggs are unaffected by ivermectin and may take up to 3 weeks to hatch, complete elimination may not occur following a single injection.

Exact dosing is important especially in pigs with low bodyweight, therefore a syringe capable of dosing in 0.1 ml steps should be used.

10. Withdrawal periods

Cattle

Meat and offal: 49 days.

Not authorised for use in animals producing milk for human consumption. Do not use in non-lactating dairy cows including pregnant dairy heifers within 60 days of calving.

Pigs

Meat and offal: 18 days.

11. Special storage precautions

Keep out of the sight and reach of children.

Protect from direct sunlight.

Store below 25 °C.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton. The expiry date refers to the last day of that month.

Shelf life after first opening the immediate packaging: 28 days.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

This veterinary medicinal product should not enter water courses as ivermectin is extremely dangerous for fish and other aquatic organisms. Do not contaminate surface waters or ditches with product or used containers.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack size

The product will be supplied in 50 ml, 100 ml, 250 ml, 500 ml and 1 L volumes, presented in high-density polyethylene vials with bromobutyl bungs and aluminium caps.

Not all pack sizes may be marketed.

15. Date on which the package leaflet was last revised

{MM/YYYY}

Detailed information on this veterinary medicinal product is available in the United Product Database (<https://medicines.health.europa.eu/veterinary>).

16. Contact details

Marketing authorisation holder and contact details to report suspected adverse reactions:

Norbrook Laboratories (Ireland) Limited
Rossmore Industrial Estate
Monaghan
Ireland
Tel: +44 (0)28 3026 4435
E-mail: phvdept@norbrook.co.uk

Manufacturer responsible for batch release:

| | |
|-----------------------------|-------------------------------|
| Norbrook Manufacturing Ltd. | Norbrook Laboratories Limited |
| Rossmore Industrial Estate | Station Works |
| Monaghan | Newry, Co. Down |
| Ireland | BT35 6JP |

Local representatives and contact details to report suspected adverse reactions:

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.