

PACKAGE LEAFLET:

TRIDERM cutaneous spray solution for dogs

TRIPOFLOX cutaneous spray solution for dogs

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder: Organit Kft., Homoksor 7., Székesfehérvár, H-8000, Hungary

Manufacturer responsible for batch release: ALPHAVET Zrt., Köves János út 13., Bábolna, H-2943, Hungary

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

TRIDERM cutaneous spray solution for dogs (CY, ES, EL, HR, LT, LV, MT, PT)

TRIPOFLOX cutaneous spray solution for dogs (PL)

Marbofloxacin, Ketoconazole, Prednisolone

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each ml contains:

Active substances:

Marbofloxacin..... 1.025 mg

Ketoconazole..... 2.041 mg

Prednisolone 0.926 mg

Excipients: q.s

Yellowish-slightly opal solution.

4. INDICATION(S)

Treatment of acute dermatitis when mixed infection caused by *Pseudomonas aeruginosa* or *Staphylococcus pseudintermedius* susceptible to marbofloxacin and *Malassezia pachydermatis* susceptible to ketoconazole is demonstrated. The veterinary medicinal product should be used based on susceptibility testing on the bacteria isolated from the animal.

5. CONTRAINDICATIONS

Do not use in case of hypersensitivity to the active substances or to any of the excipients.

6. ADVERSE REACTIONS

Mild erythematous lesions have been observed following the application. The frequency of adverse reactions is very rare (less than 1 animal in 10.000 animals treated, including isolated reports).

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1.000 animals treated)
- rare (more than 1 but less than 10 animals in 10.000 animals treated)
- very rare (less than 1 animal in 10.000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon. Alternatively you can report via your national reporting system. For details regarding the national system please contact NCA.

7. TARGET SPECIES

Dogs

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Cutaneous use. Shake well before use.

The recommended dosage is 2.26-9.18 µg of marbofloxacin, 4.52-18.36 µg of ketoconazole and 2.08-8.45 µg of prednisolone per cm² of affected skin per day. This dosage can be achieved with two pump spray activations (equivalent to approximately 0.2 ml/ treatment) over a surface to be treated equivalent to a square of 5 cm x 5 cm when spraying from a distance of about 10 cm; of 10 cm x 10 cm when spraying from a distance of about 30 cm. Repeat the application twice a day for 7-14 days, depending on clinical and microbiological healing. Before the application of the veterinary medicinal product the hair or dirt on the treated surface has to be removed.

Period of treatment depends on clinical convalescence of skin inflammations of bacterial and of fungal origin. In case the dog treated doesn't recover until 7th day the treatment should be followed until 14th day. In those cases in which the treatment was extended to 14 days and dog still didn't recover within 14 days, it is recommendable to change to another, adequate veterinary medicinal product.

9. ADVICE ON CORRECT ADMINISTRATION

10. WITHDRAWAL PERIOD(S)

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the label, and carton after EXP. The expiry date refers to the last day of that month.

Shelf life after first opening the container: 28 days

12. SPECIAL WARNING(S)

None.

Special warnings for each target species:

Collar should be fixed on the treated dogs in order to prevent licking. Keep the animals to be treated separated from each other in order to prevent licking each other. Bacterial and fungal

dermatitis is often secondary in nature and appropriate diagnosis should be used to determine the primary factors involved.

Unnecessary use of pharmacologically active substance in terms of any active substance should be avoided. Treatment is indicated only if mixed infection with *Pseudomonas aeruginosa* or *Staphylococcus pseudintermedius* and *Malassezia pachydermatis* has been proved. If one of the pharmacologically active substances is no longer indicated due to the different characteristics of bacterial and fungal infections, the application of pharmacologically active substance should be discontinued and replaced by an appropriate treatment option.

Special precautions for use in animals:

If hypersensitivity to any of the active components occurs, treatment should be discontinued and appropriate therapy should be instituted.

Use of the veterinary medicinal product should be based on identification of infecting organisms and susceptibility testing and take into account official and local antimicrobial policies.

Heavy reliance on a single class of antibiotic may result in the induction of resistance in a bacterial population. It is prudent to reserve fluoroquinolones for the treatment of clinical conditions, which have responded poorly or are expected to respond poorly to other classes of antibiotics. However, microbiological diagnosis and sensitivity test must be performed.

Prolonged and intensive use of topical corticosteroids preparation is known to trigger local and systemic effects, including suppression of adrenal function, thinning of the epidermis and delayed healing.

Spraying on open lesions and wounds is must be avoided.

During the administration do not bath or shampoo the animals.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

As with all medicinal formulation operators are advised to use the product according to label instructions.

Flammable solution. Do not spray on naked flame or any incandescent material.

Do not smoke, drink or eat while handling the product.

Do not inhale spray mist. Use only in well ventilated areas

Several components of the product may cause hypersensitivity reactions as well as skin and/or eye irritation.

People with known hypersensitivity to (fluoro)quinolones, ketoconazole, prednisolone or any of the excipients should avoid contact with the veterinary medicinal product.

Avoid contact with skin or eyes. In case of accidental spillage, skin or eyes should be rinsed immediately with plenty of water.

Wash hands after use.

Seek medical advice if signs of cutaneous erythema, exanthema, or persistent ocular irritation appear after exposure. Swelling of face, lips and eyes, or respiratory difficulties are more serious signs that need urgent medical action.

In case of accidental ingestion seek medical advice immediately and show the package leaflet or the label to the physician.

Treated animals should not be handled, and children should not be allowed to play with treated animals until the fur is dry.

Animals treated should not be allowed to sleep with owners, especially children.

Product should be kept out of sight and reach of children.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy and lactation. The use is not recommended during pregnancy and lactation.

Interaction with other medicinal products and other forms of interaction:

No data available.

Overdose (symptoms, emergency procedures, antidotes):

At 5 times of the recommended dose, no local or general adverse reactions were observed.

Incompatibilities:

Not applicable.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Medicines should not be disposed of via wastewater or household waste.

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

15. OTHER INFORMATION

Pack size:

Box with 1 bottle of 30 ml

Dispensing conditions: Veterinary medicinal product subject to veterinary prescription.

Administration conditions: Administration under the control or direct responsibility of a veterinary surgeon.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.