

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Cortavance 0.584 mg/ml cutaneous spray solution for dogs

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml contains:

Active substance:

Hydrocortisone aceponate 0.584 mg

Excipient:

Qualitative composition of excipients and other constituents
Propylene glycol methyl ether

Clear colourless or very slightly yellow solution.

3. CLINICAL INFORMATION

3.1 Target species

Dogs.

3.2 Indications for use for each target species

For symptomatic treatment of inflammatory and pruritic dermatoses in dogs.
For alleviation of clinical signs associated with atopic dermatitis in dogs.

3.3 Contraindications

Do not use on cutaneous ulcers.
Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

3.4 Special warnings

None.

3.5 Special precautions for use

Special precautions for safe use in the target species:

Clinical signs of atopic dermatitis such as pruritus and skin inflammation are not specific for this disease and therefore other causes of dermatitis such as ectoparasitic infestations and infections which cause dermatological signs should be ruled out before treatment is started, and underlying causes should be investigated.

In the case of concurrent microbial disease or parasitic infestation, the dog should receive appropriate treatment for such condition.

In the absence of specific information, the use in animal suffering from Cushing's syndrome shall be based on the risk-benefit assessment.

Since glucocorticosteroids are known to slow growth, use in young animals (under 7 months of age) shall be based on the risk-benefit assessment and subject to regular clinical evaluations.

Total body surface treated should not exceed approximately 1/3 of the dog's surface corresponding for example to a treatment of two flanks from the spine to the mammary chains including the shoulders and the thighs (see also section 3.10). Otherwise, use only according to the risk-benefit assessment of the responsible veterinary surgeon and subject the dog to regular clinical evaluations as further described in section 3.9.

Care should be taken to avoid spraying into the eyes of the animal.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

The active substance is potentially pharmacologically active at high doses of exposure.
The formulation may cause eye irritation following accidental ocular contact.
The formulation is flammable.

Wash hands after use. Avoid contact with eyes.
To avoid skin contact, recently treated animals should not be handled until the application site is dry.
To avoid inhalation of the product, apply the spray in a well-ventilated area.
Do not spray on naked flame or any incandescent material.
Do not smoke while handling the veterinary medicinal product.
Replace the bottle in the outer carton and in a safe place out of the sight and the reach of children immediately after use.

In case of accidental skin contact, avoid hand-to-mouth contact and wash the exposed area immediately with water.
In case of accidental eye contact, rinse with abundant quantities of water.
If eye irritation persists, seek medical advice.
In case of accidental ingestion, especially by children, seek medical advice immediately and show the leaflet or the label to the physician.

Special precautions for the protection of the environment:

Not applicable.

Other precautions:

The solvent in this product may stain certain materials including painted, varnished or other household surfaces or furnishings. Allow the application site to dry before permitting contact with such materials.

3.6 Adverse events

Dogs:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Application site pruritus ¹ Application site erythema ¹
--	--

¹Transient local

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing

authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

The safety of the veterinary medicinal product has not been established during pregnancy and lactation.

Systemic absorption of hydrocortisone aceponate being negligible, it is unlikely for teratogenic, foetotoxic, maternotoxic effects to happen at the recommended dosage in dogs.

Pregnancy and lactation:

Use only according to the benefit-risk assessment by the responsible veterinarian.

3.8 Interaction with other medicinal products and other forms of interaction

No data available.

In the absence of information, it is recommended not to apply other topical preparations simultaneously on the same lesions.

3.9 Administration routes and dosage

Cutaneous use.

Before administration, screw the pump spray on the bottle.

The veterinary medicinal product is then applied by activating the pump spray, from a distance of about 10 cm of the area to be treated.

The recommended dosage is 1.52 mcg of hydrocortisone aceponate/cm² of affected skin per day. This dosage can be achieved with two pump spray activations over a surface to be treated equivalent to a square of 10 cm x 10 cm.

- For treatment of inflammatory and pruritic dermatoses, repeat the treatment daily for 7 consecutive days.

In case of conditions requiring an extended treatment, the responsible veterinarian should subject the use of the veterinary medicinal product to the risk-benefit assessment.

If signs fail to improve within 7 days, treatment should be re-evaluated by the veterinarian.

- For alleviation of clinical signs associated with atopic dermatitis, repeat the treatment daily for at least 14 and up to 28 consecutive days.

An intermediary control by the veterinarian at day 14 should be made to decide if further treatment is needed. The dog should be re-evaluated regularly with regard to hypothalamic-pituitary axis (HPA) suppression or skin atrophy, both being possibly asymptomatic.

Any prolonged use of this product, to control atopy, should be at the benefit risk assessment of the responsible veterinary surgeon. It should take place after a re-evaluation of the diagnosis and also a consideration of the multi-modal treatment plan in the individual animal.

Presented as a volatile spray, this veterinary medicinal product does not require any massage.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

Tolerance studies of multiple doses were assessed over a period of 14 days in healthy dogs using 3 and 5 times the recommended dosage corresponding to the two flanks, from the spine to the mammary chains including the shoulder and the thighs (1/3 of the dog's body surface area). These resulted in a reduced capacity for production of cortisol that is fully reversible within 7 to 9 weeks after the end of treatment.

In 12 dogs suffering from atopic dermatitis, after topical application once a day at the recommended therapeutic dosage for 28 to 70 (n=2) consecutive days, no noticeable effect on the systemic cortisol level was observed.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Not applicable.

4. PHARMACOLOGICAL INFORMATION

4.1 ATCvet code: QD07AC16.

4.2 Pharmacodynamics

The veterinary medicinal product contains the active substance hydrocortisone aceponate. Hydrocortisone aceponate is a dermocorticoid with a potent intrinsic glucocorticoid activity which means a relief of both inflammation and pruritus leading to a quick improvement of skin lesions observed in case of inflammatory and pruritic dermatosis. In case of atopic dermatitis, improvement will be slower.

4.3 Pharmacokinetics

Hydrocortisone aceponate belongs to the diesters class of the glucocorticosteroids.

The diesters are lipophilic components ensuring an enhanced penetration into the skin associated to a low plasma availability. Hydrocortisone aceponate thus accumulates in the dog's skin allowing local efficacy at low dosage. The diesters are transformed inside the skin structures. This transformation is responsible for the potency of the therapeutic class. In laboratory animals, hydrocortisone aceponate is eliminated the same way as hydrocortisone (other name for endogenous cortisol) through urine and faeces.

Topical application of diesters results in high therapeutic index: high local activity with reduced systemic secondary effects.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

None known.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

Shelf life after first opening the immediate packaging: 6 months.

5.3 Special precautions for storage

This veterinary medicinal product does not require any special storage conditions.

5.4 Nature and composition of immediate packaging

Box containing a polyethylene terephthalate (PET) or high density polyethylene (HDPE) bottle filled with 31 ml or 76 ml of solution, closed with an aluminium screw cap or a white plastic screw cap and a pump spray.

Carton box with a PET bottle of 31 ml

Carton box with a PET bottle of 76 ml

Carton box with a HDPE bottle of 31 ml

Carton box with a HDPE bottle of 76 ml

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

VIRBAC

7. MARKETING AUTHORISATION NUMBER(S)

EU/2/06/069/001

EU/2/06/069/002

EU/2/06/069/003

EU/2/06/069/004

8. DATE OF FIRST AUTHORISATION

Date of first authorisation: 09/01/2007

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

{DD/YYYY}

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

ANNEX II

OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

None

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

CARTON BOX (31 ml)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Cortavance 0.584 mg/ml cutaneous spray solution

2. STATEMENT OF ACTIVE SUBSTANCES

Each ml contains 0.584 mg of hydrocortisone aceponate

3. PACKAGE SIZE

31 ml

4. TARGET SPECIES

Dogs.

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Cutaneous use.

7. WITHDRAWAL PERIOD(S)

8. EXPIRY DATE

Exp. {mm/yyyy}
Once opened, use by 6 months.

9. SPECIAL STORAGE PRECAUTIONS

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

VIRBAC

14. MARKETING AUTHORISATION NUMBERS

Carton with a PET bottle of 31 ml: EU/2/06/069/002

Carton with a HDPE bottle of 31 ml: EU/2/06/069/003

15. BATCH NUMBER

Lot {number}

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

CARTON BOX (76 ml)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Cortavance 0.584 mg/ml cutaneous spray solution

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Each ml contains 0.584 mg of hydrocortisone aceponate

3. PACKAGE SIZE

76 ml

4. TARGET SPECIES

Dogs.

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Cutaneous use.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {mm/yyyy}
Once opened, use by 6 months.

9. SPECIAL STORAGE PRECAUTIONS

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

VIRBAC

14. MARKETING AUTHORISATION NUMBERS

Carton box with a PET bottle of 76 ml: EU/2/06/069/001

Carton box with a HDPE bottle of 76 ml: EU/2/06/069/004

15. BATCH NUMBER

Lot {number}

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

BOTTLE OF 76 ML

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Cortavance 0.584 mg/ml cutaneous spray solution

2. STATEMENT OF ACTIVE SUBSTANCES

Hydrocortisone aceponate 0.584 mg/ml

3. TARGET SPECIES

Dogs.

4. ROUTES OF ADMINISTRATION

Cutaneous use.
Read the package leaflet before use.

5. WITHDRAWAL PERIODS

6. EXPIRY DATE

Exp. {mm/yyyy}
Once opened, use by 6 months.

7. SPECIAL STORAGE PRECAUTIONS

8. NAME OF THE MARKETING AUTHORISATION HOLDER

VIRBAC

9. BATCH NUMBER

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

BOTTLE OF 31 ML

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Cortavance

2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

0.584 mg/ml

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Cortavance 0.584 mg/ml cutaneous spray solution for dogs

2. Composition

Hydrocortisone aceponate 0.584 mg/ml

Clear colourless or very slightly yellow solution.

3. Target species

Dogs.

4. Indications for use

For symptomatic treatment of inflammatory and pruritic dermatoses in dogs.

For alleviation of clinical signs associated with atopic dermatitis in dogs.

5. Contraindications

Do not use on cutaneous ulcers.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

6. Special warnings

Special precautions for safe use in the target species:

Clinical signs of atopic dermatitis such as pruritus and skin inflammation are not specific for this disease and therefore other causes of dermatitis such as ectoparasitic infestations and infections which cause dermatological signs should be ruled out before treatment is started, and underlying causes should be investigated.

In the case of concurrent microbial disease or parasitic infestation, the dog should receive appropriate treatment for such condition.

In the absence of specific information, the use in animal suffering from Cushing's syndrome shall be based on the risk-benefit assessment.

Since glucocorticosteroids are known to slow growth, use in young animals (under 7 months of age) shall be based on the risk-benefit assessment and subject to regular clinical evaluations.

Total body surface treated should not exceed approximately 1/3 of the dog's surface corresponding for example to a treatment of two flanks from the spine to the mammary chains including the shoulders and the thighs. See also section "Overdose". Otherwise, use only according to the risk-benefit assessment of the responsible veterinary surgeon and subject the dog to regular clinical evaluations as further described in section "Administration routes and dosage".

Care should be taken to avoid spraying into the eyes of the animal.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

The active substance is potentially pharmacologically active at high doses of exposure.
The formulation may cause eye irritation following accidental ocular contact.
The formulation is flammable.

Wash hands after use. Avoid contact with eyes.

To avoid skin contact, recently treated animals should not be handled until the application site is dry.

To avoid inhalation of the product, apply the spray in a well-ventilated area.

Do not spray on naked flame or any incandescent material.

Do not smoke while handling the veterinary medicinal product.

Replace the bottle in the outer carton and in a safe place out of the sight and the reach of children immediately after use.

In case of accidental skin contact, avoid hand-to-mouth contact and wash the exposed area immediately with water.

In case of accidental eye contact, rinse with abundant quantities of water.

If eye irritation persists, seek medical advice.

In case of accidental ingestion, especially by children, seek medical advice immediately and show the leaflet or the label to the physician.

Other precautions:

The solvent in this product may stain certain materials including painted, varnished or other household surfaces or furnishings. Allow the application site to dry before permitting contact with such materials.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy and lactation.

Systemic absorption of hydrocortisone aceponate being negligible, it is unlikely for teratogenic, foetotoxic, maternotoxic effects to happen at the recommended dosage in dogs.

Use only according to the benefit-risk assessment by the responsible veterinarian.

Interaction with other medicinal products and other forms of interaction:

In the absence of information, it is recommended not to apply other topical preparations simultaneously on the same lesions.

Overdose:

Tolerance studies of multiple doses were assessed over a period of 14 days in healthy dogs using 3 and 5 times the recommended dosage corresponding to the two flanks, from the spine to the mammary chains including the shoulder and the thighs (1/3 of the dog's body surface area). These resulted in a reduced capacity for production of cortisol that is fully reversible within 7 to 9 weeks after the end of treatment.

In 12 dogs suffering from atopic dermatitis, after topical application once a day at the recommended therapeutic dosage for 28 to 70 (n=2) consecutive days, no noticeable effect on the systemic cortisol level was observed.

7. Adverse events

Dogs:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):
Application site pruritus ¹ Application site erythema ¹

¹Transient local

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: {national system details}.

8. Dosage for each species, routes and method of administration

Cutaneous use.

Before administration, screw the pump spray on the bottle.

The veterinary medicinal product is then applied by activating the pump spray, from a distance of about 10 cm of the area to be treated.

The recommended dosage is 1.52 mcg of hydrocortisone aceponate/cm² of affected skin per day. This dosage can be achieved with two pump spray activations over a surface to be treated equivalent to a square of 10 cm x 10 cm.

- For treatment of inflammatory and pruritic dermatoses, repeat the treatment daily for 7 consecutive days.

In case of conditions requiring an extended treatment, the responsible veterinarian should subject the use of the veterinary medicinal product to the risk-benefit assessment.

If signs fail to improve within 7 days, treatment should be re-evaluated by the veterinarian.

- For alleviation of clinical signs associated with atopic dermatitis, repeat the treatment daily for at least 14 and up to 28 consecutive days.

An intermediary control by the veterinarian at day 14 should be made to decide if further treatment is needed. The dog should be re-evaluated regularly with regard to HPA suppression or skin atrophy, both being possibly asymptomatic.

Any prolonged use of this product, to control atopy, should be at the benefit risk assessment of the responsible veterinary surgeon. It should take place after a re-evaluation of the diagnosis and also a consideration of the multi-modal treatment plan in the individual animal.

9. Advice on correct administration

Presented as a volatile spray, this veterinary medicinal product does not require any massage.

10. Withdrawal periods

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp. The expiry date refers to the last day of that month.

Shelf life after first opening the immediate packaging: 6 months.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack sizes

EU/2/06/069/001-004

Pack sizes:

Carton box with a PET bottle of 31 ml

Carton box with a PET bottle of 76 ml

Carton box with a HDPE bottle of 31 ml

Carton box with a HDPE bottle of 76 ml

Not all pack sizes may be marketed.

15. Date on which the package leaflet was last revised

{MM/YYYY}

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

16. Contact details

Marketing authorisation holder and manufacturer responsible for batch release:

VIRBAC
1ère Avenue 2065m LID
06516 Carros
France

Local representatives and contact details to report suspected adverse reactions:

Österreich

VIRBAC Österreich GmbH
Hildebrandgasse 27
A-1180 Wien
Österreich
Tel: +43-(0)1 21 834 260

Република България

ЕРГОН МИЛАНОВА ЕООД
с. Бърложница 2222, Софийска област
Република България
Тел: + 359 359888215520
ergonood@gmail.com

Česká republika

VIRBAC Czech Republic s.r.o.
Žitavského 496
156 00 Praha 5
Česká republika
Tel.: +420 608 836 529

Danmark

VIRBAC Danmark A/S
Profilvej 1
6000 Kolding
Danmark
Tel: +45 75521244
virbac@virbac.dk

España

VIRBAC España SA
Angel Guimerá 179-181
ES-08950 Esplugues de Llobregat (Barcelona)
España
Tel.: + 34-(0)93 470 79 40

France

VIRBAC France
13^{ème} rue LID
06517 Carros
service-conso@virbac.fr

Hrvatska

CENTRALNA VETERINARSKA AGENCIJA
d.o.o. (CVA)
Prve Ravnice 2e, 10000 Zagreb
Republika Hrvatska
Tel.: + 385 91 46 55 115
kz@cva.hr

Ireland

VIRBAC IRELAND
McInerney & Saunders
38, Main Street

België/Belgique

VIRBAC Belgium NV
Esperantolaan 4
BE-3001 Leuven
België / Belgique / Belgien
Tel: +32-(0)16 387 260
phv@virbac.be

Κύπρος

VET2VETSUPPLIES LTD
Γαλιλαίου 60
3011 Λεμεσός
Κύπρος
Τηλ: + 357 96116730
info@vet2vetsupplies.com

Deutschland

VIRBAC Tierarzneimittel GmbH
Rögen 20
23843 Bad Oldesloe
Deutschland
Tel: +49-(4531) 805 111

Eesti

OÜ ZOOVETVARU
Uusaru 5
EE-76505 Saue/Harjumaa
Eesti
Tel: + 372 56480207
pv@zoovet.eu

Suomi/Finland

BIOFARM OY
Yrittäjätie 20
FI-03600 Karkkila
Tel: +358-9-225 2560
haittavaikutukset@biofarm.fi

Ελλάδα

VIRBAC ΕΛΛΑΣ ΜΟΝΟΠΡΟΣΩΠΗ Α.Ε.
13^ο χλμ Ε.Ο. Αθηνών - Λαμίας
EL-14452, Μεταμόρφωση
Τηλ.: +30 2106219520
info@virbac.gr

Magyarország

VIRBAC HUNGARY KFT
Váci utca 81. 4 emelet.
HU-1056 Budapest
Tel.: +36703387177
akos.csoman@virbac.hu

Ísland

Vistor hf.
Hörgatún 2,
IS-210 Garðabær

Swords, Co Dublin
K67E0A2
Republic Of Ireland
Tel: +44 (0)-1359 243243

Italia

VIRBAC SRL
Via Ettore Bugatti, 15
IT-20142 Milano
Italia
Tel: + 39 02 40 92 47 1

Luxembourg/Luxemburg

VIRBAC BELGIUM NV
Esperantolaan 4
BE-3001 Leuven
Belgique / Belgien
Tél/Tel: +32-(0)16 387 260
info@virbac.be

Malta

SEROLF TRADING AGENCY LTD.
8, Adelaide Cini Street
Pembroke PBK 1231
Malta
Tel: + 35621337231
info@serolf.com

Norge

VIRBAC Danmark A/S
Profilvej 1
DK-6000 Kolding
Danmark
Tel: + 45 75521244
virbac@virbac.dk

Portugal

VIRBAC de Portugal Laboratórios, Lda.
Rua do Centro Empresarial
Edif.13 - Piso 1- Escrit.3
Quinta da Beloura
PT-2710-693 Sintra (Portugal)
Tel: + 351 219 245 020

Slovenija

MEDICAL INTERTRADE d.o.o.
Brodišče 12, 1236 Trzin
Slovenija
Tel: + 386 1 2529 113
farmakovigilanca@medical-intertrade.si

Sverige

VIRBAC Danmark A/S Filial Sverige
Box 1027
SE-171 21 Solna
Tel: +45 75521244

Sími: +354 535 7000
safety@vistor.is

Lietuva

OŪ ZOOVETVARU
Uusaru 5
EE-76505 Saue/Harjumaa
Estija
Tel: + 372 56480207
pv@zoovet.eu

Latvija

OŪ ZOOVETVARU
Uusaru 5
EE-76505 Saue/Harjumaa
Igaunija
Tel: + 372 56480207
pv@zoovet.eu

Nederland

VIRBAC Nederland BV
Hermesweg 15
3771 ND-Barneveld
Nederland
Tel: +31-(0)342 427 127
phv@virbac.nl

Polska

VIRBAC Sp. z o.o.
ul. Puławska 314
PL-02 - 819 Warszawa
Polska
Tel.: + 48 22 855 40 46

România

Altius SA
Str. Zăgazului nr. 21-25, Corp A, et 8 si 8A
Ap. A.8.2, sect 1, București,
Romania
Tel: + 40 021 310 88 80

Slovenská republika

VIRBAC Czech Republic s.r.o.
Žitavského 496
156 00 Praha 5
Česká republika
Tel.: +420 608 836 529

United Kingdom (Northern Ireland)

VIRBAC IRELAND
McInerney & Saunders
38, Main Street
Swords, Co Dublin

virbac@virbac.dk

K67E0A2
Republic Of Ireland
Tel: +44 (0)-1359 243243

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

17. Other information

Hydrocortisone aceponate administered topically accumulates and is metabolised in skin, as suggested by radioactivity distribution studies and pharmacokinetic data. This results in minimal amounts to reach the blood stream. This particularity will increase the ratio between the desired local anti-inflammatory effect in the skin and the undesirable systemic effects.

Hydrocortisone aceponate applications on the skin lesions provide rapid reduction of the skin redness, irritation and scratching while minimising the general effects.