

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

EVANT suspension and solvent for oral spray for chickens

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each dose (0.007 ml) of undiluted vaccine contains:

Active substances:

| | |
|--|------------|
| <i>Eimeria acervulina</i> , strain 003 | 332 – 450* |
| <i>Eimeria maxima</i> , strain 013 | 196 – 265* |
| <i>Eimeria mitis</i> , strain 006 | 293 – 397* |
| <i>Eimeria praecox</i> , strain 007 | 293 – 397* |
| <i>Eimeria tenella</i> , strain 004 | 276 – 374* |

* Number of sporulated oocysts derived from precocious attenuated lines of coccidia, according to *in vitro* procedures of the manufacturer at the time of blending.

Adjuvants:

Montanide IMS
Light mineral oil

Excipients:

| Qualitative composition of excipients and other constituents |
|---|
| EVANT (suspension) |
| Potassium chloride |
| Disodium phosphate dodecahydrate |
| Potassium dihydrogen phosphate |
| Sodium chloride |
| Polysorbate 80 |
| Purified water |
| HIPRAMUNE T (solvent) |
| Brilliant blue (E 133) |
| Red AC (E 129) |
| Vanillin |
| Montanide IMS |
| HIPRACELL (solvent) |
| Brilliant blue (E 133) |
| Red AC (E 129) |
| Vanillin |
| Light mineral oil |
| Polysorbate 80 |
| Sorbitan mono-oleato |
| Potassium chloride |
| Disodium phosphate dodecahydrate |
| Potassium dihydrogen phosphate |
| Sodium chloride |
| Water for injections |

Suspension: white turbid suspension.

Solvent: dark brownish solution.

3. CLINICAL INFORMATION

3.1 Target species

Chickens.

3.2 Indications for use for each target species

For the active immunisation of chicks from 1 day of age to reduce intestinal lesions and oocysts output associated with coccidiosis caused by *Eimeria acervulina*, *Eimeria maxima*, *Eimeria mitis*, *Eimeria praecox* and *Eimeria tenella* and to reduce clinical signs (diarrhoea) associated with *Eimeria acervulina*, *Eimeria maxima* and *Eimeria tenella*.

Onset of immunity: 2 weeks post-vaccination.

Duration of immunity: 9 weeks post-vaccination in an environment that permits oocysts recycling.

3.3 Contraindications

None.

3.4 Special warnings

The vaccine will not protect species other than chickens against coccidiosis and is only effective against the *Eimeria* species indicated. This product is intended for the vaccination of short-lived chickens only.

No data is available on protection of longer-lived birds such as future layers/breeders.

Vaccinate healthy animals only.

3.5 Special precautions for use

Special precautions for safe use in the target species:

Chickens must be strictly floor-reared in the first 3 weeks after vaccination.

In order to reduce field infections, it is recommended that all litter should be removed and facilities and related equipment in contact with vaccinated chickens should be cleaned between production cycles.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Wash and disinfect hands and equipment after use.

Special precautions for the protection of the environment:

Not applicable.

3.6 Adverse events

Chickens:

None.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Laying birds:

The safety of the veterinary medicinal product has not been established during lay. Do not use in birds in lay or in breeding birds, or within 4 weeks before the start of the laying period.

3.8 Interaction with other medicinal products and other forms of interaction

No information is available on the safety and efficacy of this immunological veterinary medicinal product when used with any other veterinary medicinal product. A decision to use this immunological veterinary medicinal product before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

No anticoccidial substances or other agents having anticoccidial activity *via* feed or water should be used for at least 3 weeks following the vaccination of chickens with this product otherwise the correct replication of the vaccine oocysts, and consequently the development of a solid immunity, could be hindered. Additionally, the duration of immunity depends on an environment that permits recycling of oocysts, therefore a decision to use any anticoccidial substances in the period after 3 weeks post-vaccination should be made taking into account the potential negative impact on the duration of immunity of this product.

3.9 Administration routes and dosage

Oral use.

The method of administration is by coarse spray.

Vaccination schedule:

One dose of vaccine (0.007 ml) from 1 day of age.

Administration:

The method of administration is by coarse spray by using a suitable device (volume delivered: 28 ml/100 chicks, droplet size: 200 – 250 µm and working pressure: 1.5 to 3 bars).

Before starting to prepare the spray solution, ensure there is a clean container with sufficient capacity for preparing the diluted vaccine suspension available. Dilute the vaccine with the relevant volumes of the solvent (HIPRAMUNE T or HIPRACELL) and water, as shown in the following table:

| Doses | Water | Vaccine | Solvent | Total |
|--------|----------|---------|---------|----------|
| 1 000 | 223 ml | 7 ml | 50 ml | 280 ml |
| 5 000 | 1 115 ml | 35 ml | 250 ml | 1 400 ml |
| 10 000 | 2 230 ml | 70 ml | 500 ml | 2 800 ml |

Shake the solvent (HIPRAMUNE T or HIPRACELL) vial. Dilute the content of the vial with clean, room temperature water into an appropriate container.

Shake the vaccine (EVANT) vial and dilute the contents of it into the solvent and water solution. A purplish suspension is obtained after dilution.

Fill the reservoir of the spraying device with all the vaccine suspension prepared.

Maintain the diluted vaccine suspension in continuous homogenisation by using a magnetic stirrer while the vaccine is being administered via coarse spray to the chicks.
To improve the uniformity of the vaccination, maintain the chicks inside the transportation box for at least 1 hour in order to let them ingest all the vaccine droplets.
After this time, place the chicks carefully into the litter and continue with regular management practices.

The device should be cleaned after each use. See the manufacturer's instructions to ensure proper disinfection and maintenance of the device.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

After the administration of a severe overdose (10-fold), mild, transient, clinical signs of coccidiosis were commonly observed without any consequences on the final performance.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Zero days.

4. IMMUNOLOGICAL INFORMATION

4.1 ATCvet code: QI01AN01.

To stimulate active immunity against coccidiosis caused by *Eimeria acervulina*, *Eimeria maxima*, *Eimeria mitis*, *Eimeria praecox* and *Eimeria tenella*.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

Do not mix with any other veterinary medicinal product, except the solvent supplied for use with the veterinary medicinal product.

5.2 Shelf life

EVANT:

Shelf life of the veterinary medicinal product as packaged for sale: 10 months.

Shelf life after dilution according to directions: 10 hours.

HIPRAMUNE T (solvent):

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

HIPRACELL (solvent):

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

5.3 Special precautions for storage

Store and transport refrigerated (2 °C – 8 °C).
Do not freeze.

5.4 Nature and composition of immediate packaging

EVANT:

Type I colourless glass vials containing 7 ml, 35 ml or 70 ml of suspension (1 000, 5 000 and 10 000 doses) closed with type I polymeric elastomer closures and aluminium caps.

HIPRAMUNE T and HIPRACELL (solvents):

Polypropylene vials containing 50 ml, 250 ml or 500 ml of solvent closed with type I polymeric elastomer closures and aluminium caps.

Pack sizes:

Cardboard box with one vial of EVANT containing 7 ml (1 000 doses) and one vial of HIPRAMUNE T containing 50 ml.

Cardboard box with one vial of EVANT containing 35 ml (5 000 doses) and one vial of HIPRAMUNE T containing 250 ml.

Cardboard box with one vial of EVANT containing 70 ml (10 000 doses) and one vial of HIPRAMUNE T containing 500 ml.

Cardboard box with one vial of EVANT containing 7 ml (1 000 doses) and one vial of HIPRACELL containing 50 ml.

Cardboard box with one vial of EVANT containing 35 ml (5 000 doses) and one vial of HIPRACELL containing 250 ml.

Cardboard box with one vial of EVANT containing 70 ml (10 000 doses) and one vial of HIPRACELL containing 500 ml.

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

LABORATORIOS HIPRA, S.A.

7. MARKETING AUTHORISATION NUMBER(S)

EU/2/18/233/001-006

8. DATE OF FIRST AUTHORISATION

Date of first authorisation: 05/02/2019.

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

{DD/MM/YYYY}

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

ANNEX II

OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

None

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Cardboard boxes

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

EVANT suspension and solvent for oral spray for chickens

2. STATEMENT OF ACTIVE SUBSTANCES

Each dose (0.007 ml) of undiluted vaccine contains the following number of sporulated oocysts:

| | |
|--|-----------|
| <i>Eimeria acervulina</i> , strain 003 | 332 – 450 |
| <i>Eimeria maxima</i> , strain 013 | 196 – 265 |
| <i>Eimeria mitis</i> , strain 006 | 293 – 397 |
| <i>Eimeria praecox</i> , strain 007 | 293 – 397 |
| <i>Eimeria tenella</i> , strain 004 | 276 – 374 |

3. PACKAGE SIZE

One vial of 7 ml (1 000 doses) of EVANT and one vial of 50 ml of HIPRAMUNE T (solvent).
One vial of 35 ml (5 000 doses) of EVANT and one vial of 250 ml of HIPRAMUNE T (solvent).
One vial of 70 ml (10 000 doses) of EVANT and one vial of 500 ml of HIPRAMUNE T (solvent).
One vial of 7 ml (1 000 doses) of EVANT and one vial of 50 ml of HIPRACELL (solvent).
One vial of 35 ml (5 000 doses) of EVANT and one vial of 250 ml of HIPRACELL (solvent).
One vial of 70 ml (10 000 doses) of EVANT and one vial of 500 ml of HIPRACELL (solvent).

4. TARGET SPECIES

Chickens.

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Oral use.
Coarse spray.

7. WITHDRAWAL PERIODS

Withdrawal period: Zero days.

8. EXPIRY DATE

Exp {mm/yyyy}

Once diluted use within 10 hours.

9. SPECIAL STORAGE PRECAUTIONS

Store and transport refrigerated.

Do not freeze.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

LABORATORIOS HIPRA, S.A.

14. MARKETING AUTHORISATION NUMBERS

EU/2/18/233/001 (1 000 doses)

EU/2/18/233/002 (5 000 doses)

EU/2/18/233/003 (10 000 doses)

EU/2/18/233/004 (1 000 doses)

EU/2/18/233/005 (5 000 doses)

EU/2/18/233/006 (10 000 doses)

15. BATCH NUMBER

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

Vaccine vial of 1 000 or 5 000 doses

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

EVANT

2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

Each dose (0.007 ml) of undiluted vaccine contains the following number of sporulated oocysts:

| | |
|--|-----------|
| <i>Eimeria acervulina</i> , strain 003 | 332 – 450 |
| <i>Eimeria maxima</i> , strain 013 | 196 – 265 |
| <i>Eimeria mitis</i> , strain 006 | 293 – 397 |
| <i>Eimeria praecox</i> , strain 007 | 293 – 397 |
| <i>Eimeria tenella</i> , strain 004 | 276 – 374 |

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

Once diluted use within 10 hours.

5. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

1 000 doses
5 000 doses

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

Vaccine vial of 10 000 doses

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

EVANT suspension and solvent for oral spray for chickens

2. STATEMENT OF ACTIVE SUBSTANCES

Each dose (0.007 ml) of undiluted vaccine contains the following number of sporulated oocysts:

| | |
|--|-----------|
| <i>Eimeria acervulina</i> , strain 003 | 332 – 450 |
| <i>Eimeria maxima</i> , strain 013 | 196 – 265 |
| <i>Eimeria mitis</i> , strain 006 | 293 – 397 |
| <i>Eimeria praecox</i> , strain 007 | 293 – 397 |
| <i>Eimeria tenella</i> , strain 004 | 276 – 374 |

3. TARGET SPECIES

Chickens.

4. ROUTES OF ADMINISTRATION

Oral use.

Coarse spray.

To be mixed with the HIPRAMUNE T or HIPRACELL (solvent).

Read the package leaflet before use.

5. WITHDRAWAL PERIODS

Withdrawal period: Zero days.

6. EXPIRY DATE

Exp. {mm/yyyy}

Once diluted use within 10 hours.

7. SPECIAL STORAGE PRECAUTIONS

Store and transport refrigerated.

Do not freeze.

8. NAME OF THE MARKETING AUTHORISATION HOLDER

LABORATORIOS HIPRA, S.A.

9. BATCH NUMBER

Lot {number}

10. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

10 000 doses

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING (LABEL) OF THE SOLVENT

Solvent vial of 50 ml, 250 ml or 500 ml: Hipramune T

1. NAME OF THE SOLVENT

HIPRAMUNE T solvent for oral spray for chickens

2. TARGET SPECIES

Chickens.

3. ROUTES OF ADMINISTRATION

Read the package leaflet before use.

4. EXPIRY DATE

Exp. {mm/yyyy}

5. SPECIAL STORAGE PRECAUTIONS

Store and transport refrigerated.
Do not freeze.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

LABORATORIOS HIPRA, S.A.

7. BATCH NUMBER

Lot {number}

8. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

50 ml
250 ml
500 ml

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING (LABEL) OF THE SOLVENT

Solvent vial of 50 ml, 250 ml or 500 ml: Hipracell

1. NAME OF THE SOLVENT

HIPRACELL solvent for oral spray for chickens

2. TARGET SPECIES

Chickens.

3. ROUTES OF ADMINISTRATION

Oral use. Coarse spray.
Read the package leaflet before use.

4. EXPIRY DATE

Exp. {mm/yyyy}

5. SPECIAL STORAGE PRECAUTIONS

Store and transport refrigerated.
Do not freeze.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

LABORATORIOS HIPRA, S.A.

7. BATCH NUMBER

Lot {number}

8. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

50 ml
250 ml
500 ml

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

EVANT suspension and solvent for oral spray for chickens

2. Composition

Active substances:

Each dose (0.007 ml) of undiluted vaccine contains:

Eimeria acervulina, strain 003332 – 450*

Eimeria maxima, strain 013196 – 265*

Eimeria mitis, strain 006293 – 397*

Eimeria praecox, strain 007293 – 397*

Eimeria tenella, strain 004276 – 374*

* Number of sporulated oocysts derived from precocious attenuated lines of coccidia, according to *in vitro* procedures of the manufacturer at the time of blending.

Suspension: white turbid suspension.

Solvent: dark brownish solution.

3. Target species

Chickens.

4. Indications for use

For the active immunisation of chicks from 1 day of age to reduce intestinal lesions and oocysts output associated with coccidiosis caused by *Eimeria acervulina*, *Eimeria maxima*, *Eimeria mitis*, *Eimeria praecox* and *Eimeria tenella* and to reduce clinical signs (diarrhoea) associated with *Eimeria acervulina*, *Eimeria maxima* and *Eimeria tenella*.

Onset of immunity: 2 weeks post-vaccination.

Duration of immunity: 9 weeks post-vaccination in an environment that permits oocysts recycling.

5. Contraindications

None.

6. Special warnings

Special warnings:

The vaccine will not protect species other than chickens against coccidiosis and is only effective against the *Eimeria* species indicated. This product is intended for the vaccination of short-lived chickens only. No data is available on protection of longer-lived birds such as future layers/breeders.

Special precautions for safe use in the target species:

Vaccinate healthy chickens only.

Chickens must be strictly floor-reared in the first 3 weeks after vaccination. In order to reduce field infections, it is recommended that all litter should be removed and facilities and related equipment in contact with vaccinated chickens should be cleaned between production cycles.

Laying birds:

The safety of the veterinary medicinal product has not been established during lay. Do not use in birds in lay or in breeding birds, or within 4 weeks before the start of the laying period.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:
Wash and disinfect hands and equipment after use.

Interaction with other medicinal products and other forms of interaction:

No information is available on the safety and efficacy of this immunological veterinary medicinal product when used with any other veterinary medicinal product. A decision to use this immunological veterinary medicinal product before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

No anticoccidial substances or other agents having anticoccidial activity *via* feed or water should be used for at least 3 weeks following the vaccination of chickens with this product otherwise the correct replication of the vaccine oocysts, and consequently the development of a solid immunity, could be hindered. Additionally, the duration of immunity depends on an environment that permits recycling of oocysts, therefore a decision to use any anticoccidial substances in the period after 3 weeks post-vaccination should be made taking into account the potential negative impact on the duration of immunity of this product.

Overdose:

After the administration of a severe overdose (10 - fold), mild, transient, clinical signs of coccidiosis were commonly observed without any consequences on the final performance.

Major incompatibilities:

Do not mix with any other veterinary medicinal product, except the solvent supplied for use with the veterinary medicinal product.

7. Adverse events

Chickens:

None.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: {national system details}.

8. Dosage for each species, routes and method of administration

One dose of vaccine (0.007 ml) from 1 day of age.

Oral use.

The method of administration is by coarse spray.

9. Advice on correct administration

The method of administration is by coarse spray by using a suitable device (volume delivered: 28 ml/100 chicks, droplet size: 200 – 250 µm and working pressure: 1.5 to 3 bars). Before starting to prepare the spray solution, ensure there is a clean container with sufficient capacity for preparing the diluted vaccine suspension available. Dilute the vaccine with the relevant volumes of the solvent (HIPRAMUNE T or HIPRACELL) and water, as shown in the following table:

| DOSES | WATER | VACCINE | Solvent | TOTAL |
|--------|----------|---------|---------|----------|
| 1 000 | 223 ml | 7 ml | 50 ml | 280 ml |
| 5 000 | 1 115 ml | 35 ml | 250 ml | 1 400 ml |
| 10 000 | 2 230 ml | 70 ml | 500 ml | 2 800 ml |

Shake the solvent vial (HIPRAMUNE T or HIPRACELL). Dilute the contents of the vial with clean, room temperature water into an appropriate container.

Shake the vaccine (EVANT) vial and dilute the contents of it into the solvent and water solution. A purplish suspension is obtained after dilution.

Fill the reservoir of the spraying device with all the vaccine suspension prepared.

Maintain the diluted vaccine suspension in continuous homogenisation by using a magnetic stirrer while the vaccine is being administered via coarse spray to the chicks.

To improve the uniformity of the vaccination, maintain the chicks inside the transportation box for at least 1 hour in order to let them ingest all the vaccine droplets.

After this time, place the chicks carefully into the litter and continue with regular management practices.

The device should be cleaned after each use. See the manufacturer's instructions to ensure proper disinfection and maintenance of the device.

10. Withdrawal periods

Zero days.

11. Special storage precautions

Keep out of the sight and reach of children.

Store and transport refrigerated (2 °C – 8 °C).

Do not freeze.

Shelf life after dilution according to directions: 10 hours.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and the label. The expiry date refers to the last day of that month.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack sizes

Marketing authorisation number: EU/2/18/233/001-006

Pack sizes:

Cardboard box with one vial of EVANT containing 7 ml (1 000 doses) and one vial of HIPRAMUNE T containing 50 ml.

Cardboard box with one vial of EVANT containing 35 ml (5 000 doses) and one vial of HIPRAMUNE T containing 250 ml.

Cardboard box with one vial of EVANT containing 70 ml (10 000 doses) and one vial of HIPRAMUNE T containing 500 ml.

Cardboard box with one vial of EVANT containing 7 ml (1 000 doses) and one vial of HIPRACELL containing 50 ml.

Cardboard box with one vial of EVANT containing 35 ml (5 000 doses) and one vial of HIPRACELL containing 250 ml.

Cardboard box with one vial of EVANT containing 70 ml (10 000 doses) and one vial of HIPRACELL containing 500 ml.

Not all pack sizes may be marketed.

15. Date on which the package leaflet was last revised

{DD/MM/YYYY}

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

16. Contact details

Marketing authorisation holder and manufacturer responsible for batch release and contact details to report suspected adverse reactions:

LABORATORIOS HIPRA, S.A.
Avda. la Selva 135
17170 Amer (Girona) SPAIN
[TEL:+34 972 43 06 60](tel:+34972430660)

Local representatives and contact details to report suspected adverse reactions:

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder

België/Belgique/Belgien

HIPRA BENELUX NV
Nieuwewandeling 62
9000 Gent
BELGIUM
Tel: +32 09 2964464

Република България

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ИСПАНИЯ
Тел: +34 972 43 06 60

Česká republika

HIPRA SLOVENSKO, s.r.o.
Zochova 5,
811 03 Bratislava,
SLOVENSKO
Tel: +421 02 32 335 223

Danmark

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPANIEN
Tel: +34 972 43 06 60

Deutschland

HIPRA DEUTSCHLAND GmbH
Am Wehrhahn 28-30
40211 Düsseldorf
DEUTSCHLAND
Tel: +49 211 698236 – 0

Eesti

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
HISPAANIA
Tel: +34 972 43 06 60

Ελλάδα

HIPRA ΕΛΛΑΣ Α.Ε.
Λεωφ. Αθηνών 80 & Μηριόνου 2-4,
104 41 Κολωνός - ΑΘΗΝΑ - ΕΛΛΑΣ
Τηλ: +30 210 4978660

Lietuva

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ISPANIJA
Tel: +34 972 43 06 60

Luxembourg/Luxemburg

HIPRA BENELUX NV
Nieuwewandeling 62
9000 Gent
BELGIUM
Tel: +32 09 2964464

Magyarország

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPANYOLORSZÁG
Tel: +34 972 43 06 60

Malta

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPANJA
Tel: +34 972 43 06 60

Nederland

HIPRA BENELUX NV
Nieuwewandeling 62
9000 Gent
BELGIUM
Tel: +32 09 2964464

Norge

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPANIA
Tlf: +34 972 43 06 60

Österreich

HIPRA DEUTSCHLAND GmbH
Am Wehrhahn 28-30
40211 Düsseldorf
DEUTSCHLAND
Tel: +49 211 698236 – 0

España

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ESPAÑA
Tel: +34 972 43 06 60

France

HIPRA FRANCE
7 rue Roland Garros, Batiment H
44700 - Orvault -
FRANCE
Tél: +33 02 51 80 77 91

Hrvatska

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ŠPANJOLSKA
Tel: +34 972 43 06 60

Ireland

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPAIN
Tel: +34 972 43 06 60

Ísland

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPÁNN
Sími: +34 972 43 06 60

Italia

Hipra Italia S.r.l.
Enrico Mattei, 2
25030 Coccaglio (BS)
ITALIA
Tel: +39 030 7241821

Κύπρος

LABORATORIOS HIPRA, S.A.
Λεωφόρος Λα Σέλβα 135, Αμερ
Τζιρόνα 17170
Ισπανία Τηλ: +34 972 43 06 60

Polska

HIPRA POLSKA Sp.z.o.o.
Ul. Wincentego Rzymowskiego 31
02-697 Warszawa - POLSKA
Tel: +48 22 642 33 06

Portugal

ARBUSET, Produtos Farmacêuticos e Sanitários
De Uso Animal, Lda
Portela de Mafra e Fontainha - Abrunheira
2665 – 191 Malveira - PORTUGAL
Tel: +351 219 663 450

România

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPANIA
Tel: +34 972 43 06 60

Slovenija

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ŠPANIJA
Tel: +34 972 43 06 60

Slovenská republika

HIPRA SLOVENSKO, s.r.o.
Zochova 5,
811 03 Bratislava,
SLOVENSKO
Tel: +421 02 32 335 223

Suomi/Finland

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
ESPANJA
Puh/Tel: +34 972 43 06 60

Sverige

LABORATORIOS HIPRA, S.A.
Avda. La Selva 135
17170 Amer (Girona)
SPANIEN
Tel: +34 972 43 06 60

Latvija

LABORATORIOS HIPRA, S.A.

Avda. La Selva 135

17170 Amer (Girona)

SPĀNIJA

Tel. +34 972 43 06 60

United Kingdom (Northern Ireland)

LABORATORIOS HIPRA, S.A.

Avda. La Selva 135

17170 Amer (Girona)

SPAIN

Tel: +34 972 43 06 60