



**FRENCH AGENCY FOR VETERINARY MEDICINAL PRODUCTS  
AGENCE NATIONALE DU MEDICAMENT VETERINAIRE**

8 rue Claude Bourgelat –  
Parc d'activités de la grande Marche –  
Javené – CS 70611 –  
35306 FOUGERES

**DECENTRALISED PROCEDURE**

**PUBLICLY AVAILABLE ASSESSMENT REPORT FOR A VETERINARY MEDICINAL PRODUCT**

**TYLOVECTIN 200 solution for injection for cattle, goats and pigs**

**DATE : 30/10/2018**

## MODULE 1

### PRODUCT SUMMARY

EU Procedure number	FR/V/0323/001/DC
Name, strength and pharmaceutical form	TYLOVECTIN 200 solution for injection for cattle, goats and pigs.
Applicant	Aniserve GmbH Geyerspergerstr. 27 80689 Munich Germany
Active substance(s)	Tylosin
ATC Vetcode	QJ01FA90
Target species	Cattle Pigs Goats
Indication for use	<p><u>For the treatment of specific infections (listed below)</u> caused by microorganisms susceptible to tylosin.</p> <p><u>Cattle (adult):</u> -Respiratory infections, metritis caused by Gram-positive microorganisms, mastitis caused by <i>Streptococcus spp</i>, <i>Staphylococcus spp</i> and interdigital necrobacillosis, i.e. panaritium or foot rot.</p> <p><u>Calves:</u> -Respiratory infections and necrobacillosis.</p> <p><u>Pigs:</u> -Enzootic pneumonia, haemorrhagic enteritis, erysipelas and metritis. -Arthritis caused by <i>Mycoplasma spp.</i> and <i>Staphylococcus spp.</i></p> <p>For information regarding swine dysentery see section 4.5.</p> <p><u>Goats:</u> - Respiratory infections and metritis caused by Gram-positive microorganisms - Mastitis caused by Gram-positive microorganisms or <i>Mycoplasma spp..</i></p>

**MODULE 2**

The Summary of Product Characteristics (SPC) for this product is available on the website <http://mri.medagencies.org/veterinary/>

**MODULE 3****PUBLIC ASSESSMENT REPORT**

Legal basis of original application	Generic application in accordance with Article 13(1) of Directive 2001/82/EC as amended.
Date of completion of the original decentralised procedure	26/09/2018
Concerned Member States for original procedure	BE, NL, UK

**I. SCIENTIFIC OVERVIEW**

The product is produced and controlled using validated methods and tests, which ensure the consistency of the product released on the market.

It has been shown that the product can be safely used in the target species; the reactions that may be observed are indicated in the SPC, with information on the frequency.

The product is safe for the user, the consumer of foodstuffs from treated animals and for the environment, when used as recommended. Suitable warnings and precautions are indicated in the SPC.

The efficacy claims for these products are equivalent to those of the reference product with taking also into account the decision of the European Commission in a referral on oral products containing Tylosin to be used in pigs.

The overall risk/benefit analysis is in favour of granting a marketing authorisation.

## II. QUALITY ASPECTS

### A. *Composition*

The product contains 200 000 IU/ml of tylosin and the following excipients: benzyl alcohol, propylene glycol and water for injection.

The product is packed in glass vials. The particulars of the containers and controls performed are provided and conform to the regulation.

The product is an established pharmaceutical form and its development is adequately described in accordance with the relevant European guidelines.

### B. *Method of Preparation of the Product*

The product is manufactured fully in accordance with the principles of good manufacturing practice from a licensed manufacturing site.

Process validation data on the product have been presented in accordance with the relevant European guidelines.

### C. *Control of Starting Materials*

The active substance is tylosin, an established active substance described in the European Pharmacopoeia. The active substance is manufactured in accordance with the principles of good manufacturing practice.

The active substance specification is considered adequate to control the quality of the material. Batch analytical data demonstrating compliance with this specification have been provided.

### D. *Specific Measures concerning the Prevention of the Transmission of Animal Spongiform Encephalopathies*

There are no substances within the scope of the TSE Guideline present or used in the manufacture of this product.

### E. *Control on intermediate products*

Not applicable.

### F. *Control Tests on the Finished Product*

The finished product specification controls the relevant parameters for the pharmaceutical form. The tests in the specification, and their limits, have been justified and are considered appropriate to adequately control the quality of the product.

Satisfactory validation data for the analytical methods have been provided.

Batch analytical data from the proposed production site have been provided demonstrating compliance with the specification.

#### ***G. Stability***

A re-test period for the active substance is set in the certificate of suitability issued by EDQM.

Stability data on the finished product have been provided in accordance with applicable European guidelines, demonstrating the stability of the product throughout its shelf life when stored under the approved conditions.

An in-use shelf-life as detailed on the SPC has been supported by appropriate data.

#### ***H. Genetically Modified Organisms***

Not applicable.

#### ***J. Other Information***

Not applicable.

### **III. SAFETY AND RESIDUES ASSESSMENT**

#### ***III.A Safety Testing***

##### ***Pharmacological Studies***

Based on exemption 7.1. b) of the “Guidelines on the conduct of bioequivalence studies for veterinary medicinal products” (EMA/CVMP/016/00-Rev.2), it is accepted that the test product is bioequivalent to the reference product TYLAN 200 marketed by LILLY FRANCE and authorized in France since 15/12/1980. As this is a generic application according to Article 13, and bioequivalence with a reference product has been demonstrated, results of pharmacological and toxicological tests are not required.

The pharmacological aspects of this product are identical to those of the reference product.

##### ***Toxicological Studies***

The toxicological aspects of this product are identical to those of the reference product.

##### ***User Safety***

The applicant has provided a user safety assessment in compliance with the relevant guideline.

Warnings and precautions as listed on the product literature are adequate to ensure safety of the product to users.

##### ***Ecotoxicity***

A phase I and a phase II Environmental Risk Assessment (ERA) has been performed according to VICH Phase I Guideline CVMP/VICH/592/98-final, VICH Phase II Guideline CVMP/VICH/790/03-final and the supporting CVMP Guideline EMEA/CVMP/ERA/ 418282/2005-rev1.

**Phase I:**

A Phase II ERA is required as the Phase I assessment showed that the initial predicted environmental concentration in soil (PECsoil initial = 174.9 µg/kg) is greater to 100 µg/kg and no mitigations exist that alter the PECsoil.

**Phase II:**

A Phase II data set was provided according to the requirements of the CVMP/VICH guideline GL38 and the CVMP guideline on the Environmental Impact Assessment for Veterinary Medicinal Products in support of the VICH guidelines GL6 and GL38 (EMEA/CVMP/ERA/418282/2005-Rev.1). The data were considered to be complete and acceptable.

<b>Physical-chemical properties</b>			
<b>Study type</b>	<b>Test protocol</b>	<b>Result</b>	<b>Remarks</b>
Water solubility	OECD 105	5 g/L	
Dissociation constants in water pKa	OECD 112	pKa = 7.73	
n-Octanol/Water Partition Coefficient logP <sub>ow</sub>	OECD 107 and 117	logK <sub>ow</sub> = 0.36 at pH 5, 1.18 at pH 7 1.36 at pH 9	

<b>Environmental fate</b>			
Soil Adsorption/Desorption	OECD 106	Koc = 3820 l/kg	
Aerobic and Anaerobic Transformation in Soil	OECD 307	DT50 (geometric mean) = 16.1 days	

<b>Effect studies</b>					
<b>Study type</b>	<b>Test protocol</b>	<b>Endpoint</b>	<b>Result</b>	<b>Unit</b>	<b>Remarks</b>
Algae and or cyanobacteria, growth inhibition test/species	OECD 201	EC50	193	µg/l	

<i>Daphnia</i> sp. immobilisation	OECD 202	EC50	680	mg/l	
Fish, acute toxicity/species	OECD 203	LC50	>46	mg/l	
Soil microorganisms: Nitrogen transformation test (28 days)	OECD 216	% effect	66.3		Trigger value: 25% deviation from the control
Soil micro-organisms: nitrogen transformation test (100 days)	OECD 216	% effect	28.5		At concentration higher than PEC x 10
Terrestrial Plants, growth test	OECD 208	EC50	124.1	mg/kg	Allium cepa Avena sativa Cucumis sativus Lycopersicon lycopersicum Pisum sativum Raphanus sativus
Earthworm/ <i>Enchytraeidae</i> reproduction	OECD 220/222	EC10	149	mg/kg	No OECD compliance

### ***Risk characterisation***

The Predicted Environmental Concentration (PEC) for each compartment was calculated in accordance with VICH guideline GL6 and the CVMP guideline on the Environmental Impact Assessment for Veterinary Medicinal Products in support of the VICH guidelines GL6 and GL38 (EMEA/CVMP/ERA/418282/2005-Rev.1)

Using the assessment factors (AF) in these VICH guidelines, predicted no effect concentrations (PNEC) were calculated and compared with the PEC values. This results in a risk quotient (RQ) for each compartment as follows:

Compartment	PNEC	PEC	RQ
surface water	46	0.31	0.0067
groundwater		0.93 FOCUS : < 0.001	
soil microorganisms: Nitrogen	28.5 %	NA	NA

transformation test	difference in N transformation at high concentration		
soil	1241	174.9	0.14

The risk characterisation resulted in risk quotients (RQs) below 1 for the surface water and soil compartments indicating that the product will not pose a risk to those compartments when used as recommended.

#### PBT assessment

PBT-assessment			
Parameter	Result relevant for conclusion		Conclusion
Bioaccumulation	BCF		not B
Persistence	DT <sub>50</sub> , compartment, 12 °C	16.1 days	not P
Toxicity	NOEC or CMR	46 mg/L	not T
<b>PBT-statement :</b>	The compound is not considered as PBT nor vPvB		

#### III.B Residues documentation

##### Residue Studies

No residue depletion studies were required, based on exemption 7.1. b) of the "Guidelines on the conduct of bioequivalence studies for veterinary medicinal products" (EMA/CVMP/016/00-Rev.2) and in accordance with the CVMP guideline "Approach towards harmonisation of withdrawal periods" (EMEA/CVMP/036/95).

##### MRLs

The active substance tylosin is listed in table 1 of Council Regulation 37/2010. The marker substance is Tylosin A.

MRLs are listed below:

Marker residue	Animal Species	MRL	Target Tissues	Other Provisions	Therapeutic Classification	Regulation
Tylosin A	All food producing species	100 µg/kg 100 µg/kg 100 µg/kg 100 µg/kg 50 µg/kg 200 µg/kg	Muscle Fat Liver Kidneys Milk Eggs	For fin fish the muscle MRL relates to « muscle and skin in natural proportions ». MRLs for fat, liver and kidney do not apply for fish. For porcine and poultry species,	Anti-infectious agents/ Antibiotics	37/2010 of 22.12.2009

				the fat MRL relates to "skin and fat in natural proportions".		
--	--	--	--	---	--	--

The MRL status of excipients is indicated in the following table:

Excipient	MRL status
Benzyl alcohol	Table1, no MRL required
Propylene glycol	Table1, no MRL required
Water for injection	Out of scope

### ***Withdrawal Periods***

The same withdrawal periods as the reference product are accepted:

Cattle:

Meat – 28 days

Milk – 108 hours

Goat:

Meat – 42 days

Milk – 108 hours

Pigs:

Meat – 14 days

## **IV. CLINICAL ASSESSMENT (EFFICACY)**

As this is a generic application according to Article 13, and bioequivalence with a reference product has been demonstrated based on exemption 7.1. b) of the "Guidelines on the conduct of bioequivalence studies for veterinary medicinal products" (EMA/CVMP/016/00-Rev.2), efficacy studies are not required.

### ***IV.A Pre-Clinical Studies***

#### ***Tolerance in the Target Species of Animals***

The product literature accurately reflects the type and incidence of adverse effects which might be expected.

#### ***Resistance***

The bibliography regarding the susceptibility or resistance to tylosin for pathogens but also for commensal bacteria, provided in a context of prudent use of antimicrobials and responsible attitude of the applicant, is difficult to interpret in the absence of recognised break points for tylosin.

Adequate warnings and precautions appear on the product literature.

#### ***IV.B Clinical Studies***

The efficacy claims for this product are equivalent to those of the reference product and take also into account the decision taken by the European Commission in a referral on oral products containing tylosin to be used in pigs.

### **V. OVERALL CONCLUSION AND BENEFIT– RISK ASSESSMENT**

The data submitted in the dossier demonstrate that when the product is used in accordance with the Summary of Product Characteristics, the risk benefit profile for the target species is favourable and the quality and safety of the product for humans and the environment are acceptable.